

11/28/00



J0970 U.S. PTO

11-29-00

PATENT



## UTILITY PATENT APPLICATION TRANSMITTAL

(only for new nonprovisional applications under 37 CFR §1.53(b))

Assistant Commissioner of Patents

Box Patent Application

Washington, D.C. 20231

Sir:

Transmitted herewith for filing is the patent application  
under 37 CFR 1.53(b) of:

AVENTIS DOCKET NO.: **A3536A-US**INVENTORS: **Jayyosi et al**

TITLE: **THERAPEUTIC USES OF TRI-ARYL ACID  
DERIVATIVES**

I hereby certify that this correspondence is being  
deposited with the United States Postal Service as  
Express Mail in an envelope addressed to Assistant  
Commissioner for Patents, Washington, D.C. 20231,  
on

Date of Deposit

Signature

Express Mail No.

## APPLICATION ELEMENTS

1. ☒ \*Fee Transmittal Form (e.g., PTO/SB/17)  
(Submit an original and a duplicate for fee processing)
2. ☒ Specification  
(preferred arrangement set forth below)
- Descriptive title of the Invention
  - Cross References to Related Applications
  - Statement Regarding Fed sponsored R & D
  - Reference to Microfiche Appendix
  - Background of the Invention
  - Brief Summary of the Invention
  - Brief Description of the Drawings (if filed)
  - Detailed Description
  - Claim(s)
  - Abstract of the Disclosure

Total Pages

250

3. ☐ Drawing(s) (35 U.S.C. 113)

Total Sheets

0

4. Oath or Declaration

Total Pages

3

- a. ☒ Newly executed (original or copy) (**unsigned**)

- b. ☐ Copy from a prior application (37 CFR § 1.63(d))  
(for continuation/divisional with Box 16 completed)

- i. ☐ **DELETION OF INVENTOR(S)**  
Signed statement attached deleting inventor(s) named in the prior  
application, see 37 CFR §§ 1.63(d)(2) and 1.33(b).

5. ☐ Microfiche Computer Program (Appendix)

6. Nucleotide and/or Amino Acid Sequence Submission  
(if applicable, all necessary)

- a. ☐ Computer Readable Copy
- b. ☐ Paper Copy (identical to computer copy)
- c. ☐ Statement verifying identity of above copies

☒ If a **CONTINUING Application**, check appropriate box, and supply the requisite information:

☒ Continuation ☐ Divisional ☐ Continuation-in-part  
of prior application no: PCT/US00/11490, filed 28 April 2000.

Examiner \_\_\_\_\_, Group/Art Unit \_\_\_\_\_.

(The cross reference has been/will be inserted on page one of the specification).

☒ This application claims the benefit of U.S. Provisional Application No. 60/131,454,  
filed 28 April 1999. (The cross reference has been/will be inserted on page one of the specification).

☐ **Incorporation By Reference** (useable if filing a continuation/divisional and a copy of the declaration from the prior application is enclosed.)

The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

### ACCOMPANYING APPLICATION PARTS


7. ☐ Assignment Papers (cover sheet & document(s))
8. ☐ 37 CFR §3.73(b) Statement ☐ Power of Attorney  
(when there is an assignee)
9. ☐ English Translation Document (if applicable)
10. ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations
11. ☐ Preliminary Amendment
12. ☒ Return Receipt Postcard (MPEP 503)  
(Should be specifically itemized)
13. ☐ \*Small Entity Statement(s) ☐ Statement filed in prior application,  
(PTO/SB/09-12) Status still proper and desired
14. ☐ Certified Copy of Priority Document(s)  
(if foreign priority is claimed)
15. ☒ Other: Fee Sheet  
EFS Form

### CORRESPONDENCE ADDRESS

☒ Customer Number or Bar Code Label

005487

(Insert Customer No. or Attach bar code label here)

  
Irving Newman  
Attorney/Agent for Applicant  
Reg. No. 22,638

Aventis Pharmaceuticals Inc.  
Patent Department  
Route #202-206 / P.O. Box 6800  
Mail Code: EMC-G1  
Bridgewater, New Jersey 08807-0800  
Telephone (908) 231-2785  
Telefax (908) 231-2626

Aventis Docket No. A3535A-US

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

**FEE TRANSMITTAL**  
**for FY 2000**

Patent fees are subject to annual revision.

Small Entity payments must be supported by a small entity statement,  
otherwise large entity fees must be paid. See Forms PT/SB/09-12.  
See 37 C.F.R. §§ 1.27 and 1.28.**TOTAL AMOUNT OF PAYMENT (\$)** 2,334.00**Complete if Known**

Application Number	
Filing Date	
First Named Inventor	Jayyosi et al
Examiner Name	
Group / Art Unit	
Attorney Docket No.	A3536A-US

**METHOD OF PAYMENT (check one)**1. ☒ The Commissioner is hereby authorized to charge indicated fees and credit any overpayments to:Deposit Account Number **18-1982**Deposit Account Name **Aventis Pharmaceuticals Products Inc.**☒ Charge Any Additional Fee Required  
Under 37 C.F.R. §§ 1.16 and 1.172. ☐ Payment Enclosed:☐ Check ☐ Money Order ☐ Other**FEE CALCULATION****1. BASIC FILING FEE**

Large Fee Code	Entity Fee (\$)	Small Fee Code	Entity Fee (\$)	Fee Description	Fee Paid
101	690	201	345	Utility filing fee	710
106	310	206	155	Design filing fee	
107	480	207	240	Plant filing fee	
108	690	208	345	Reissue filing fee	
109	150	214	75	Provisional filing fee	

**SUBTOTAL (1) (\$)** 710**2. EXTRA CLAIM FEES**

Extra Claims				Fee from below	Fee Paid
Total Claims	88	-20**=	68	x 18	= 1224
Independent Claims	8	-3**=	5	x 80	= 400
Multiple Dependent					

\*\*or number previously paid, if greater; For Reissues, see below

Large Fee Code	Entity Fee (\$)	Small Fee Code	Entity Fee (\$)	Fee Description
103	18	203	9	Claims in excess of 20
102	78	202	39	Independent claims in excess of 3
104	260	204	130	Multiple dependent claim, if not paid
109	78	209	39	**Reissue independent claims over original patent
110	18	210	9	**Reissue claims in excess of 20 and over original patent

**SUBTOTAL (2) (\$)** 1624**FEE CALCULATION (continued)****3. ADDITIONAL FEES**

Large Fee Code	Entity Fee (\$)	Small Fee Code	Entity Fee (\$)	Fee Description	Fee Paid
105	130	205	65	Surcharge - late filing fee or oath	
127	50	227	25	Surcharge - late provisional filing fee or cover sheet	
139	130	139	130	Non-English Specification	
147	2,520	147	2,520	For filing a request for reexamination	
112	920*	112	920*	Requesting publication of SIR prior to Examiner action	
113	1,840*	113	1,840*	Requesting publication of SIR after Examiner action	
115	110	215	55	Extension for reply within first month	
116	380	216	190	Extension for reply within second month	
117	870	217	435	Extension for reply within third month	
118	1,360	218	680	Extension for reply within fourth month	
128	1,850	228	925	Extension for reply within fifth month	
119	300	219	150	Notice of Appeal	
120	300	220	150	Filing a brief in support of an appeal	
121	260	221	130	Request for oral hearing	
138	1,510	138	1,510	Petition to institute a public use proceeding	
140	110	240	55	Petition to revive - unavoidable	
141	1,210	241	605	Petition to revive - unintentional	
142	1,210	242	605	Utility issue fee (or reissue)	
143	430	243	215	Design issue fee	
144	580	244	290	Plant issue fee	
122	130	122	130	Petitions to the Commissioner	
123	50	123	50	Petitions related to provisional applications	
126	240	126	240	Submission of Information Disclosure Stmt	
581	40	581	40	Recording each patent assignment per property (times number of properties)	
146	690	246	345	Filing a submission after final rejection (37 CFR § 1.129(a))	
149	690	249	345	For each additional invention to be examined (37 CFR § 1.129(b))	

Other fee (specify) \_\_\_\_\_

Other fee (specify) \_\_\_\_\_

\*Reduced by Basic Filing Fee Paid

**SUBTOTAL (3)** 0**SUBMITTED BY**Name (Print/Type) **Irving Newman, Reg. No. 22,638**

Registration No. (Attorney/Agent)

22,638

**Complete (if applicable)**

Telephone (908) 231-2785

Signature

Date

11/15/00

**WARNING:**

Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

INVENTOR INFORMATION

Inventor One Given Name:: Zaid  
Family Name:: Joyyosi  
Postal Address Line One:: 108 Cherrywood Court  
City:: Collegeville  
State or Province:: PA  
Country:: US  
Postal or Zip Code:: 19426  
City of Residence:: Collegeville  
State or Province of Residence:: PA  
Country of Residence:: US  
Citizenship Country:: FR  
Inventor Two Given Name:: Gerard M  
Family Name:: McGeehan  
Postal Address Line One:: 1711 Spring House Road  
City:: Chester Springs  
State or Province:: PA  
Country:: US  
Postal or Zip Code:: 19425  
City of Residence:: Chester Springs  
State or Province of Residence:: PA  
Country of Residence:: US  
Citizenship Country:: US  
Inventor Three Given Name:: Michael F  
Family Name:: Kelley  
Postal Address Line One:: 1086 Heartsease Drive  
City:: West Chester  
State or Province:: PA  
Country:: US  
Postal or Zip Code:: 19382  
City of Residence:: West Chester  
State or Province of Residence:: PA  
Country of Residence:: US  
Citizenship Country:: US  
Inventor Four Given Name:: Richard F  
Family Name:: Labaudiniere  
Postal Address Line One:: 220 Richard Way  
City:: Collegeville  
State or Province:: PA  
Country:: US  
Postal or Zip Code:: 19426  
City of Residence:: Collegeville  
State or Province of Residence:: PA  
Country of Residence:: US  
Citizenship Country:: FR  
Inventor Five Given Name:: Litao  
Family Name:: Zhang  
Postal Address Line One:: 456 Shakespere Drive  
City:: Collegeville

State or Province:: PA  
Country:: US  
Postal or Zip Code:: 19426  
City of Residence:: Collegeville  
State or Province of Residence:: PA  
Country of Residence:: US  
Citizenship Country:: US  
Inventor Six Given Name:: Robert D  
Family Name:: Groneberg  
Postal Address Line One:: 4173 Ironbridge Drive  
City:: Collegeville  
State or Province:: PA  
Country:: US  
Postal or Zip Code:: 19426  
City of Residence:: Collegeville  
State or Province of Residence:: PA  
Country of Residence:: US  
Citizenship Country:: US  
Inventor Seven Given Name:: Daniel G  
Family Name:: McGarry  
Postal Address Line One:: 3000 Valley Forge Circle, Apt. 1148  
City:: King of Prussia  
State or Province:: PA  
Country:: US  
Postal or Zip Code:: 19406  
City of Residence:: King of Prussia  
State or Province of Residence:: PA  
Country of Residence:: US  
Citizenship Country:: GB  
Inventor Eight Given Name:: Thomas J  
Family Name:: Caulfield  
Postal Address Line One:: 362 Vista Drive  
City:: Phoenixville  
State or Province:: PA  
Country:: US  
Postal or Zip Code:: 19460  
City of Residence:: Phoenixville  
State or Province of Residence:: PA  
Country of Residence:: US  
Citizenship Country:: US  
Inventor Nine Given Name:: Anne  
Family Name:: Minnich  
Postal Address Line One:: 2107 Goodwin Lane  
City:: Montgomeryville  
State or Province:: PA  
Country:: US  
Postal or Zip Code:: 19454  
City of Residence:: Montgomeryville  
State or Province of Residence:: PA  
Country of Residence:: US

Citizenship Country:: US  
Inventor Ten Given Name:: Mark  
Family Name:: Bobko  
Postal Address Line One:: 526 Summerville Drive  
City:: Exton  
State or Province:: PA  
Country:: US  
Postal or Zip Code:: 19341  
City of Residence:: Exton  
State or Province of Residence:: PA  
Country of Residence:: US  
Citizenship Country:: US  
Inventor Eleven Given Name:: Robert  
Family Name:: Morris  
Postal Address Line One:: 125 Conestoga Road  
City:: Wayne  
State or Province:: PA  
Country:: US  
Postal or Zip Code:: 19087  
City of Residence:: Wayne  
State or Province of Residence:: PA  
Country of Residence:: US  
Citizenship Country:: US

CORRESPONDENCE INFORMATION

Correspondence Customer Number:: 005487  
Name Line One:: Irving Newman  
Address Line One:: Route 202-206 EMC-G1  
City:: Bridgewater  
State or Province:: NJ  
Country:: US  
Postal or Zip Code:: 08807  
Fax One:: (908)231-2626  
Electronic Mail One:: Irving.Newman@Aventis.com

APPLICATION INFORMATION

Title Line One:: THERAPEUTIC USES OF TRI-ARYL DERIVATIVES  
Total Drawing Sheets:: 0  
Formal Drawings?:: No  
Application Type:: Utility  
Docket Number:: A3536A-US  
Secrecy Order in Parent Appl.?:: No

CONTINUITY INFORMATION

This application is a:: CONTINUATION OF  
> Application One:: PCT/US00/11490

SCANNED # 12

Filing Date:: 04-28-2000

This application is a:: NON PROV. OF PROVISIONAL  
> Application Two:: 60/131,454  
Filing Date:: 04-28-1999

PRIOR FOREIGN APPLICATIONS

Foreign Application One:: PCT/US00/11490  
Filing Date:: 04-28-2000  
Country:: PCT  
Priority Claimed:: Yes

Source:: PrintEFS Version 1.0.1

SCANNED, # 12

## THERAPEUTIC USES OF TRI-ARYL ACID DERIVATIVES

Cross-Reference to Related Applications

This application is a continuation of International Application No. PCT/US00/11490, filed on April 28, 2000, which application, in turn, claims priority from U.S. Provisional Application No. 60/131,454, filed on April 28, 1999.

Background of the Invention

This invention is directed to the use of triaryl acid derivatives and their pharmaceutical compositions as PPAR ligand receptor binders. The PPAR ligand receptor binders of this invention are useful as agonists or antagonists of the PPAR receptor.

Field of the Invention

Peroxisome proliferator-activated receptors (PPAR) can be subdivided into three subtypes, namely: PPAR $\alpha$ , PPAR $\delta$ , and PPAR $\gamma$ . These are encoded by different genes (Motojima, Cell Structure and Function, 18:267-277, 1993). Moreover, 2 isoforms of PPAR $\gamma$  also exist, PPAR $\gamma_1$  and  $\gamma_2$ . These 2 proteins differ in their NH<sub>2</sub>-terminal-30 amino acids and are the result of alternative promoter usage and differential mRNA splicing (Vidal-Puig, Jimenez, Linan, Lowell, Hamann, Hu, Spiegelman, Flier, Moller, J. Clin. Invest., 97:2553-2561, 1996).

Biological processes modulated by PPAR are those modulated by receptors, or receptor combinations, which are responsive to the PPAR receptor ligands described herein. These processes include, for example, plasma lipid transport and fatty acid catabolism, regulation of insulin sensitivity and blood glucose levels, which are involved in hypoglycemia/hyperinsulinism (resulting from, for example, abnormal pancreatic beta cell function, insulin secreting tumors and /or autoimmune hypoglycemia due to autoantibodies to insulin, the insulin receptor, or autoantibodies that are stimulatory to pancreatic beta cells), macrophage differentiation which lead to the formation of atherosclerotic plaques, inflammatory response, carcinogenesis, hyperplasia or adipocyte differentiation.

Obesity is an excessive accumulation of adipose tissue. Recent work in this area indicates that PPAR $\gamma$  plays a central role in the adipocyte gene expression and differentiation. Excess adipose tissue is associated with the development of serious medical conditions, for example, non-insulin-dependent diabetes mellitus (NIDDM), hypertension, coronary artery disease, hyperlipidemia and certain malignancies. The adipocyte may also influence glucose homeostasis through the production of tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) and other molecules.

Non-insulin-dependent diabetes mellitus (NIDDM), or Type II diabetes, is the more common form of diabetes, with 90-95% of hyperglycemic patients experiencing this form of the disease. In NIDDM there appears to be a reduction in the pancreatic  $\beta$ -cell mass, several distinct defects in insulin secretion or a decrease in tissue sensitivity to insulin. The symptoms of this form of diabetes include



fatigue, frequent urination, thirst, blurred vision, frequent infections and slow healing of sores, diabetic nerve damage and renal disease.

Resistance to the metabolic actions of insulin is one of the key features of non-insulin dependent diabetes (NIDDM). Insulin resistance is characterised by impaired uptake and utilization of glucose in insulin-sensitive target organs, for example, adipocytes and skeletal muscle, and by impaired inhibition of hepatic glucose output. The functional insulin deficiency and the failure of insulin to suppress hepatic glucose output results in fasting hyperglycemia. Pancreatic  $\beta$ -cells compensate for the insulin resistance by secreting increased levels of insulin. However, the  $\beta$ -cells are unable to maintain this high output of insulin, and, eventually, the glucose-induced insulin secretion falls, leading to the deterioration of glucose homeostasis and to the subsequent development of overt diabetes.

Hyperinsulinemia is also linked to insulin resistance, hypertriglyceridaemia and increased plasma concentration of low density lipoproteins. The association of insulin resistance and hyperinsulinemia with these metabolic disorders has been termed "Syndrome X" and has been strongly linked to an increased risk of hypertension and coronary artery disease.

Metformin is known in the art to be used in the treatment of diabetes in humans (US Patent No. 3,174,901). Metformin acts primarily to decrease liver glucose production. Troglitazone® is known to work primarily on enhancing the ability of skeletal muscle to respond to insulin and take up glucose. It is known that combination therapy comprising metformin and troglitazone can be used in the treatment of abnormalities associated with diabetes (DDT 3:79-88, 1998).

PPAR  $\gamma$  activators, in particular Troglitazone®, have been found to convert cancerous tissue to normal cells in liposarcoma, a tumor of fat (PNAS 96:3951-3956, 1999). Furthermore, it has been suggested that PPAR  $\gamma$  activators may be useful in the treatment of breast and colon cancer (PNAS 95:8806-8811, 1998, Nature Medicine 4:1046-1052, 1998).

Moreover, PPAR $\gamma$  activators, for example Troglitazone®, have been implicated in the treatment of polycystic ovary syndrome (PCO). This is a syndrome in women that is characterized by chronic anovulation and hyperandrogenism. Women with this syndrome often have insulin resistance and an increased risk for the development of noninsulin-dependent diabetes mellitus. (Dunaif, Scott, Finegood, Quintana, Whitcomb, J. Clin. Endocrinol. Metab., 81:3299, 1996).

Furthermore, PPAR $\gamma$  activators have recently been discovered to increase the production of progesterone and inhibit steroidogenesis in granulosa cell cultures and therefore may be useful in the treatment of climacteric. (United States Patent 5,814,647 Urban et al. September 29, 1998; B. Lohrke et al. Journal of Endocrinology, 159, 429-39, 1998). Climacteric is defined as the syndrome of endocrine, somatic and psychological changes occurring at the termination of the reproductive period in the female.

Peroxisomes are cellular organelles which play a role in controlling the redox potential and oxidative stress of cells by metabolizing a variety of substrates such as hydrogen peroxide. There are a

number of disorders associated with oxidative stress. For example, inflammatory response to tissue injury, pathogenesis of emphysema, ischemia-associated organ injury (shock), doxorubicin-induced cardiac injury, drug-induced hepatotoxicity, atherosclerosis, and hyperoxic lung injuries, are each associated with the production of reactive oxygen species and a change in the reductive capacity of the cell. Therefore, it is envisaged that PPAR $\alpha$  activators, among other things, regulate the redox potential and oxidative stress in cells, would be effective in the treatment of these disorders (Poynter et al, J. Biol. Chem. 273, 32833-41, 1998).

It has also been discovered that PPAR $\alpha$  agonists inhibit NF $\kappa$ B-mediated transcription thereby modulating various inflammatory responses such as the inducible nitric oxide synthase (NOS) and cyclooxygenase-2 (COX-2) enzyme pathways (Pineda-Torra, I. T al, 1999, Curr. Opinion in Lipidology, 10,151-9 ) and thus can be used in the therapeutic intervention of a wide variety of inflammatory diseases and other pathologies (Colville-Nash, et al., Journal of Immunology, 161, 978-84, 1998; Staels et al, Nature, 393, 790-3, 1998).

Peroxisome proliferators activate PPAR , which in turn, acts as a transcription factor, and causes differentiation, cell growth and proliferation of peroxisomes. PPAR activators are also thought to play a role in hyperplasia and carcinogenesis as well as altering the enzymatic capability of animal cells, such as rodent cells, but these PPAR activators appear to have minimal negative effects in human cells (Green, Biochem. Pharm. 43(3):393, 1992). Activation of PPAR results in the rapid increase of gamma glutamyl transpeptidase and catalase.

PPAR $\alpha$  is activated by a number of medium and long-chain fatty acids and is involved in stimulating  $\beta$ -oxidation of fatty acids in tissues such as liver, heart, skeletal muscle, and brown adipose tissue (Isseman and Green, supra; Beck et al., Proc. R. Soc. Lond. 247:83-87, 1992; Gottlicher et al., Proc. Natl. Acad. Sci. USA 89:4653-4657, 1992). Pharmacological PPAR $\alpha$  activators, for example fenofibrate, clofibrate, genfibrozil, and bezafibrate, are also involved in substantial reduction in plasma triglycerides along with moderate reduction in LDL cholesterol, and they are used particularly for the treatment of hypertriglyceridemia, hyperlipidemia and obesity. PPAR $\alpha$  is also known to be involved in inflammatory disorders. (Schoonjans, K., Current Opinion in Lipidology, 8, 159-66, 1997).

The human nuclear receptor PPAR $\delta$  has been cloned from a human osteosarcoma cell cDNA library and is fully described in A. Schmidt et al., Molecular Endocrinology, 6:1634-1641 (1992), the contents of which are hereby incorporated herein by reference. It should be noted that PPAR $\delta$  is also referred to in the literature as PPAR $\beta$  and as NUC1, and each of these names refers to the same receptor. For example, in A. Schmidt et al., Molecular Endocrinology, 6: pp. 1634-1641, 1992, the receptor is referred to as NUC1. PPAR $\delta$  is observed in both embryo and adult tissues. This receptor has been reported to be involved in regulating the expression of some fat-specific genes, and plays a role in the adipogenic process (Amri, E. et al., J. Biol. Chem. 270, 2367-71, 1995).

Atherosclerotic disease is known to be caused by a number of factors, for example, hypertension, diabetes, low levels of high density lipoprotein (HDL), and high levels of low density lipoprotein (LDL). In addition to risk reduction via effects on plasma lipid concentrations and other risk factors, PPAR $\alpha$  agonists exert direct atheroprotective effects (Frick, M. H., et al. 1997.. Circulation 96:2137-2143, de Faire, et al. 1997. Cardiovasc. Drugs Ther. 11 Suppl 1:257-63:257-263).

It has recently been discovered that PPAR $\delta$  agonists are useful in raising HDL levels and therefore useful in treating atherosclerotic diseases. (Leibowitz et al.; WO/9728149). Atherosclerotic diseases include vascular disease, coronary heart disease, cerebrovascular disease and peripheral vessel disease. Coronary heart disease includes CHD death, myocardial infarction, and coronary revascularization. Cerebrovascular disease includes ischemic or hemorrhagic stroke and transient ischemic attacks.

PPAR $\gamma$  subtypes are involved in activating adipocyte differentiation, and are not involved in stimulating peroxisome proliferation in the liver. Activation of PPAR $\gamma$  is implicated in adipocyte differentiation through the activation of adipocyte-specific gene expression (Lehmann, Moore, Smith-Oliver, Wilkison, Willson, Kliewer, J. Biol. Chem., 270:12953-12956, 1995). The DNA sequences for the PPAR $\gamma$  receptors are described in Elbrecht et al., BBRC 224:431-437 (1996). Although peroxisome proliferators, including fibrates and fatty acids, activate the transcriptional activity of PPAR's, only prostaglandin J<sub>2</sub> derivatives such as the arachidonic acid metabolite 15-deoxy- $\Delta^{12,14}$ -prostaglandin J<sub>2</sub> (15d-PGJ<sub>2</sub>) have been identified as natural ligands specific for the PPAR $\gamma$  subtype, which also binds thiazolidinediones. This prostaglandin activates PPAR $\gamma$ -dependent adipogenesis, but activates PPAR $\alpha$  only at high concentrations (Forman, Tontonoz, Chen, Brun, Spiegelman, Evans, Cell, 83:803-812, 1995; Kliewer, Lenhard, Wilson, Patel, Morris, Lehman, Cell, 83:813-819, 1995). This is further evidence that the PPAR family subtypes are distinct from one another in their pharmacological response to ligands.


It has been suggested that compounds activating both PPAR $\alpha$  and PPAR $\gamma$  should be potent hypotriglyceridemic drugs, which could be used in the treatment of dyslipidemia associated with atherosclerosis, non-insulin dependent diabetes mellitus, Syndrome X, (Staels, B. et al., Curr. Pharm. Des., 3 (1), 1-14 (1997)) and familial combined hyperlipidemia (FCH). Syndrome X is the syndrome characterized by an initial insulin resistant state, generating hyperinsulinaemia, dyslipidaemia and impaired glucose tolerance, which can progress to non-insulin dependent diabetes mellitus (Type II diabetes), characterized by hyperglycemia. FCH is characterized by hypercholesterolemia and hypertriglyceridemia within the same patient and family.

The present invention is directed to a series of compounds that are useful in modulating PPAR receptors, as well as to a number of other pharmaceutical uses associated therewith.

#### Summary of the Invention

This invention provides new aromatic compounds and pharmaceutical compositions prepared therewith that are PPAR ligand receptor binders, and which are useful as agonists or antagonists of the PPAR receptors. The invention also includes the discovery of new uses for previously known compounds.

The compounds for use according to the invention, including the new compounds of the present invention, are of Formula I




wherein:




are independently aryl, fused arylcycloalkenyl, fused arylcycloalkyl, fused arylheterocyclenyl, fused arylheterocyclyl, heteroaryl, fused heteroaryl/cycloalkenyl, fused heteroaryl/cycloalkyl, fused heteroaryl/heterocyclenyl, or fused heteroaryl/heterocyclyl;

A is -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sub>13</sub>-, -C(O)-, -N(R<sub>14</sub>)C(O)-, -C(O)N(R<sub>15</sub>)-, -N(R<sub>14</sub>)C(O)N(R<sub>15</sub>)-, -C(R<sub>14</sub>)=N-,



a chemical bond,



15 B is -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sub>17</sub>-, a chemical bond, ethynylene, -C(O)-, -N(R<sub>18</sub>)C(O)-, or -C(O)NR<sub>18</sub>-;

D is -O-, -S-, -NR<sub>19</sub>-, a chemical bond, ethynylene, -C(O)-, -N(R<sub>20</sub>)C(O)-, or -C(O)N(R<sub>20</sub>)-;

E is a chemical bond or an ethylene group;

a is 0-4;

b is 0-4;

20 c is 0-4;

d is 0-5;

e is 0-4;

f is 0-6;

g is 1-4:

25 h is 1-4;

$R_1, R_3, R_5, R_7, R_9$ , and  $R_{11}$ , are independently hydrogen, halogen, alkyl, carboxyl, alkoxycarbonyl or aralkyl;

$R_2, R_4, R_6, R_8, R_{10}$  and  $R_{12}$ , are independently  $-(CH_2)_q-X$ ;

q is 0-3;

- 5 X is hydrogen, halogen, alkyl, alkenyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, aralkyl, heteroaralkyl, hydroxy, alkoxy, aralkoxy, heteroaralkoxy, carboxyl, alkoxycarbonyl, tetrazolyl, acyl, acylHNSO<sub>2</sub>-, -SR<sub>23</sub>, Y<sup>1</sup>Y<sup>2</sup>N- or Y<sup>3</sup>Y<sup>4</sup>NCO-;

Y<sup>1</sup> and Y<sup>2</sup> are independently hydrogen, alkyl, aryl, aralkyl or heteroaralkyl, or one of Y<sup>1</sup> and Y<sup>2</sup> is hydrogen or alkyl and the other of Y<sup>1</sup> and Y<sup>2</sup> is acyl or aryl;

- 10 Y<sup>3</sup> and Y<sup>4</sup> are independently hydrogen, alkyl, aryl, aralkyl or heteroaralkyl;

Z is R<sub>21</sub>O<sub>2</sub>C-, R<sub>21</sub>OC-, cyclo-imide, -CN, R<sub>21</sub>O<sub>2</sub>SHNCO-, R<sub>21</sub>O<sub>2</sub>SHN-, (R<sub>21</sub>)<sub>2</sub>NCO-, R<sub>21</sub>O- 2,4-thiazolidinedionyl, or tetrazolyl; and

R<sub>15</sub> and R<sub>21</sub> are independently hydrogen, alkyl, aryl, cycloalkyl, or aralkyl;

R<sub>13</sub>, R<sub>17</sub>, R<sub>19</sub> and R<sub>23</sub> are independently R<sub>22</sub>OC-, R<sub>22</sub>NHOC-, hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, heteroaralkyl, or aralkyl;

R<sub>14</sub>, R<sub>15</sub>, R<sub>16</sub>, R<sub>18</sub> and R<sub>20</sub> are independently hydrogen, alkyl, aralkyl, carbonyl, or alkoxycarbonyl;

or R<sub>14</sub>, and R<sub>15</sub> taken together with the carbon and nitrogen atoms through which they are linked form a 5 or 6-membered azaheterocyclyl group; or

when a is 2-4, then vicinal R<sub>1</sub> radicals taken together with the carbon atoms to which the R<sub>1</sub> radicals are linked form an ethylene group; or

when b is 2-4, then vicinal R<sub>3</sub> radicals taken together with the carbon atoms to which the R<sub>3</sub> radicals are linked form an ethylene group; or

when c is 2-4, then vicinal R<sub>5</sub> radicals taken together with the carbon atoms to which the R<sub>5</sub> radicals are linked form an ethylene group; or

- 25 when d is 2-5, then vicinal R<sub>7</sub> radicals taken together with the carbon atoms to which the R<sub>7</sub> radicals are linked form an ethylene group; or

when e is 2-4, then vicinal R<sub>9</sub> radicals taken together with the carbon atoms to which the R<sub>9</sub> radicals are linked form an ethylene group; or

when f is 2-6, then vicinal R<sub>11</sub> radicals taken together with the carbon atoms to which the R<sub>11</sub> radicals are linked form an ethylene group; and

- 30 R<sub>22</sub> is hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, heteroaralkyl, or aralkyl; or a pharmaceutically acceptable salt thereof, an N-oxide thereof, a hydrate thereof or a solvate thereof.

#### DETAILED DESCRIPTION OF THE INVENTION

As employed above and throughout the disclosure, the following terms, unless otherwise indicated, shall be understood to have the following meanings:

### Definitions

In the present specification, the term "compounds for use according to the invention", and equivalent expressions, are meant to embrace compounds of general Formula (I) as hereinbefore described, which expression includes the prodrugs, the pharmaceutically acceptable salts, and the solvates, e.g. hydrates, where the context so permits. Similarly, reference to intermediates, whether or not they themselves are claimed, is meant to embrace their salts, and solvates, where the context so permits.

For the sake of clarity, particular instances when the context so permits are sometimes indicated in the text, but these instances are purely illustrative and it is not intended to exclude other instances when the context so permits.

"Prodrug" means a compound which is convertible in vivo by metabolic means (e.g. by hydrolysis) to a compound of Formula (I), including N-oxides thereof. For example an ester of a compound of Formula (I) containing a hydroxy group may be convertible by hydrolysis in vivo to the parent molecule. Alternatively an ester of a compound of Formula (I) containing a carboxy group may be convertible by hydrolysis in vivo to the parent molecule.

"Patient" includes both human and other mammals.

"Chemical bond" means a direct single bond between atoms.

"Acyl" means an H-CO- or alkyl-CO- group wherein the alkyl group is as herein described. Preferred acyls contain a lower alkyl. Exemplary acyl groups include formyl, acetyl, propanoyl, 2-methylpropanoyl, butanoyl and palmitoyl.

"Alkenyl" means an aliphatic hydrocarbon group containing a carbon-carbon double bond and which may be a straight or branched chain having about 2 to about 15 carbon atoms in the chain.

Preferred alkenyl groups have 2 to about 12 carbon atoms in the chain and more preferably about 2 to about 4 carbon atoms in the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl are attached to a linear alkenyl chain. "Lower alkenyl" means about 2 to about 4 carbon atoms in the chain, which may be straight or branched. The alkenyl group is optionally substituted by one or more halo groups. Exemplary alkenyl groups include ethenyl, propenyl, *n*-butenyl, *i*-butenyl, 3-methylbut-2-enyl, *n*-pentenyl, heptenyl, octenyl and decenyl.

"Alkoxy" means an alkyl-O- group wherein the alkyl group is as herein described. Exemplary alkoxy groups include methoxy, ethoxy, *n*-propoxy, *i*-propoxy, *n*-butoxy and heptoxy.

"Alkoxy carbonyl" means an alkyl-O-CO- group, wherein the alkyl group is as herein defined. Exemplary alkoxy carbonyl groups include methoxycarbonyl, ethoxycarbonyl, or *t*-butoxycarbonyl.

"Alkyl" means an aliphatic hydrocarbon group which may be a straight or branched chain having about 1 to about 20 carbon atoms in the chain. Preferred alkyl groups have 1 to about 13 carbon atoms in

the chain. Branched means that one or more lower alkyl groups such as methyl, ethyl or propyl are attached to a linear alkyl chain. "Lower alkyl" means that there are about 1 to about 4 carbon atoms in the chain, which may be straight or branched. The alkyl is optionally substituted with one or more "alkyl group substituents" which may be the same or different, and include halo, carboxy, cycloalkyl, cycloalkenyl, heterocyclyl, heterocyclyl, aryl, alkoxy, alkoxycarbonyl, aralkoxycarbonyl,

heteroaralkoxycarbonyl,  $Y^1 Y^2 NCO-$ , wherein  $Y^1$  and  $Y^2$  are independently hydrogen, alkyl, aryl, aralkyl or heteroaralkyl, or  $Y^1$  and  $Y^2$  taken together with the nitrogen atom to which  $Y^1$  and  $Y^2$  are attached form heterocyclyl. Exemplary alkyl groups include methyl, trifluoromethyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *t*-butyl, *n*-pentyl, and 3-pentyl. Preferably, the alkyl group substituent is selected from acyl, halo, carboxy, carboxymethyl, methoxycarbonyl, benzyloxycarbonylmethyl, and pyridylmethyloxycarbonylmethyl and alkoxycarbonyl.

"Alkylsulfinyl" means an alkyl-SO- group wherein the alkyl group is as defined above. Preferred groups are those wherein the alkyl group is lower alkyl.

"Alkylsulfonyl" means an alkyl-SO<sub>2</sub>-group wherein the alkyl group is as defined above. Preferred groups are those wherein the alkyl group is lower alkyl.

"Alkylthio" means an alkyl-S- group wherein the alkyl group is as defined above. Exemplary alkylthio groups include methylthio, ethylthio, *i*-propylthio and heptylthio.

"Aralkoxy" means an aralkyl-O- group wherein the aralkyl group is as defined herein. Exemplary aralkoxy groups include benzyloxy and 1- and 2-naphthalenemethoxy.

"Aralkoxycarbonyl" means an aralkyl-O-CO- group wherein the aralkyl group is as defined herein. An exemplary aralkoxycarbonyl group is benzyloxycarbonyl.

"Aralkyl" means an aryl-alkyl- group wherein the aryl and alkyl groups are as defined herein. Preferred aralkyls contain a lower alkyl moiety. Exemplary aralkyl groups include benzyl, 2-phenethyl and naphthalenemethyl.

"Aralkylsulfonyl" means an aralkyl-SO<sub>2</sub>- group wherein the aralkyl group is as defined herein.

"Aralkylsulfinyl" means an aralkyl-SO- group wherein the aralkyl group is as defined herein.

"Aralkylthio" means an aralkyl-S- group wherein the aralkyl group is as defined herein. An exemplary aralkylthio group is benzylthio.

"Aroyl" means an aryl-CO- group wherein the aryl group is as defined herein. Exemplary aroyl groups include benzoyl and 1- and 2-naphthoyl.

"Aryl" means an aromatic monocyclic or multicyclic ring system of about 6 to about 14 carbon atoms, preferably of about 6 to about 10 carbon atoms. The aryl is optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined herein. Exemplary aryl groups include phenyl, naphthyl, substituted phenyl, and substituted naphthyl.

"Aryldiazo" means an aryl-diazo- group wherein the aryl and diazo groups are as defined herein.

“Fused arylcycloalkenyl” means a fused aryl and cycloalkenyl as defined herein. Preferred fused arylcycloalkenyls are those wherein the aryl thereof is phenyl and the cycloalkenyl consists of about 5 to about 6 ring atoms. A fused arylcycloalkenyl group may be bonded to the rest of the compound through any atom of the fused system capable of such bondage. The fused arylcycloalkenyl may be optionally substituted by one or more ring system substituents, wherein the “ring system substituent” is as defined herein. Exemplary fused arylcycloalkenyl groups include 1,2-dihydronaphthylenyl; indenyl; 1,4-naphthoquinonyl, and the like.

“Fused arylcycloalkyl” means a fused aryl and cycloalkyl as defined herein. Preferred fused arylcycloalkyls are those wherein the aryl thereof is phenyl and the cycloalkyl consists of about 5 to about 6 ring atoms. A fused arylcycloalkyl group may be bonded to the rest of the compound through any atom of the fused system capable of such bonding. The fused arylcycloalkyl may be optionally substituted by one or more ring system substituents, wherein the “ring system substituent” is as defined herein. Exemplary fused arylcycloalkyl and substituted fused arylcycloalkyl groups include 1,2,3,4-tetrahydronaphthyl; 1,4-dimethyl-2,3-dihydronaphthyl; 2,3-dihydro-1,4-naphthoquinonyl,  $\alpha$ -tetralonyl,  $\beta$ -tetralonyl and the like.

“Fused arylheterocyclenyl” means a fused aryl and heterocyclenyl wherein the aryl and heterocyclenyl groups are as defined herein. Preferred fused arylheterocyclenyl groups are those wherein the aryl thereof is phenyl and the heterocyclenyl consists of about 5 to about 6 ring atoms. A fused arylheterocyclenyl group may be bonded to the rest of the compound through any atom of the fused system capable of such bonding. The designation of aza, oxa or thia as a prefix before the heterocyclenyl portion of the fused arylheterocyclenyl means that a nitrogen, oxygen or sulfur atom respectively, is present as a ring atom. The fused arylheterocyclenyl may be optionally substituted by one or more ring system substituents, wherein the “ring system substituent” is as defined herein. The nitrogen atom of a fused arylheterocyclenyl may be a basic nitrogen atom. The nitrogen or sulphur atom of the heterocyclenyl portion of the fused arylheterocyclenyl is also optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Exemplary fused arylheterocyclenyl and substituted fused arylheterocyclenyl groups include 3H-indolyl, 2(1H)quinolinyl, 4-oxo-1,4-dihydroquinolinyl, 2H-1-oxoisquinolyl, 1,2-dihydroquinolinyl, (2H)quinolinyl N-oxide, 3,4-dihydroquinolinyl, 1,2-dihydroisquinolinyl, 3,4-dihydroisquinolinyl, chromonyl, 3,4-dihydroisquinoxalyl, 4-(3H)quinazolinonyl, 4H-chromen-2yl, and the like. Preferably, 2(1H)quinolinonyl, 1,2-dihydroquinolinyl, (2H)quinolinyl N-oxide, or 4-(3H)quinazolinonyl.

“Fused arylheterocyclyl” means a fused aryl and heterocyclyl wherein the aryl and heterocyclyl groups are as defined herein. Preferred fused arylheterocyclyls are those wherein the aryl thereof is phenyl and the heterocyclyl consists of about 5 to about 6 ring atoms. A fused arylheterocyclyl may be bonded to the rest of the compound through any atom of the fused system capable of such bonding. The designation of aza, oxa or thia as a prefix before the heterocyclyl portion of the fused arylheterocyclyl



means that a nitrogen, oxygen or sulphur atom respectively is present as a ring atom. The fused arylheterocyclyl group may be optionally substituted by one or more ring system substituents, wherein the "ring system substituent" is as defined herein. The nitrogen atom of a fused arylheterocyclyl may be a basic nitrogen atom. The nitrogen or sulphur atom of the heterocyclyl portion of the fused

arylheterocyclyl is also optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Exemplary fused arylheterocyclyl and substituted fused arylheterocyclyl groups include indolinyl, o-benzoic sulfimidyl, 4-chromanonyl, oxindole, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinolinyl, 1H-2,3-dihydroisoindol-2-yl, 2,3-dihydrobenz[f]isoindol-2-yl, 1,2,3,4-tetrahydrobenz[g]isoquinolin-2-yl, chromanyl, isochromanonyl, 2,3-dihydrochromonyl, 1,4-benzodioxan, 1,2,3,4-tetrahydroquinoxaliny, and the like. Preferably, 1,2,3,4-tetrahydroisoquinolinyl, 1,2,3,4-tetrahydroquinoxaliny, and 1,2,3,4-tetrahydroquinolinyl.

"Aryloxy" means an aryl-O- group wherein the aryl group is as defined herein. Exemplary groups include phenoxy and 2-naphthylloxy.

"Aryloxycarbonyl" means an aryl-O-CO- group wherein the aryl group is as defined herein.

Exemplary aryloxycarbonyl groups include phenoxycarbonyl and naphthoxycarbonyl.

"Arylsulfonyl" means an aryl-SO<sub>2</sub>- group wherein the aryl group is as defined herein.

"Arylsulfinyl" means an aryl-SO- group wherein the aryl group is as defined herein.

"Arylthio" means an aryl-S- group wherein the aryl group is as defined herein. Exemplary arylthio groups include phenylthio and naphthylthio.

"Carbamoyl" is an NH<sub>2</sub>-CO- group.

"Carboxy" means a HO(O)C- (carboxylic acid) group.

"Compounds of the invention," and equivalent expressions, are meant to embrace compounds of general Formula (I) as hereinbefore described, which expression includes the prodrugs, the pharmaceutically acceptable salts, and the solvates, e.g. hydrates, where the context so permits.

Similarly, reference to intermediates, whether or not they themselves are claimed, is meant to embrace their salts, and solvates, where the context so permits. For the sake of clarity, particular instances when the context so permits are sometimes indicated in the text, but these instances are purely illustrative and it is not intended to exclude other instances when the context so permits.

"Cycloalkoxy" means an cycloalkyl-O- group wherein the cycloalkyl group is as defined herein.

Exemplary cycloalkoxy groups include cyclopentylloxy and cyclohexylloxy.


"Cycloalkenyl" means a non-aromatic mono- or multicyclic ring system of about 3 to about 10 carbon atoms, preferably of about 5 to about 10 carbon atoms, and which contains at least one carbon-carbon double bond. Preferred ring sizes of rings of the ring system include about 5 to about 6 ring atoms. The cycloalkenyl is optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined herein. Exemplary monocyclic cycloalkenyl include

cyclopentenyl, cyclohexenyl, cycloheptenyl, and the like. An exemplary multicyclic cycloalkenyl is norbornenyl.

"Cycloalkyl" means a non-aromatic mono- or multicyclic ring system of about 3 to about 10 carbon atoms, preferably of about 5 to about 10 carbon atoms. Preferred ring sizes of rings of the ring system include about 5 to about 6 ring atoms. The cycloalkyl is optionally substituted with one or more "ring system substituents" which may be the same or different, and are as defined herein. Exemplary monocyclic cycloalkyl include cyclopentyl, cyclohexyl, cycloheptyl, and the like. Exemplary multicyclic cycloalkyl include 1-decalin, norbornyl, adamant-(1- or 2-)yl, and the like.

"Cycloalkylene" means a bivalent, saturated carbocyclic group having about 3 to about 6 carbon atoms. Preferred cycloalkylene groups include 1,1-, 1,2-, 1,3-, and 1,4- cis or trans-cyclohexylene; and 1,1-, 1,2-, and 1,3-cyclopentylene.

"Cyclo-imide" means a compound of formulae



The cyclo-imide moiety may be attached to the parent molecule through either a carbon atom or nitrogen atom of the carbamoyl moiety. An exemplary imide group is N-phthalimide.

"Diazo" means a bivalent  $-N=N-$  radical.

"Halo" means fluoro, chloro, bromo, or iodo. Preferred are fluoro, chloro and bromo, more preferably fluoro and chloro.

"Heteroalkyl" means a heteroaryl-alkyl- group wherein the heteroaryl and alkyl groups are as defined herein. Preferred heteroalkyls contain a lower alkyl moiety. Exemplary heteroalkyl groups include thienylmethyl, pyridylmethyl, imidazolylmethyl and pyrazinylmethyl.

"Heteroalkylthio" means a heteroalkyl-S- group wherein the heteroalkyl group is as defined herein. An exemplary heteroalkylthio group is 3-pyridinepropanthiol.

"Heteroalkoxy" means a heteroalkyl-O- group wherein the heteroalkyl group is as defined herein. An exemplary heteroalkoxy group is 4-pyridylmethoxy.

"Heteroaryl" means an means an heteroaryl-CO- group wherein the heteroaryl group is as defined herein. Exemplary heteroaryl groups include thiophenyl, nicotinyl, pyrrol-2-ylcarbonyl and 1- and 2-naphthoyl and pyridinoyl.

"Heteroaryldiazo" means an heteroaryl-diazo- group wherein the heteroaryl and diazo groups are as defined herein.

"Heteroaryl" means an aromatic monocyclic or multicyclic ring system of about 5 to about 14 carbon atoms, preferably about 5 to about 10 carbon atoms, in which at least one of the carbon atoms in the ring system is replaced by a hetero atom, i.e., other than carbon, for example nitrogen, oxygen or sulfur. Preferred ring sizes of rings of the ring system include about 5 to about 6 ring atoms. The heteroaryl ring is optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The designation of aza, oxa or thia as a prefix before the heteroaryl means that a nitrogen, oxygen or sulfur atom is present, respectively, as a ring atom. A nitrogen atom of a heteroaryl may be a basic nitrogen atom and also may be optionally oxidized to the corresponding N-oxide. Exemplary heteroaryl and substituted heteroaryl groups include pyrazinyl, thienyl, isothiazolyl, oxazolyl, pyrazolyl, cinnolyl, pteridinyl, benzofuryl, furazanyl, pyrrolyl, 1,2,4-oxadiazolyl, benzoxazole, 1,2,4-thiadiazolyl, pyridazinyl, indazolyl, quinoxalyl, phthalazinyl, imidazo[1,2-a]pyridine, imidazo[2,1-b]thiazolyl, benzofurazanyl, azaindolyl, benzimidazolyl, benzothienyl, thienopyridyl, thienopyrimidyl, pyrrolopyridyl, imidazopyridyl, naphthyridinyl, benzoazaindole, 1,2,4-triazinyl, benzothiazolyl, furyl, imidazolyl, indolyl, isoindolyl, indolizyl, isoxazolyl, isoquinolyl, isothiazolyl, oxadiazolyl, pyrazinyl, pyridazinyl, pyrazolyl, pyridyl, pyrimidinyl, pyrrolyl, quinazolinyl, quinolyl, 1,3,4-thiadiazolyl, thiazolyl, thienyl and triazolyl. Preferred heteroaryl and substituted heteroaryl groups include quinolyl, indazolyl, indolyl, quinazolinyl, pyridyl, pyrimidinyl, furyl, benzothiazolyl, benzoxazole, benzofuryl, quinoxalyl, benzimidazolyl, 1,2,4-oxadiazolyl, benzothienyl, and isoquinolyl.

"Fused heteroarylcycloalkenyl" means a fused heteroaryl and cycloalkenyl wherein the heteroaryl and cycloalkenyl groups are as defined herein. Preferred fused heteroarylcycloalkenyls are those wherein the heteroaryl thereof is phenyl and the cycloalkenyl consists of about 5 to about 6 ring atoms. A fused heteroarylcycloalkenyl may be bonded to the rest of the compound through any atom of the fused system capable of such bonding. The designation of aza, oxa or thia as a prefix before the heteroaryl portion of the fused heteroarylcycloalkenyl means that a nitrogen, oxygen or sulfur atom is present, respectively, as a ring atom. The fused heteroarylcycloalkenyl may be optionally substituted by one or more ring system substituents, wherein the "ring system substituent" is as defined herein. The nitrogen atom of a fused heteroarylcycloalkenyl may be a basic nitrogen atom. The nitrogen atom of the heteroaryl portion of the fused heteroarylcycloalkenyl may also be optionally oxidized to the corresponding N-oxide. Exemplary fused heteroarylcycloalkenyl groups include 5,6-dihydroquinolyl; 5,6-dihydroisoquinolyl; 5,6-dihydroquinoxalyl; 5,6-dihydroquinazolinyl; 4,5-dihydro-1H-benzimidazolyl; 4,5-dihydrobenzoxazolyl; 1,4-naphthoquinolyl, and the like.

"Fused heteroarylcycloalkyl" means a fused heteroaryl and cycloalkyl wherein the heteroaryl and cycloalkyl groups are as defined herein. Preferred fused heteroarylcycloalkyls are those wherein the

heteroaryl thereof consists of about 5 to about 6 ring atoms and the cycloalkyl consists of about 5 to about 6 ring atoms. A fused heteroarylcycloalkyl may be bonded to the rest of the compound through any atom of the fused system capable of such bonding. The designation of aza, oxa or thia as a prefix before the heteroaryl portion of the fused heteroarylcycloalkyl means that a nitrogen, oxygen or sulfur atom is present respectively as a ring atom. The fused heteroarylcycloalkyl may be optionally substituted by one or more ring system substituents, wherein the "ring system substituent" is as defined herein. The nitrogen atom of a fused heteroarylcycloalkyl may be a basic nitrogen atom. The nitrogen atom of the heteroaryl portion of the fused heteroarylcycloalkyl may also be optionally oxidized to the corresponding N-oxide. Exemplary fused heteroarylcycloalkyl include 5,6,7,8-tetrahydroquinolyl; 5,6,7,8-tetrahydroisoquinolyl; 5,6,7,8-tetrahydroquinazolyl; 4,5,6,7-tetrahydro-1H-benzimidazolyl; 4,5,6,7-tetrahydrobenzoxazolyl; 1H-4-oxa-1,5-diazanaphthalen-2-yl; 1,3-dihydroimidazole-[4,5]-pyridin-2-yl; 2,3-dihydro-1,4-dinaphthoquinonyl and the like, preferably, 5,6,7,8-tetrahydroquinolyl or 5,6,7,8-tetrahydroisoquinolyl.

"Fused heteroarylheterocyclenyl" means a fused heteroaryl and heterocyclenyl wherein the heteroaryl and heterocyclenyl groups are as defined herein. Preferred fused heteroarylheterocyclenyls are those wherein the heteroaryl thereof consists of about 5 to about 6 ring atoms and the heterocyclenyl consists of about 5 to about 6 ring atoms. A fused heteroarylheterocyclenyl may be bonded to the rest of the compound through any atom of the fused system capable of such bonding. The designation of aza, oxa or thia as a prefix before the heteroaryl or heterocyclenyl portion of the fused heteroarylheterocyclenyl means that a nitrogen, oxygen or sulfur atom is present respectively as a ring atom. The fused heteroarylheterocyclenyl may be optionally substituted by one or more ring system substituent, wherein the "ring system substituent" is as defined herein. The nitrogen atom of a fused heteroarylazaheterocyclenyl may be a basic nitrogen atom. The nitrogen or sulphur atom of the heteroaryl or heterocyclenyl portion of the fused heteroarylheterocyclenyl may also be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Exemplary fused heteroarylheterocyclenyl groups include 7,8-dihydro[1,7]naphthyridinyl; 1,2-dihydro[2,7]naphthyridinyl; 6,7-dihydro-3H-imidazo[4,5-c]pyridyl; 1,2-dihydro-1,5-naphthyridinyl; 1,2-dihydro-1,6-naphthyridinyl; 1,2-dihydro-1,7-naphthyridinyl; 1,2-dihydro-1,8-naphthyridinyl; 1,2-dihydro-2,6-naphthyridinyl, and the like.

"Fused heteroarylheterocyclyl" means a fused heteroaryl and heterocyclyl wherein the heteroaryl and heterocyclyl groups are as defined herein. Preferred fused heteroarylheterocyclyls are those wherein the heteroaryl thereof consists of about 5 to about 6 ring atoms and the heterocyclyl consists of about 5 to about 6 ring atoms. A fused heteroarylheterocyclyl may be bonded to the rest of the compound through any atom of the fused system capable of such bonding. The designation of aza, oxa or thia as a prefix before the heteroaryl or heterocyclyl portion of the fused heteroarylheterocyclyl means that a nitrogen, oxygen or sulfur atom is present respectively as a ring atom. The fused heteroarylheterocyclyl may be

optionally substituted by one or more ring system substituent, wherein the "ring system substituent" is as defined herein. The nitrogen atom of a fused heteroaryl/heterocyclyl may be a basic nitrogen atom. The nitrogen or sulphur atom of the heteroaryl or heterocyclyl portion of the fused heteroaryl/heterocyclyl may also be optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Exemplary fused heteroaryl/heterocyclyl groups include 2,3-dihydro-1H-pyrrol[3,4-b]quinolin-2-yl; 1,2,3,4-tetrahydrobenz[b][1,7]naphthyridin-2-yl; 1,2,3,4-tetrahydrobenz[b][1,6]naphthyridin-2-yl; 1,2,3,4-tetrahydro-9H-pyrido[3,4-b]indol-2-yl; 1,2,3,4-tetrahydro-9H-pyrido[4,3-b]indol-2-yl, 2,3-dihydro-1H-pyrrolo[3,4-b]indol-2-yl; 1H-2,3,4,5-tetrahydroazepino[3,4-b]indol-2-yl; 1H-2,3,4,5-tetrahydroazepino[4,3-b]indol-3-yl; 1H-2,3,4,5-tetrahydroazepino[4,5-b]indol-2-yl, 5,6,7,8-tetrahydro[1,7]naphthyridinyl; 1,2,3,4-tetrahydro[2,7]naphthyridyl; 2,3-dihydro[1,4]dioxino[2,3-b]pyridyl; 2,3-dihydro[1,4]dioxino[2,3-b]pyridyl; 3,4-dihydro-2H-1-oxa[4,6]diazanaphthalenyl; 4,5,6,7-tetrahydro-3H-imidazo[4,5-c]pyridyl; 6,7-dihydro[5,8]diazanaphthalenyl; 1,2,3,4-tetrahydro[1,5]naphthyridinyl; 1,2,3,4-tetrahydro[1,6]naphthyridinyl; 1,2,3,4-tetrahydro[1,7]naphthyridinyl; 1,2,3,4-tetrahydro[1,8]naphthyridinyl; 1,2,3,4-tetrahydro[2,6]naphthyridinyl, and the like.

"Heteroarylsulfonyl" means an heteroaryl-SO<sub>2</sub>- group wherein the heteroaryl group is as defined herein. An exemplary heteroarylsulfonyl group is 3-pyridinepropane-sulfonyl.

"Heteroarylsulfinyl" means an heteroaryl-SO- group wherein the heteroaryl group is as defined herein.

"Heteroarylthio" means an heteroaryl-S- group wherein the heteroaryl group is as defined herein. Exemplary heteroaryl thio groups include pyridylthio and quinolinylthio.

"Heterocyclenyl" means a non-aromatic monocyclic or multicyclic hydrocarbon ring system of about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms, in which at least one or more of the carbon atoms in the ring system is replaced by a hetero atom, for example a nitrogen, oxygen or sulfur atom, and which contains at least one carbon-carbon double bond or carbon-nitrogen double bond. Preferred ring sizes of rings of the ring system include about 5 to about 6 ring atoms. The designation of aza, oxa or thia as a prefix before the heterocyclenyl means that a nitrogen, oxygen or sulfur atom is present respectively as a ring atom. The heterocyclenyl may be optionally substituted by one or more ring system substituents, wherein the "ring system substituent" is as defined herein. The nitrogen atom of an heterocyclenyl may be a basic nitrogen atom. The nitrogen or sulphur atom of the heterocyclenyl is also optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide.

Exemplary monocyclic azaheterocyclenyl and substituted monocyclic azaheterocyclenyl groups include 1,2,3,4-tetrahydrohydropyridine, 1,2-dihydropyridyl, 1,4-dihydropyridyl, 1,2,3,6-tetrahydropyridine, 4(3H)pyrimidone, 1,4,5,6-tetrahydropyrimidine, 2-pyrrolinyl, 3-pyrrolinyl, 2-imidazolinyl, 2-pyrazolinyl, and the like. Exemplary oxaheterocyclenyl groups include 3,4-dihydro-2H-pyran, dihydrofuryl, and fluorodihydrofuryl. An exemplary multicyclic oxaheterocyclenyl group is 7-oxabicyclo[2.2.1]heptenyl. Exemplary monocyclic thiaheterocyclenyl rings include dihydrothiophenyl and dihydrothiopyranyl.

"Heterocyclyl" means a non-aromatic saturated monocyclic or multicyclic ring system of about 3 to about 10 carbon atoms, preferably about 5 to about 10 carbon atoms, in which at least one of the carbon atoms in the ring system is replaced by a hetero atom, for example nitrogen, oxygen or sulfur. Preferred ring sizes of rings of the ring system include about 5 to about 6 ring atoms. The designation of aza, oxa or thia as a prefix before the heterocyclyl means that a nitrogen, oxygen or sulfur atom is present respectively as a ring atom. The heterocyclyl may be optionally substituted by one or more "ring system substituents" which may be the same or different, and are as defined herein. The nitrogen atom of an heterocyclyl may be a basic nitrogen atom. The nitrogen or sulphur atom of the heterocyclyl is also optionally oxidized to the corresponding N-oxide, S-oxide or S,S-dioxide. Exemplary monocyclic heterocyclyl rings include piperidyl, pyrrolidinyl, piperazinyl, morpholinyl, thiomorpholinyl, thiazolidinyl, 1,3-dioxolanyl, 1,4-dioxanyl, tetrahydrofuryl, tetrahydrothiophenyl, tetrahydrothiopyranyl, and the like. Exemplary multicyclic heterocyclyl rings include 1,4 diazabicyclo-[2.2.2]octane and 1,2-cyclohexanedicarboxylic acid anhydride.

"Ring system substituent" includes hydrogen, alkyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, aralkyl, heteroaralkyl, hydroxy, alkoxy, aryloxy, aralkoxy, acyl, aroyl, halo, nitro, cyano, carboxy, alkoxycarbonyl, aryloxy carbonyl, aralkoxycarbonyl, alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, alkylsulfinyl, arylsulfinyl, heteroarylsulfinyl, alkylthio, arylthio, heteroarylthio, aralkylthio, heteroaralkylthio, fused cycloalkyl, fused cycloalkenyl, fused heterocyclyl, fused heterocyclenyl, arylazo, heteroarylazo,  $R^aR^bN-$ ,  $R^cR^dNCO-$ ,  $R^eO_2CN-$ , and  $R^fR^gNSO_2-$  wherein  $R^a$  and  $R^b$  are independently hydrogen, alkyl, aryl, aralkyl or heteroaralkyl, or one of  $R^a$  and  $R^b$  is hydrogen or alkyl and the other of  $R^a$  and  $R^b$  is aroyl or heteroaroyl.  $R^c$  and  $R^d$  are independently hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocyclyl, heterocyclenyl, aralkyl or heteroaralkyl. Where the ring is cycloalkyl, cycloalkenyl, heterocyclyl or heterocyclenyl, the ring system substituent may also include methylene ( $H_2C=$ ), oxo ( $O=$ ), thioxo ( $S=$ ), on carbon atom(s) thereof. Preferably, the ring substituents are selected from oxo ( $O=$ ), (lower) alkyl, aryl, alkoxy, aralkoxy, halo, trifluoromethyl, carboxy, alkoxycarbonyl, optionally substituted phenyl, optionally substituted benzyloxy, optionally substituted cyclohexyl, optionally substituted cyclobutyl, optionally substituted heteroaryl, and  $R^eO_2CN-$ , wherein  $R^e$  is cycloalkyl.

"Tetrazolyl" means a group of formula



wherein the hydrogen atom thereof is optionally replaced by alkyl, carboxyalkyl or alkoxycarbonylalkyl.

"PPAR ligand receptor binder" means a ligand which binds to the PPAR receptor. PPAR ligand receptor binders of this invention are useful as agonists or antagonists of the PPAR- $\alpha$ , PPAR- $\delta$ , or PPAR- $\gamma$  receptor.

The term "pharmaceutically acceptable salt" refers to a relatively non-toxic, inorganic or organic acid addition salt of a compound of the present invention. A salt can be prepared *in situ* during the final isolation and purification of a compound or by separately reacting the purified compound in its free base form with a suitable organic or inorganic acid and isolating the salt thus formed. Representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, phosphate, nitrate, acetate, oxalate, valerate, oleate, palmitate, stearate, laurate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate, mesylate, glucoheptonate, lactobionate, laurylsulphonate salts, and the like. (See, for example S. M. Berge, et al., "Pharmaceutical Salts," *J. Pharm. Sci.*, 66: 1-19, 1977, the contents of which are hereby incorporated herein by reference.)

"Treating" means the partial or complete relieving or preventing of one or more physiological or biochemical parameters associated with PPAR activity.

The term "modulate" refers to the ability of a compound to either directly (by binding to the receptor as a ligand) or indirectly (as a precursor for a ligand or an inducer which promotes production of a ligand from a precursor) induce expression of gene(s) maintained under hormone control, or to repress expression of gene (s) maintained under such control.

The term "obesity" refers generally to individuals who are at least about 20-30% over the average weight for the person's age, sex and height. Technically, "obese" is defined, for males, as individuals whose body mass index is greater than 27.3 kg/m<sup>2</sup>. Those skilled in the art readily recognize that the invention method is not limited to those who fall within the above criteria. Indeed, the invention method can also be advantageously practiced by individuals who fall outside of these traditional criteria, for example by those who are prone to obesity.

The phrase "amount effective to lower blood glucose levels" refers to levels of a compound sufficient to provide circulating concentrations high enough to accomplish the desired effect. Such a concentration typically falls in the range of about 10nM up to 2 $\mu$ M, with concentrations in the range of about 100nm up to about 500nm being preferred.

The phrase "amount effective to lower tricyceride levels" refers to levels of a compound sufficient to provide circulating concentrations high enough to accomplish the desired effect. Such a concentration typically falls in the range of about 10nM up to 2 $\mu$ M; with concentrations in the range of about 100nm up to about 500nm being preferred.

#### Preferred Embodiments

Preferred embodiments according to the invention includes the use of compounds of Formula I (and their pharmaceutical compositions) as binders for PPAR receptors.

More particularly, the use of compounds of Formula I that bind to the PPAR- $\alpha$  receptor,  
compounds of Formula I that bind to the PPAR- $\delta$  receptor,  
compounds of Formula I that bind to the PPAR- $\gamma$  receptor,  
compounds of Formula I that bind to the PPAR- $\alpha$  and the PPAR- $\gamma$  receptor,  
compounds of Formula I that bind to the PPAR- $\alpha$  and the PPAR- $\delta$  receptor,  
compounds of Formula I that bind to the PPAR- $\gamma$  and the PPAR- $\delta$  receptor,  
compounds of Formula I that act as PPAR receptor agonists,  
compounds of Formula I that act as PPAR- $\alpha$  receptor agonists,  
compounds of Formula I that act as PPAR- $\delta$  receptor agonists,  
compounds of Formula I that act as PPAR- $\gamma$  receptor agonists,  
compounds of Formula I that act as both PPAR- $\alpha$  and PPAR- $\gamma$  receptor agonists,  
compounds of Formula I that act as both PPAR- $\alpha$  and PPAR- $\delta$  receptor agonists,  
compounds of Formula I that act as both PPAR- $\gamma$  and PPAR- $\delta$  receptor agonists,  
compounds of Formula I that act as both PPAR- $\alpha$  receptor antagonists and PPAR- $\gamma$  receptor

agonists,

compounds of Formula I that act as both PPAR- $\alpha$  receptor antagonists and PPAR- $\delta$  receptor  
agonists,

compounds of Formula I and act as both PPAR- $\gamma$  receptor antagonists and PPAR- $\delta$  receptor  
agonists,

compounds of Formula I that act as both PPAR- $\alpha$  receptor agonists and PPAR- $\gamma$  receptor  
antagonists,

compounds of Formula I that act as both PPAR- $\alpha$  receptor agonists and PPAR- $\delta$  receptor  
antagonists,

compounds of Formula I that act as both PPAR- $\gamma$  receptor agonists and PPAR- $\delta$  receptor  
antagonists,

compounds of Formula I that act as PPAR receptor antagonists,

compounds of Formula I that act as PPAR- $\alpha$  receptor antagonists,

compounds of Formula I that act as PPAR- $\delta$  receptor antagonists,

compounds of Formula I that act as PPAR- $\gamma$  receptor antagonists,

compounds of Formula I that act as both PPAR- $\alpha$  and PPAR- $\gamma$  receptor antagonists,

compounds of Formula I that act as both PPAR- $\alpha$  and PPAR- $\delta$  receptor antagonists, and

compounds of Formula I that act as both PPAR- $\gamma$  and PPAR- $\delta$  receptor antagonists.

An embodiment according to the invention is directed to treating a patient suffering from a  
physiological disorder capable of being modulated by a compound of Formula I having PPAR ligand  
binding activity, comprising administering to the patient a pharmaceutically effective amount of the



compound, or a pharmaceutically acceptable salt thereof. Physiological disorders capable of being so modulated include, for example, cell differentiation to produce lipid accumulating cells, regulation of insulin sensitivity and blood glucose levels, which are involved in hypoglycemia/hyperinsulinism (resulting from, for example, abnormal pancreatic beta cell function, insulin secreting tumors and /or autoimmune hypoglycemia due to autoantibodies to insulin, autoantibodies to the insulin receptor, or autoantibodies that are stimulatory to pancreatic beta cells), macrophage differentiation which leads to the formation of atherosclerotic plaques, inflammatory response, carcinogenesis, hyperplasia, adipocyte gene expression, adipocyte differentiation, reduction in the pancreatic  $\beta$ -cell mass, insulin secretion, tissue sensitivity to insulin, liposarcoma cell growth, chronic anovulation, hyperandrogenism, progesterone production, steroidogenesis, redox potential and oxidative stress in cells, nitric oxide synthase (NOS) production, increased gamma glutamyl transpeptidase, catalase, plasma triglycerides, HDL and LDL cholesterol levels and the like.

Another embodiment according to the invention is directed to a method of treating a disease state in a patient with a pharmaceutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein the disease is associated with a physiological detrimental blood level of insulin, glucose, free fatty acids (FFA), or tricyclerides.

An embodiment according to the invention is directed to treating a patient suffering from a physiological disorder associated with physiologically detrimental levels of tricyclerides in the blood, by administering to the patient a pharmaceutically effective amount of the compound, or of a pharmaceutically acceptable salt thereof.

An embodiment according to the invention is the use of compounds of Formula I and their pharmaceutical compositions as anti-diabetic, anti-lipidemic, anti-hypertensive or anti-arteriosclerotic agents, or in the treatment of obesity.

Another embodiment according to the invention is directed to a method of treating hyperglycemia in a patient, by administering to the patient a pharmaceutically effective amount to lower blood glucose levels of a compound of Formula I, or a pharmaceutically acceptable salt thereof. Preferably, the form of hyperglycemia treated in accordance with this invention is Type II diabetes.

Another embodiment according to the invention is directed to a method of reducing triglyceride levels in a patient, comprising administering to the patient a therapeutically effective amount (to lower triglyceride levels) of a compound of Formula I, or a pharmaceutically acceptable salt thereof.

Another embodiment according to the invention is directed to a method of treating hyperinsulinism in a patient, comprising administering to the patient a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof.

Another embodiment according to the invention is directed to a method of treating insulin resistance in a patient, comprising administering to the patient a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof.

Another embodiment according to the invention is directed to a method of treating cardiovascular disease, such as atherosclerosis in a patient, comprising administering to the patient a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof.

Another embodiment according to the invention is directed to treating of hyperlipidemia in a patient, comprising administering to the patient a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof.

Another embodiment according to the invention is directed to treating of hypertension in a patient, comprising administering to the patient a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof.

Another embodiment according to the invention is directed to treating eating disorders in a patient, comprising administering to the patient a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof. Treatment of eating disorders includes the regulation of appetite and/or food intake in patients suffering from under-eating disorders such as anorexia nervosa as well as over-eating disorders such as obesity and anorexia bulimia.

Another embodiment according to the invention is directed to treating a disease state associated with low levels of HDL comprising administering to the patient a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof. Diseases associated with low levels of HDL include atherosclerotic diseases.

Another embodiment according to the invention is directed to treating polycystic ovary syndrome comprising administering to the patient a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof.

Another embodiment according to the invention is directed to treating climacteric comprising administering to the patient a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof.

Another embodiment according to the invention is directed to treating inflammatory diseases such as rheumatoid arthritis, chronic obstructive pulmonary disease (emphysema or chronic bronchitis), or asthma comprising administering to the patient a therapeutically effective amount of a compound of Formula I, or a pharmaceutically acceptable salt thereof.

Another aspect of the invention is to provide a novel pharmaceutical composition which is effective, in and of itself, for utilization in a beneficial combination therapy because it includes a plurality of active ingredients which may be utilized in accordance with the invention.

In another aspect, the present invention provides a method for treating a disease state in a patient, wherein the disease is associated with a physiological detrimental level of insulin, glucose, free fatty acids (FFA), or triglycerides, in the blood, comprising administering to the patient a therapeutically effective amount of a compound of Formula I, and also administering a therapeutically effective amount of an additional hypoglycemic agent.

In another aspect, the present invention provides a method for treating a disease state in a patient, wherein the disease is associated with a physiological detrimental level of insulin, glucose, free fatty acids (FFA), or triglycerides, in the blood, comprising administering to the patient a therapeutically effective amount of a compound of Formula I, and also administering a therapeutically effective amount of a biguanidine compound.

In another aspect, the present invention provides a method for treating a disease state in a patient, wherein the disease is associated with a physiological detrimental level of insulin, glucose, free fatty acids (FFA), or triglycerides, in the blood, comprising administering to the patient a therapeutically effective amount of a compound of Formula I, and also administering a therapeutically effective amount of metformin.

The invention also provides kits or single packages combining two or more active ingredients useful in treating the disease. A kit may provide (alone or in combination with a pharmaceutically acceptable diluent or carrier), a compound of Formula (I) and an additional hypoglycaemic agent (alone or in combination with diluent or carrier).

There are many known hypoglycemic agents in the art, for example, insulin; biguanidines, such as metformin and buformin; sulfonylureas, such as acetohexamide, chlorpropamide, tolazamide, tolbutamide, glyburide, glypizide and glyclazide; thiazolidinediones, such as troglitazone;  $\alpha$ -glycosidase inhibitors, such as acarbose and miglatol; and  $B_3$  adrenoreceptor agonists such as CL-316, 243.

Since sulfonylureas are known to be capable of stimulating insulin release, but are not capable of acting on insulin resistance, and compounds of Formula I are able to act on insulin resistance, it is envisaged that a combination of these medicaments could be used as a remedy for conditions associated with both deficiency in insulin secretion and insulin-resistance.

Therefore, the invention also provides a method of treating diabetes mellitus of type II in a patient comprising administering a compound of Formula I and one or more additional hypoglycemic agents selected from the group consisting of sulfonylureas, biguanidines, thiazolidinediones,  $B_3$ -adrenoreceptor agonists,  $\alpha$ -glycosidase inhibitors and insulin.

The invention also provides a method of treating diabetes mellitus of type II in a patient comprising administering a compound of Formula I and a sulfonylurea selected from the group consisting of acetohexamide, chlorpropamide, tolazamide, tolbutamide, glyburide, glypizide and glyclazide.

The invention also provides a method of treating diabetes mellitus of type II in a patient comprising administering a compound of Formula I and a biguanidine selected from the group consisting of metformin and buformin.

The invention also provides a method of treating diabetes mellitus of type II in a patient comprising administering a compound of Formula I and an  $\alpha$ -glycosidase inhibitor selected from the group consisting of acarbose and miglatol.


The invention also provides a method of treating diabetes mellitus of type II in a patient comprising administering a compound of Formula I and an thiazolidinedione, for example, troglitazone.


As indicated above, a compound of Formula I may be administered alone or in combination with one or more additional hypoglycemic agents. Combination therapy includes administration of a single pharmaceutical dosage formulation which contains a compound of Formula I and one or more additional hypoglycemic agent, as well as administration of the compound of Formula I and each additional hypoglycemic agents in its own separate pharmaceutical dosage formulation. For example, a compound of Formula I and hypoglycemic agent can be administered to the patient together in a single oral dosage composition such as a tablet or capsule, or each agent administered in separate oral dosage formulations. Where separate dosage formulations are used, the compound of Formula I and one or more additional hypoglycemic agents can be administered at essentially the same time, i.e., concurrently, or at separately staggered times, i.e., sequentially.

For example, the compound of Formula I may be administered in combination with one or more of the following additional hypoglycemic agents: insulin; biguanidines such as metformin or buformin; sulfonylureas such as acetohexamide, chlorpropamide, tolazamide, tolbutamide, glyburide, glypizide or glyclazide; thiazolidinediones such as troglitazone;  $\alpha$ -glycosidase inhibitors such as acarbose or miglatol; or B<sub>3</sub> adrenoreceptor agonists such as CL-316, 243.


The compound of Formula I is preferably administered with a biguanidine, in particular, metformin.


The compounds of Formula I contain at least three aromatic or hetero-aromatic rings, which may be designated as shown in Formula II below, and for which their substitution pattern along the chain with respect to each other also is shown below.





A preferred aspect of the compounds of Formula II, is a compound wherein  is selected from quinolinyl, benzothiophenyl, benzoimidazolyl, quinazolinyl, benzothiazolyl, quinoxalinyl, naphthyl,


pyridyl, 1H-indazolyl, 1,2,3,4-tetrahydroquinoliny, benzofuranyl, thienyl, or indolyl, and one end of the


linker, Linker I, is attached to  preferably at the 2-position of the ring moiety.


Another aspect of the compounds of Formula II is a compound wherein  is a 6-

membered aryl or heteroaryl group and Linker I and Linker II are attached to  at positions 1,2-,  
1,3-, or 1,4- to each other.


Another aspect of the compounds of Formula II is a compound wherein  is a naphthyl

group, Linker I and Linker II are attached to  at positions 1,4-,  
or 2,4- to each other on the naphthyl moiety.


Another aspect of the compounds of Formula II, is a compound wherein  is 6-  
membered aryl or heteroaryl, and has a preferred position of attachment of Linker II and Linker III to  
Ring III at positions 1,2-, to each other.

Another aspect of the compounds of Formula II, is a compound wherein  is 6-  
membered aryl or heteroaryl, and has a preferred position of attachment of Linker II and Linker III to  
Ring III at positions 1,2-,

1,3-, to each other.




Another aspect of the compounds of Formula II, is a compound wherein  is 6-  
membered aryl or heteroaryl, and has a preferred position of attachment of Linker II and Linker III to  
Ring III at positions 1,4- to each other.

A further preferred aspect of the compound of Formula II is described by Formula V below:




where  $R_1$ ,  $R_2$ ,  $c$ ,  $d$ ,  $e$ ,  $f$ ,  $D$ ,  $E$  and  $Z$  are as defined above,  $c + d = 1-3$ , and  $R'$  and  $R''$  are ring system substituents.

5

Another aspect of this invention is a compound of the invention wherein  is optionally substituted aryl, optionally substituted azaheteroaryl, or optionally substituted fused arylheterocyclyl;  is optionally substituted aryl, optionally substituted heteroaryl, or optionally substituted fused arylheterocyclyl; and  is optionally substituted aryl, optionally substituted heteroaryl, optionally substituted fused arylheterocyclyl or optionally substituted fused arylheterocyclyl.

Another aspect of this invention is a compound of the invention wherein  $a = 1$  or  $2$ ;  $R_1$  and  $R_2$  is hydrogen;  $A$  is a chemical bond; and  $b = 0$ .

Another aspect of this invention is a compound of the invention wherein  $a = 0$ ;  $A$  is



15 ;  $R_{15}$  and  $R_{16}$  are hydrogen;  $g$  is 1, 2, or 3; and  $b = 0$ .

Another aspect of this invention is a compound of the invention wherein  $a = 0$ ;  $A$  is - $NR_{13}$ -,  $b = 1$ ,  $R_3$  and  $R_4$  are hydrogen.

Another aspect of this invention is a compound of the invention wherein  $a = 2$ ; vicinal  $R_1$  radicals taken together with the carbon atoms to which the  $R_1$  radicals are linked form an ethylene group;  $R_2$  is hydrogen;  $A$  is a chemical bond; and  $b = 0$ .

20

Another aspect of this invention is a compound of the invention wherein  $a = 1, 2$  or  $3$ ;  $R_1$  and  $R_2$  are hydrogen;  $A$  is - $O$ -; and  $b = 0$ .

Another aspect of this invention is a compound of the invention wherein  $a = 1$ ;  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  are hydrogen; A is  $-O-$ ; and  $b = 1$ .

Another aspect of this invention is a compound of the invention wherein  $c = 1$  or  $2$ ;  $R_5$  and  $R_6$  are hydrogen or alkyl; B is a chemical bond; and  $d = 0$ .

Another aspect of this invention is a compound of the invention wherein  $c = 2$ ; vicinal  $R_5$  radicals taken together with the carbon atoms to which the  $R_5$  radicals are linked form an ethylene group;  $R_6$  is hydrogen; B is a chemical bond; and  $d = 0$ .

Another aspect of this invention is a compound of the invention wherein  $c = 0$  or  $1$ ;  $R_5$  and  $R_6$  are hydrogen; B is  $-O-$ ; and  $d = 0$  or  $1$ .

Another aspect of this invention is a compound of the invention wherein  $c = 0$ ; B is  $-C(O)-$  or  $-S(O)_2-$ ;  $d = 1$  and  $R_7$  and  $R_8$  are independently hydrogen or alkyl.

Another aspect of this invention is a compound of the invention wherein  $e = 0$ ;  $f = 0$ ; D and E is a chemical bond; Z is  $R_{21}O_2SHNCO-$ , and  $R_{21}$  is phenyl.

Another aspect of this invention is a compound of the invention wherein  $e = 0$ ;  $f = 0$  or  $1$ ; D and E is a chemical bond; Z is tetrazolyl,  $NH_2CO-$  or  $-CO_2R_{21}$ ; and  $R_{21}$  is hydrogen or lower alkyl.

Another aspect of this invention is a compound of the invention wherein  $e = 0$ ;  $f = 0$  or  $1$ ; D is  $-O-$  or a chemical bond; E is a chemical bond; and Z is tetrazolyl,  $NH_2CO-$  or  $-CO_2R_{21}$ ; and  $R_{21}$  is hydrogen or lower alkyl.


Another aspect of this invention is a compound of the invention wherein  $e = 0$ ;  $f = 1$ ; D is  $-O-$  or a chemical bond; E is a chemical bond;  $R_{11}$  and  $R_{12}$  are hydrogen or alkyl; and Z is tetrazolyl,  $NH_2CO-$  or  $-CO_2R_{21}$ ; and  $R_{21}$  is hydrogen or lower alkyl.

Another aspect of this invention is a compound of the invention wherein  $e = 2$ , then vicinal  $R_9$  radicals taken together with the carbon atoms to which the  $R_9$  radicals are linked form an ethylene group;  $f = 0$ ; D and E is a chemical bond; and Z is  $-CO_2R_{21}$ ; and  $R_{21}$  is hydrogen.

Another aspect of this invention is a compound of the invention wherein  $e = 0$ ;  $f = 3$ ; D is  $-O-$ ; E is a chemical bond;  $R_{11}$  and  $R_{12}$  are hydrogen or alkyl, or at least one of  $R_{11}$  is carboxyl or alkoxycarbonyl; Z is tetrazolyl, or  $-CO_2R_{21}$ ; and  $R_{21}$  is hydrogen or lower alkyl.

Another aspect of this invention is a compound of the invention wherein  $e = 0$ ;  $f = 1, 2$ , or  $3$ ; D is  $-C(O)-$ ; E is a chemical bond;  $R_{11}$  and  $R_{12}$  are hydrogen or alkyl; Z is tetrazolyl or  $-CO_2R_{21}$ ; and  $R_{21}$  is hydrogen or lower alkyl.



A preferred aspect of this invention is a compound of the invention wherein  is an optionally substituted quinolinyl, quinoxaliny, quinazolinyl, isoquinolinyl, *N*-alkyl-quinolin-4-onyl, quinazolin-4-onyl, benzoxazolyl, benzimidazolyl, benzothiazolyl, benzofuranyl, benzothiophenyl, indolinyl oxazolyl, thiazolyl, oxadiazolyl isoxazolyl, imidazolyl, pyrazol-yl, thiadiazolyl, triazolyl, pyridyl pyrimidinyl, pyrazinyl, pyridazinyl, phenyl, or naphthalenyl group, wherein the substituent is a ring system substituent as defined herein, more preferably a substituent selected from the group consisting of phenyl, substituted-phenyl, thienyl, substituted thienyl, cycloalkyl, lower alkyl, branched alkyl, fluoro, chloro, alkoxy, aralkyloxy, trifluoromethyl and trifluoromethyloxy.

A more preferred aspect of this invention is a compound of the invention wherein



is unsubstituted quinolin-2-yl, 3-substituted quinolin-2-yl, 4-substituted quinolin-2-yl, 6-substituted quinolin-2-yl or 7 substituted quinolin-2-yl; an unsubstituted quinoxalin-2-yl, 3-substituted quinoxalin-2-yl, 6-substituted quinoxalin-2-yl or 3,6-disubstituted quinoxalin-2-yl; unsubstituted quinazolin-2-yl, 4-substituted quinazolin-2-yl or 6-substituted quinazolin-2-yl; unsubstituted isoquinolin-3-yl, 6-substituted isoquinolin-3-yl or 7-substituted isoquinolin-3-yl; 3-substituted-quinazolin-4-on-2-yl; *N*-substituted quinolin-4-on-2-yl; 2-substituted-oxazol-4-yl or 2,5 disubstituted-oxazol-4-yl; 4-substituted oxazol-2-yl or 4,5-disubstituted-oxazol-2-yl; 2-substituted thiazol-4-yl or 2,5-disubstituted thiazol-4-yl; 4-substituted thiazol-2-yl or 4,5-disubstituted-thiazol-2-yl; 5-substituted-[1,2,4]oxadiazol-3-yl; 3-substituted-[1,2,4] oxadiazol-5-yl; 5-substituted-imidazol-2-yl or 3,5-disubstituted-imidazol-2-yl; 2-substituted-imidazol-5-yl or 2,3-disubstituted-imidazol-5-yl; 3-substituted-isoxazol-5-yl; 5-substituted-isoxazol-3-yl; 5-substituted-[1,2,4] thiadiazol-3-yl; 3-substituted-[1,2,4]-thiadiazol-5-yl; 2-substituted-[1,3,4]-thiadiazol-5-yl; 2-substituted-[1,3,4]-oxadiazol-5-yl; 1-substituted-pyrazol-3-yl; 3-substituted-pyrazol-5-yl; 3-substituted-[1,2,4]-triazol-5-yl; 1-substituted-[1,2,4]-triazol-3-yl; 3-substituted pyridin-2-yl, 5-substituted pyridin-2-yl, 6-substituted pyridin-2-yl or 3,5-disubstituted pyridin-2-yl; 3-substituted pyrazin-2-yl, 5-substituted pyrazin-2-yl, 6-substituted pyrazin-2-yl or 3,5 disubstituted-pyrazin-2-yl; 5-substituted pyrimidin-2-yl or 6-substituted-pyrimidin-2-yl; 6-substituted-pyridazin-3-yl or 4,6-disubstituted-pyridazin-3-yl; unsubstituted naphthalen-2-yl, 3-substituted naphthalen-2-yl, 4-substituted naphthalen-2-yl, 6-substituted naphthalen-2-yl or 7 substituted naphthalen-2-yl; 2-substituted phenyl, 4-substituted phenyl or 2,4-disubstituted phenyl;



unsubstituted -benzothiazol-2-yl or 5-substituted-benzothiazol-2-yl; unsubstituted benzoxazol-2-yl or 5-substituted-benzoxazol-2-yl; unsubstituted -benzimidazol-2-yl or 5-substituted-benzimidazol-2-yl; unsubstituted -thiophen-2-yl, 3-substituted -thiophen-2-yl, 6-substituted -thiophen-2-yl or 3,6-disubstituted-thiophen-2-yl; unsubstituted -benzofuran-2-yl, 3-substituted-benzofuran-2-yl, 6-substituted-benzofuran-2-yl or 3,6-disubstituted-benzofuran-2-yl; 3-substituted-benzofuran-6-yl or 3,7-disubstituted-benzofuran-6-yl, wherein the substituent is a ring system substituent as defined herein, more preferably a substituent selected from the group consisting of phenyl, substituted-phenyl, thienyl, substituted thienyl, cycloalkyl, lower alkyl, branched alkyl, fluoro, chloro, alkoxy, aralkyloxy, trifluoromethyl and trifluoromethyloxy.


Another more preferred aspect of this invention is a compound of the invention wherein  $R_1$  and  $R_2$  are both H,  $a = 1$ , A is -O- and  $b = 0$ .

Another more preferred aspect of this invention is a compound of the invention wherein  $R_1$  and  $R_2$  are both H,  $a = 2$ , A is -O- and  $b = 0$ .

Another more preferred aspect of this invention is a compound of the invention wherein  $a = 0$ , A is -O- or -NR<sub>13</sub>-; R<sub>13</sub> is hydrogen or alkyl; R<sub>3</sub> and R<sub>4</sub> are both independently hydrogen; and  $b = 1$ .

Another more preferred aspect of this invention is a compound of the invention wherein  $a = 0$ , A is -O- or -NR<sub>13</sub>-; R<sub>13</sub> is hydrogen or alkyl; R<sub>3</sub> and R<sub>4</sub> are both independently hydrogen;  $b = 1$ ; and Ar1 is 3-substituted quinolin-2-yl, 4-substituted quinolin-2-yl, 6-substituted quinolin-2-yl, 7 substituted quinolin-2-yl, unsubstituted quinoxalin-2-yl, 3-substituted quinoxalin-2-yl, 6-substituted quinoxalin-2-yl, 3,6-disubstituted quinoxalin-2-yl, unsubstituted quinazolin-2-yl, 4-substituted quinazolin-2-yl, 6-substituted quinazolin-2-yl, unsubstituted isoquinolin-3-yl, 6-substituted isoquinolin-3-yl, 7-substituted isoquinolin-3-yl, 4-substituted oxazol-2-yl, 4,5-disubstituted-oxazol-2-yl, 4-substituted-thiazol-2-yl, 4,5-disubstituted-thiazol-2-yl, 5-substituted -imidazol-2-yl, 3,5-disubstituted-imidazol-2-yl, 1-substituted-pyrazol-3-yl, 3-substituted-pyrazol-5-yl, 3-substituted pyridin-2-yl, 5-substituted pyridin-2-yl, 6-substituted pyridin-2-yl or 3,5-disubstituted pyridin-2-yl, 3-substituted pyrazin-2-yl, 5-substituted pyrazin-2-yl, 6-substituted pyrazin-2-yl, 3,5 disubstituted-pyrazin-2-yl, 5-substituted pyrimidin-2-yl, 6-substituted-pyrimidin-2-yl, 6-substituted-pyridazin-3-yl, 4,6-disubstituted-pyridazin-3-yl, unsubstituted-benzothiazol-2-yl, 5-substituted-benzothiazol-2-yl, unsubstituted-benzoxazol-2-yl, 5-substituted-benzoxazol-2-yl, unsubstituted benzimidazol-2-yl, 5-substituted-benzimidazol-2-yl, 3-substituted-benzofuran-6-yl or 3,7-disubstituted-benzofuran-6-yl.





wherein



and

and are independently aryl, fused arylcycloalkenyl, fused arylcycloalkyl, fused arylheterocyclenyl, fused arylheterocyclyl, heteroaryl, fused heteroaryl cycloalkenyl, fused heteroaryl cycloalkyl, fused heteroaryl heterocyclenyl, or fused heteroaryl heterocyclyl;

 $c+d = 1 \text{ or } 2;$ 

B is  $-\text{O}-$ ;

$R_5, R_6, R_7, R_8$  are independently hydrogen:


$$e = 0;$$
$$f = 0;$$

D and E are a chemical bond;

Z is  $-\text{CO}_2\text{H}$ ;

R' and R'' are ring system substituents as defined herein, more preferably, R' is lower alkyl, halo, alkoxy, aryloxy or aralkyl; and R'' is lower alkyl or halo.

Another aspect of this invention is a compound of formula I as described by formula (Ia) below:



wherein:

a = 0-2;

b = 0-1;

A is -O- or -NR<sub>13</sub>-;

c+d = 1 or 2;

B is -O-;

R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, and R<sub>8</sub> are independently hydrogen;

R<sub>13</sub> is hydrogen, R<sub>22</sub>OC-, or alkyl;

e = 0;

f = 0;

D and E are a chemical bond;

Z is -CO<sub>2</sub>H;

R' and R'' are ring system substituents as defined herein, more preferably, R' is lower alkyl,

halo, alkoxy, aryloxy or aralkyl; and R'' is lower alkyl or halo.

A more preferred aspect of this invention is a compound of formula I as described by formula (Ia) wherein:

a = 1 or 2;

A is -O-;

b = 0;

R<sub>1</sub>, R<sub>2</sub>, R<sub>7</sub> and R<sub>8</sub> are independently hydrogen;



is optionally substituted phenyl;

c = 0;

B is -O-;

d = 1;

5 e = 0;

f = 0;

D and E are a chemical bond;

R<sup>1</sup> is hydrogen, halo or benzyloxy;

R<sup>''</sup> is lower alkyl, preferably methyl;

Z is -CO<sub>2</sub>H.

A more preferred aspect of this invention is a compound of formula I as described by formula (Ia) wherein:

a = 1 or 2;

A is -O-;

b = 0;

R<sub>1</sub>, R<sub>2</sub>, R<sub>5</sub> and R<sub>6</sub> are independently hydrogen;



is optionally substituted phenyl;

c = 1;

20 B is -O-;

d = 0;

e = 0;

f = 0;

D and E are a chemical bond;

25 R<sup>1</sup> is hydrogen, halo or benzyloxy;

R<sup>''</sup> is lower alkyl, preferably methyl;

Z is -CO<sub>2</sub>H.

A more preferred aspect of this invention is a compound of formula I as described by formula (Ia) wherein:

a = 1 or 2;

A is -O-;

b = 0;

R<sub>1</sub>, R<sub>2</sub>, R<sub>7</sub>, R<sub>8</sub>, R<sub>11</sub> and R<sub>12</sub> are independently hydrogen;



is optionally substituted phenyl;

c = 0;

B is -O-;

d = 1;

e = 0;

f = 1;

D and E are a chemical bond;

R' is halo;

R'' is lower alkyl, preferably methyl;

Z is -CO<sub>2</sub>H.

A more preferred aspect of this invention is a compound of formula I as described by formula (Ia) wherein:

a = 1;

A is -O-;

b = 0;

c = 0-1;

B is -O-;

d = 0 or 1, wherein c+d = 1 or 2;

e = 0;

f = 0;

D and E are a chemical bond;

R' is hydrogen, aralkoxy, or halo;

R'' is lower alkyl, preferably methyl;

Z is  $-\text{CO}_2\text{H}$ .

A more preferred aspect of this invention is a compound of formula I as described by formula (Ia) wherein:

a = 1;

A is  $-\text{O}-$ ;

b = 0;

c = 0;

B is  $-\text{O}-$ ;

d = 1;

e = 0;

f = 0;


D and E are a chemical bond;

R' is hydrogen;

R'' is lower alkyl;

Z is  $-\text{CO}_2\text{H}$ .

A more preferred aspect of this invention is a compound of formula I as described by formula (Ia) wherein:

 are aryl or heteroaryl;

a = 1;

A is  $-\text{O}-$ ;

b = 0;

c = 0;

B is  $-\text{O}-$ ;

d = 1;

e = 0;

f = 0;

D and E are a chemical bond;

R' is hydrogen;

R'' is lower alkyl;

Z is  $-\text{CO}_2\text{H}$ .

A more preferred aspect of this invention is a compound of formula I as described by formula (Ia) wherein:



is optionally substituted azaheteroaryl;



is optionally substituted phenyl;

a = 1;

A is  $-\text{O}-$ ;

b = 0;

c = 0;

B is  $-\text{O}-$ ;

d = 1;

e = 0;

f = 0;

D and E are a chemical bond;

R' is hydrogen;

R'' is lower alkyl;

Z is  $\text{CO}_2\text{H}$ .

A more preferred aspect of this invention is a compound of formula I as described by formula (Ia) wherein:



is optionally substituted quinoliny, or a 5-membered heteroaryl group wherein the heteroaryl group is substituted by optionally substituted phenyl or optionally substituted cyclohexyl;



is optionally substituted phenyl;

a = 1;



A is -O-;

b = 0;

c = 0;

B is -O-;

5 d = 1;

e = 0;

f = 0;


D and E are a chemical bond;








R' is hydrogen;

10 R'' is lower alkyl;


Z is CO<sub>2</sub>H.



Compounds according to the invention are selected from the group consisting of
















201-203°C





96-97°C




191-196°C












169-173°C





237-239°C









;




;




;




;




;




;




143-147°C





;



;




;





;


;






















A preferred compound according to the invention is selected from the group consisting of




101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115

10




10

A more preferred compound according to the invention is selected from the group consisting




15

A preferred compound according to the invention having PPAR $\alpha$  and PPAR $\gamma$  activity is of the formula:




5

A preferred compound according to the invention that is selective for PPAR $\alpha$  is selected from the group consisting of




A preferred compound according to the invention that is selective for PPAR $\delta$  is of the formula:



15


A preferred compound according to the invention that is selective for PPAR $\delta$  and PPAR $\gamma$  is selected from the group consisting of:




20

A preferred compound according to the invention that is selective for PPAR $\alpha$  and PPAR $\delta$  is selected from the group consisting of:






5 More preferred compounds of the invention having PPAR $\gamma$  activity have the formula:




This invention also encompasses all combinations of preferred aspects of the invention noted herein.


Compounds useful according to this invention can be prepared in segments as is common to a long chain molecule. Thus it is convenient to synthesize these molecules by employing condensation reactions at the A, B and D sites of the molecule. Compounds of Formula I can be prepared by the application or adaptation of known methods, by which is meant methods used heretofore or described in the literature. Compounds of Formula I are preparable by recognized procedures from known compounds or readily preparable intermediates. Thus, in order to prepare a compound of the below formula, where Z is generally CN or CO<sub>2</sub>R




The following reactions or combinations of reactions are employable:



Scheme 1



Scheme 2



Scheme 3

In the above schemes (1-3) a displacement reaction can be used when A, B and D are O, S or NR, and L is a leaving group such as halo, tosylate or mesylate. A base such as sodium hydride, sodium hydroxide, potassium carbonate or triethylamine may be used when A, B or D is O or S.

An alternative coupling reaction is the Mitsunobu reaction (diethylazodicarboxylate/triphenylphosphine see *Synthesis*, 1981, 1). This chemistry can be used to condense fragments when the functionality is amenable to this reaction. An example of this would be the coupling of compounds of scheme 1 where formula VI (L = OH, a > 0) and formula VII (A = O, b = 0)

Reaction temperatures are in the range of about -78 °C to 80 °C and reaction times vary from about 1 to 48 hours. The reactions are usually carried out in an inert solvent that will dissolve the reactants. Solvents include, but are not limited to N,N-dimethylformamide, acetone, acetonitrile, tetrahydrofuran.

Alternatively, the reactions shown in Schemes 1-3 can be accomplished using a fragment of the described Formula. For example, as shown in Scheme 4, a compound of Formula VI (Scheme 1) may be combined with a compound of Formula VII, where Formula VII optionally contains Ar III and Z. This notation used for Formula VII in Scheme 4 is used throughout this document and is used to generalize the described reaction. Therefore, all of the reactions of Schemes 1-3 may be accomplished as shown or by using a fragment of the described formula. In some instances, the use of a protecting group may be required when a fragment of a formula is used.



Scheme 4


One method for the preparation of compounds where Z = tetrazole is the reaction of an intermediate where Z = CN with sodium azide and ammonium chloride at an elevated temperature.

One method for the preparation of compounds where  $Z = \text{CO}_2\text{H}$ , is the hydrolysis of an intermediate where  $Z$  is  $\text{CN}$  or  $\text{CO}_2\text{R}$ . This can be accomplished under acidic or basic conditions, with the preferred method generally being sodium or potassium hydroxide in a protic solvent such as aqueous ethanol at a temperature of about  $20^\circ\text{C}$  to  $100^\circ\text{C}$ .

An alternative method for the conversion of a nitrile to a carboxylic acid is to reduce the nitrile to the corresponding aldehyde using a reducing agent such as diisobutylaluminum hydride, followed by oxidation of the aldehyde to the carboxylic acid using a reagent such as sodium chlorite, sodium dihydrogen phosphate, isobutene (See, JACS **1980**, *45*, 1176) or other standard conditions.

Another alternative method for the preparation of compounds where  $Z = \text{CO}_2\text{H}$  is the oxidation of a primary alcohol using an appropriate oxidant such as PDC in DMF,  $\text{RuCl}_3 / \text{NaIO}_4$  in 3:2:2 water acetonitrile :  $\text{CCl}_4$  or the Swern system (to produce the intermediate aldehyde then oxidation of this functionality to the carboxylic acid as described above).


Some other methods for the preparation of compounds where  $Z = \text{CO}_2\text{H}$  are shown in Scheme 5. A carboxylic acid (2) can be generated directly by halogen-metal exchange of the corresponding aromatic halide (1) with an alkyl lithium reagent such as *n*-butyllithium, followed by quenching the resulting anion with carbon dioxide. Alternatively, alkoxy-carbonylation of an aromatic bromide, iodide or triflate can be accomplished in a carbon monoxide atmosphere in the presence of a suitable alcohol (usually methanol) using a catalyst such as  $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2 / \text{Et}_3\text{NH}$ ,  $\text{Pd}(\text{Ph}_2\text{P}(\text{CH}_2)_3\text{PPh}_2)_2 / \text{Et}_3\text{N}$  or alternatively cobalt, ie.  $\text{Co}(\text{OAc})_2$ , plus a base ( $\text{NaH}$  or  $\text{K}_2\text{CO}_3$ ). The resulting benzoate (3) is then converted to the benzoic acid by hydrolysis as described above.



Scheme 5

Preparation of phenylacetic acid derivatives can be accomplished from an appropriate aryl halide or triflate as illustrated in Scheme 6. Stille coupling of this type of compound with a vinyl stannane using a palladium catalyst such as  $\text{Pd}(\text{OAc})_2$ ,  $\text{P}(\text{o-tolyl})_3$  provides an olefin, (5). Hydroboration of this

vinylbenzene derivative, followed by oxidation of the resulting primary alcohol with an oxidant, such as Jones' reagent, provides the phenylacetic acid (6).



Scheme 6

In one embodiment of the current invention, ArI can be a five membered ring heterocycle thus generating astructures of the general form shown in Figure 1.






Figure 1

In particular, the heterocycle can be a substituted thiazole, oxazole, oxadiazole, imidazole, isoxazole, pyrazole, thiadiazole or triazole. These systems can be prepared using methods known in the chemical literature (for reviews see Katritzky, A.R.; Rees, C.W. Eds. *Comprehensive Heterocyclic Chemistry*, Vol. 5; Pergamon Press (1984); Katritzky, A.R.; Rees, C.W.; Scriven, E.F.V. Eds. *Comprehensive Heterocyclic Chemistry II*; Vols 3 & 4, Pergamon Press (1996)). More specifically, oxazoles, imidazoles and thiazoles can be prepared by fusion of an amide, amidine or a thioamide, respectively, with an  $\alpha$ -halo-ketone at temperatures ranging from about 40 °C to 150 °C (Scheme 7).




Scheme 7

These reactions may be carried out neat, or in a solvent such as toluene, dichlorobenzene, or ethanol. Substituted oxazoles can also be prepared from a diazoketone and a nitrile using BF<sub>3</sub> etherate (Scheme 8), Iyata, T.; Isogami, Y. *Bull. Chem. Soc. Jpn.* **1989**, *62*, 618).




Scheme 8

1,2,4-Oxadiazoles can be prepared by reaction of a nitrile with hydroxylamine followed by condensation of the resulting hydroxy-amidine with an acid chloride in the presence of a base and heating the adduct in a solvent such as toluene or dichlorobenzene to effect ring closure. (Scheme 9, Banavara, L.M.; Beyer, T.A.; Scott, P.J.; Aldinger, C.E.; Dee, M.F.; siegel, T.W.; Zembrowsky, W.J. *J. Med. Chem.*, **1992**, *35*, 457).




Scheme 9

1,3,4-oxadiazoles are prepared (Scheme 10) by condensation of an acylhydrazide with an acid synthon (such as an ester, acid chloride acyl azide) then cyclization of the resulting diacyl-hydrazide by heating in a solvent such as benzene or ethanol with or without an acid catalyst such as sulphuric acid (for examples see Weidinger, H.; Kranz, *J. Chem. Ber.*, **1963**, *96*, 1049 and Vakula, T.R.; Srinivarsan, V.R. *Indian J. Chem.*, **1973**, *11*, 732).




Scheme 10

Substituted 1,2,4-thiadiazoles can be prepared by condensation of a thioamide with an *N,N*-dimethylamide dimethyl acetal derivative in a solvent such as benzene (Scheme 11, MacLeod, A.; Baker, R.; Freedman, S.F.; Patel, S.; Merchant, K.J.; Roe, M. Saunders, J. *J. Med. Chem.*, **1990**, 33, 2052. also Patzel, M. Liebscher, J.; Siegfried, A. Schmitz, E. *Synthesis*, **1990**, 1071) followed by reaction with an electrophilic aminating agent such as mesitylsulphonyloxyamine in methanol or dialkylloxaziridine in a solvent such as toluene.




Scheme 11

In another embodiment of this invention ArI can be a 1,3,4-thiadiazole. This system is assembled by condensation of a dithioester with an imidate ester salt in a solvent such as ethanol at a temperature between room temperature and reflux (Scheme 12. Stillings, M.R.; Welbourn, A.P.; Walter, D.S. *J. Med. Chem.*, **1986**, 29, 2280). The dithioester precursor is obtained from the corresponding Grignard reagent and carbon disulphide / MeI. The imidate ester is prepared from the corresponding nitrile by reaction with HCl gas in the presence of an appropriate alcohol.




Scheme 12

Pyrazoles can be prepared by condensation of a 1,3-diketone (Scheme 13) or a synthetic equivalent with a substituted hydrazine (for example, a  $\beta$ -aminoenone. Alberola, A.; Calvo, L.; Ortega, A.G.; Sadaba, M.L.; Sanudo, M.C.; Granda, S.G.; Rodriguez, E.G., *Heterocycles*, **1999**, *51*, 2675).



Scheme 13

Similarly, isoxazoles can be prepared by reaction of a 1,3 dicarbonyl compound with hydroxyl amine (Scheme 14. Pei, Y.; Wickham, B.O.S.; *Tetrahedron Letts*, **1993**, *34*, 7509) in a solvent such as ethanol at a temperature between  $20^\circ C$  to reflux temperature.




Scheme 14

Alternatively, isoxazoles can be prepared by condensation of a hydroxamyl chloride with an alkyne (Scheme 15, Kano, H.; Adachi, I.; Kido, R.; Hirose, K. *J. Med. Chem.* **1967**, *10*, 411) in the presence of




a base such as triethylamine. The hydroxamyl chloride unit can, in turn, be prepared from the corresponding oxime by oxidation with chlorine gas at low temperature (such as  $-60^{\circ}\text{C}$ ) in a solvent such as ether (Casanti, G. Ricca, A. *Tetrahedron Lett.*, **1967**, 4, 327).




Scheme 15

Triazoles are prepared by the Einhorn-Brunner reaction or a variant thereof (Scheme 16)



Scheme 16

In addition, 5-hydroxymethyl substituted 1,2,4-triazoles can be prepared by condensation of an imidate ester with an 2-hydroxy-acetohydrazide unit (Scheme 17. Browne, E. J.; Nunn, E. E.; Polya, J. B. *J. Chem. Soc., C* **1970**, 1515).




Scheme 17

The five membered heterocycles, so formed, can, in certain cases, be coupled directly with a fragment containing ArII using standard methodology detailed elsewhere in the description of this invention (Schemes 1-4). These methods include alkylation of metal alkoxide containing ArII with a chloromethyl substituted heterocycle, or conversely, alkylation of a hydroxyl appended heterocycle (in the presence of a base) with a chloromethyl reagent containing ArII.


In another approach to fragment condensation, the substituents on the preformed heterocycle are first modified to incorporate suitable reactive functionality then this system is coupled to a fragment

containing ArII. For example (Scheme 18), treatment of a 1,4-disubstituted imidazole with a base such as n-butyl lithium at a temperature of around -78 °C followed by alkylation of the resulting anion with an electrophile such as ethylene oxide provides the hydroxyethyl-substituted imidazole (other useful electrophiles are DMF or formaldehyde. For example see Manoharan, T.S., Brown, R.S. *J. Org. Chem.*, **1989**, *54*, 1439). This intermediate can then be coupled to an ArII fragment containing an aromatic alcohol via a Mitsunobu reaction.




Scheme 18

Another example of this general approach is shown in Scheme 19. A ring substituent such as an ester can be reduced to the corresponding alcohol using a reagent such as lithium aluminum hydride or lithium borohydride in a solvent such as THF or ether. Followed by halogenation of the resulting alcohol with a reagent system such as NCS /  $\text{Ph}_3\text{P}$ ,  $\text{Ph}_3\text{P}$  /  $\text{Br}_2$  or  $\text{PBr}_3$  (Pei, Y.; Wickham, B.O.S.; *Tetrahedron Lett.*, **1993**, *34*, 7509). The alkyl halide produced in this manner can be coupled with a nucleophilic substituent attached to ArII, using a base such as  $\text{K}_2\text{CO}_3$  in the case of an aromatic alcohol, (thiol) or NaH in the case of an aliphatic alcohol (thiol).




Scheme 19

In a third general approach to fragment condensation, Ar II can be incorporated into a precursor of the five membered heterocycle. For example (Scheme 20), amination of the 3-aryl-propionate and thionation of the resulting amide provides a suitable functionalized system for thiazole ring synthesis. Similarly formation of the thio-urea from the arylmethyl amine (path B) provides a suitable precursor for fusion with an  $\gamma$ -haloketone leading to a 2-amino-substituted thiazole (Collins, J.L.; Blanchard, S. G.; Boswell, G. E.; Charifson, P. S.; Cobb, J. E.; Henke, B. R.; Hull-Ryde, E. A.; Kazmierski, W. M.; Lake, D. H.; Leesnitzer, L. M.; Lehmann, J.; Lenhard, J. M.; Orband-Miller L.A.; Gray-Nunez, Y.; Parks, D. J.; Plunkett, K. D.; Tong, Wei-Qin. *J. Med. Chem.* **1998**, *41*, 5037).



Scheme 20



In another embodiment of this invention, ArI is a five membered heterocycle of general formula shown in Figure 2.



Provided that when A = O, N, or S then "a" is > 1


Figure 2

In particular, this heterocycle can be a pyrazole, an imidazole or a triazole. These systems can be prepared by alkylation of an *N*-unsubstituted heterocycle using a base such as sodium hydride, in a solvent such as DMF, THF, DMPU or a combination of these solvents, at or around 0 °C and an electrophile such as an alkyl halide, cyclic carbonate or an epoxide (Scheme 21).



Scheme 21

These electrophiles can incorporate ArII or the alkylation products can be further modified and coupled to a fragment containing ArII in a subsequent step as described above. For example (Scheme 22), The 3,5-disubstituted pyrazole is prepared by reacting an aldehyde with a  $\beta$ -tosylhydrazinophosphonate, using the literature procedure (Almirante, N.; Benicchio, A.; Cerri, A.; Fedrizzi, G.; Marazzi, G.; Santagostino, M. *Synlett* **1999**, 299). This intermediate can then be alkylated with sodium hydride / ethylene carbonate in DMF (for a specific example see Collins, J.L.; Blanchard, S. G.; Boswell, G. E.; Charifson, P. S.; Cobb, J. E.; Henke, B. R.; Hull-Ryde, E. A.; Kazmierski, W. M.; Lake, D. H.; Leesnitzer, L. M.; Lehmann, J.; Lenhard, J. M.; Orband-Miller L.A.; Gray-Nunez, Y.; Parks, D. J.; Plunkett, K. D.; Tong, Wei-Qin. *J. Med. Chem.* **1998**, 41, 5037). This intermediate can in turn be coupled to a fragment containing ArII by a Mitsunobu reaction as described above.



Scheme 22

In another embodiment of this invention ArI is a substituted benzene, pyridine, pyrimidine, pyrazine or pyridazine (Figure 3). These systems can be prepared by applying several of the general synthetic methods detailed elsewhere in the description of this invention.






Figure 3

More specifically (Scheme 23), treatment of the known 5-bromo-2-methyl-pyridine (Graf, *J. Prakt. Chem.*, **1932**, 133, 19.) with LDA then formaldehyde in THF at low temperature (typically around -78 °C) followed by Mitsunobu coupling of the resulting alcohol to an aromatic alcohol containing ArII gives the bromo-pyridine derivative which can be further modified to give various alkyl and aryl substituted pyridines by cross coupling with an appropriate alkyl or aryl organometallic under palladium or nickel catalysis (For general reviews see Knight, D.W. and Billington, D.C. in *Comprehensive Organic Synthesis Vol. 3*, p 413 and 481, Trost, B.M. and Fleming, I, Eds. Pergamon Press 1993).




Scheme 23

Similar procedures using the appropriate 5-bromo-2-methyl-pyrimidine (Kosolapoff, G.M.; Roy, C.H. *J. Org. Chem.*, **1961**, 26, 1895), 2-iodo-5-methyl-pyrazine (Hirshberg, A.; Spoerr, P.E.; *J. Org. Chem.*, **1961**, 26, 1907) and 3-bromo-6-methyl-pyridazine (Counotte-Potman, A.; van der Plas, H.C.; *J.*

*Heterocyclic Chem.*, **1983**, *20*, 1259) provide access to the corresponding pyrimidines, pyrazines and pyridazines respectively.

In another variant of this general class, ArI is a 3-heteroatom-substituted pyridazine. For example (Scheme 24), treatment of the known 3,6-dibromo-pyridazine with a metal alkoxide (containing ArII and derived from the corresponding alcohol and sodium hydride) in a solvent such as DMSO provides the alkoxy-substituted bromo-pyridazine. The bromide can be converted into a range of substituents as described above for pyridines. In particular, Suzuki coupling with a boronic acid in the presence of a base and a palladium catalyst provides the corresponding aryl substituted pyridazines.



Scheme 24

In another embodiment of this invention ArI can be a substituted quinoxaline (Figure 4). These systems are assembled by condensation of a 1,2-dicarbonyl compound with a 1,2 diamino-benzene (for a review, see Katritzky, A.R.; Rees, C.W.; Scriven, E.F.V. Eds. *Comprehensive Heterocyclic Chemistry II*; Vol 6 Pergamon Press (1996).





Figure 4


Functionalization of these systems and coupling to ArII can be effected using procedures described for the related pyrazines. For example (Scheme 25), condensation of 1,2-diamino-benzene with 2,3-butanedione provides the 2,3-dimethyl quinoxaline. *N*-oxidation of this intermediate with a peroxy-carboxylic acid and treatment of the product with acetyl chloride gives the 2-chloromethyl-3-methyl quinoxaline (Ahmed,

Y.; Habib, M.S.; Bakhtiari, B. Bakhtiari, *Z. J. Org. Chem.*, **1966**, *31*, 2613). This intermediate is then coupled to a fragment containing Ar II under standard conditions.



Scheme 25


In another embodiment of the current invention ArI can be a quinazoline (Scheme 26). Such systems are commonly prepared by condensation of an o-amino-benzaldehyde or o-amino-aryl-ketone derivative with an acid chloride followed by heating with ammonia. For example, condensation of o-amino-benzaldehyde with chloroacetyl chloride in the presence of pyridine followed by reaction of the product with ethanolic ammonia at room temperature (Armarego, W.L.F.; Smith, J. I. C. *J. Chem. Soc.*, C, **1966**, 234) provides a 2-chloromethyl substituted quinazoline which can be coupled to a fragment containing ArII as described above.



Scheme 26


The related quinazolin-4-one ring system (Scheme 27) can be prepared by condensation of an o-amino-benzonitrile and an acid chloride followed by ring closure using a reagent such as urea hydrogen peroxide in the presence of a base such as potassium carbonate (Bavetsias, V. *Synth. Commun.* **1998**, *28*, 4547). In another variant of the quinazoline system, 4-heteroatom substituted quinazolines can be prepared by condensation of an amino-benzonitrile with chloroacetonitrile in the presence of an acid such as HCl or HBr (Chhabria, M. T.; Shishoo, C. J. *Heterocycles* **1999**, *51*, 2723.). The resulting system can be coupled to ArII as described above. The 4-halo-substituent can then be modified by nucleophilic displacement by a metal alkoxide in a solvent such as DMSO.






Scheme 27

A general reaction for the preparation of reagents such compounds of formulas VI, IX, X, XIII and XIV (schemes 1-3) is shown in Scheme 28. Halogenation of a methyl substituted aromatic with a reagent such as N-bromosuccinimide under free radical conditions provides the halomethyl substituted aromatic reagents.




Scheme 28

An alternative preparation of certain alkylating reagents is shown in Scheme 29. For example, substituted 2-chloromethyl quinoline derivatives can be prepared using a two step procedure (See, J. Med. Chem. **1991**, 34, 3224). Oxidation of the nitrogen to form the N-oxide can be achieved with an oxidant such as m-chloroperbenzoic acid or hydrogen peroxide. Reaction of the N-oxide with a reagent such as toluene sulfonylchloride at elevated temperatures can produce the target chloromethyl derivative. This chemistry can also be extended to 2-picoline derivatives where the 6-position is non-hydrogen.



Scheme 29

Preparation of reagents which can be used as alkylating agents of formula VI (scheme 1) are shown in Scheme 30. With the quinoxaline ring system the use of trichloroisocyanuric acid (TCC) can produce the corresponding chloromethyl analog (See, Chem. Ber. **1987**, 120, 649).



Scheme 30

In a particular embodiment of this invention, B of Formula I can be an amide linker of either of the general forms shown in Figure 5. Compounds of this formula can be prepared from a carboxylic acid fragment and an amine fragment using standard peptide coupling reagents. They can also be prepared by reaction with an activated carboxylic acid derivative such as, but not limited to, an acid chloride or anhydride in combination with an amine fragment in the presence of a suitable base such as triethylamine. It should be clear that essentially the same procedures can be used in the case where group A of Formula I is an amide using the appropriate carboxylic acid and amine fragments.





Figure 5


More specifically, the 2-aminomethyl-6-substituted-benzoic acid system (7) can be prepared using the chemistry shown in Scheme 31. Selective reduction of the substituted phthalic anhydride (9) with a

sterically hindered lithium trialkylborohydride, such as L-selectride, provides lactone (10) regioselectively (See Krishnamurthy, *Heterocycles*, **1982**, *18*, 4525). Reaction of this lactone with potassium phthalimide provides the protected amine according to the procedure of Bornstein, *Org.Syn. Collective Vol IV*, **1963**, 810. Removal of the phthalimide protecting group using standard deprotection conditions with hydrazine can provide the amino acid (7).




Scheme 31

1,2-Carboxylic acid-half esters such as 12 (Scheme 32) are precursors to amide linked structures (Figure 5) in which the ArIII containing fragment is the acyl donor. These systems can be prepared in several ways: Alcoholysis of the phthalic anhydride (9) can provide selectively isomer (11) plus smaller amounts of isomer (12). Esterification to provide the diester (13) can be accomplished using a variety of conditions, such as Fisher esterification. Hydrolysis of the diester can provide the regioisomer (12) as the major isomer in addition to isomer (11).




Scheme 32

Phthalic anhydride derivatives such as (9) can, in turn, be prepared from the corresponding diacid (14) as shown in Scheme 33 using dehydrating conditions such as, but not limited to, hot acetic anhydride.




Scheme 33

In one embodiment of this invention,  $\text{ArIII}-(\text{CR}_9\text{R}_{10})_2-\text{D}-(\text{CR}_{11}\text{R}_{12})_2-\text{E}-\text{Z}$  taken together constitutes a substituted benzoic acid. A useful sequence of reactions for constructing this kind of system is shown in Scheme 34. A lactone (obtained as described in Scheme 31) can be heated with hydrobromic acid to provide the bromomethyl carboxylic acid. The carboxylate can be esterified by preparing the acid chloride, followed by reaction with an alcohol to provide an intermediate bromide which can be used as outlined in Scheme 2, Formula XIII.




Scheme 34

An alternative benzoate substitution pattern can be accessed using the Alder-Rickert reaction as shown in Scheme 35 (See, *J. Org. Chem.* **1995**, 60, 560). A silyloxydiene can be formed from an enone using a strong base such as LDA and trapping the enolate with a silylating reagent. Heating this diene with an acetylenedicarboxylate at elevated temperatures can then yield the Alder-Rickert product. Alkylation of the phenolic hydroxyl under standard conditions (using the alkylating reagent  $\text{R}'\text{-L}$ , where L is a leaving group) followed by saponification of the diester can provide a diacid intermediate which can be manipulated according to the chemistry described in figures 31-34 to provide useful intermediates for the preparation of compounds of Formula I.




Scheme 35

Another particular embodiment of this invention is one in which the substituted benzoic acid moiety described above has a 6-alkyl-2-alkoxy substituent pattern. The preparation of this kind of system is illustrated using the 6-methyl derivative shown in Scheme 36 (See Hamada, *Tetrahedron*, **1991**, *47*, 8635). Ethyl acetoacetate and crotonaldehyde can be condensed to form the cyclic  $\beta$ -ketoester. Subsequent lithium chloride / cupric chloride mediated aromatization of this intermediate can be accomplished at elevated temperatures to provide the target salicylate ester. The phenolic hydroxyl of this system can be further derivatized by alkylation as outlined elsewhere in the description of this invention.



Scheme 36


Additionally, 6-alkyl-2-alkoxy benzoate systems can be prepared by aromatic nucleophilic substitution of a 2-fluorobenzaldehyde, at elevated temperatures, with an alkoxide (Scheme 37) to produce a 2-alkoxy-benzaldehyde. Oxidation of the aldehyde to the acid can then be accomplished using conditions such as sodium chlorite, sodium dihydrogen phosphate, isobutene (See, *JACS* **1980**, *45*, 1176).



Scheme 37


In another embodiment of this invention, B of Formula I can be a sulfur atom forming a thioether linkage (Scheme 2, Formulas (XI) and (XII)). This type of system can be prepared by standard thiol alkylation using a suitable base (such as sodium or potassium carbonate, hydroxide, hydride or an amine such as triethylamine) to form an anion of the thiol and then reaction of this species with an appropriate electrophile such as an alkyl halide or sulphonate ester. Similarly, the groups A and D of Formula I can also be, independently, a sulfur atom. It should be clear that the same transformations described in the Schemes below can be applied to compounds of Formulas (VII, VIII, XI, XII and XVI; schemes 1-3).

Aromatic thiols can be prepared from the corresponding phenols. For example, preparation of a 2-thiobenzoate (10) from the salicylate (7), can be carried out as shown in Scheme 38. (See, Guise J. Chem Soc., Perkin Trans. 1, **1982**, 8, 1637). The thionocarbamate (8) can be obtained from the corresponding phenol (7) using a thiocarbamoyl chloride. Pyrolysis of (8) (>300 °C) can yield the rearrangement product (9) which upon hydrolysis can yield the thiol (10).



Scheme 38

Another useful ring substituent transformation is the conversion of an aniline to an aromatic thiol. As shown in Scheme 39, diazotization of an aniline such as (11) is followed by conversion of the diazonium salt (12) to a disulfide (13) using sodium sulfide. Reduction of the disulfide with zinc/acetic acid can yield the thiol (14) (See, Guise J. Chem Soc., Perkin Trans. 1, **1982**, 8, 1637).



Scheme 39


In a particular embodiment of this invention, ArIII can be a halo-substituted aromatic. Synthesis of a particularly relevant system is outlined in scheme 40. Regioselective halogenation of a 2,6-disubstituted-phenol to provide the 4-halophenol system can be accomplished with a halogenating reagent such as sulfonyl chloride (See, J. Het. Chem. **1989**, 26, 1547). The phenolic hydroxyl group can be further derivitized as outlined elsewhere in the description of this invention.



Scheme 40


An alternative method for the preparation of a halo (or alkoxy)-substituted benzoate is shown in Scheme 41. An aniline is first converted to its diazonium salt using nitrous acid, followed by transformation to the corresponding nitrile using a reagent such as cuprous cyanide (See Chem. Ber. **1983**, 116, 1183). The cyano group is then hydrolyzed to the acid (see, Fuson, JACS **1941**, 63, 1679). The acid may then be protected as an ester to allow further derivitization of the system as outlined elsewhere in the description of this invention.

X = halogen or alkoxy



Scheme 41

An ortho-halo-benzoate can be prepared similarly, by diazotization of an o-carboxy aniline, followed by reaction with a copper halide (Scheme 42).



Scheme 42

In another embodiment of this invention, ArIII is a benzofuranyl or dihydrobenzofuranyl carboxylic acid as illustrated in Figure 6






Figure 6

Benzofuran-2-carboxylate derivatives can be prepared as shown in Scheme 43 by cyclization of the appropriately substituted 2-carbonyl-phenoxyacetate under basic conditions. Reduction of the resulting benzofuran to the corresponding 2,3-dihydro-benzofuran can be accomplished using sodium mercury amalgam under basic conditions. (See, *J. Med. Chem.* **1984**, 27, 570).



Scheme 43

An alternative synthesis of the 2,3-dihydrobenzofuran-2-carboxylate ring system is shown in Scheme 44 (See, *J. Med. Chem.* **1981**, 865). Claisen rearrangement of a substituted allyl-phenyl ether at an elevated temperature such as 250 °C either neat or in a solvent such as dimethylaniline, can provide the o-allyl-phenol. Peracid oxidation of this intermediate provides the 2-hydroxymethyl-2,3-dihydrobenzofuran, which can be further oxidized to the carboxylic acid using a variety of oxidizing agents such as Jones' reagent.



Scheme 44

In another embodiment of this invention "A" can be an imidazolidin-2-one, a tetrahydropyrimidin-2-one an imidazoline-2,4-dione, or a tetrahydropyrimidin-2,4-dione (Figure 7).





Figure 7



These systems are prepared from an amine containing ArI by sequential acylation, aminolysis, ring closure and reduction, as illustrated in Scheme 45 (For examples see Kitazaki, T.; asaka, A.; Tamura, N.; Matsushita, Y.; Hosono, H.; Hayashi, R.; Okonogi, K.; Itoh, K. *Chem. Pharm. Bull.*, **1999**, *47*, 351 and Basha, A.; *Tetrahedron Lett.*, **1988**, *29*, 2525). Coupling to ArII can be effected by derivitization of the cyclic urea N by treatment with a base such as NaH in THF at around 0 °C then alkylation of the resulting anion with an electrophile such as an alkyl bromide / triflate containing ArII).



Scheme 45

In another embodiment of this invention ArII is a six membered ring aromatic of general form substituted shown in Figure 8. In particular, ArII is a substituted benzene, pyridine, pyrimidine, pyridazine or pyrazine.





Figure 8


In principle, appropriately functionalized ring systems of this kind can be prepared by functionalization of specific precursors followed by ring synthesis or by derivitization of a preformed ring system. There are numerous approaches to the synthesis and functionalization of the aforementioned cyclic frameworks in the chemical literature (for examples, see (a) Katritzky, A.R.; Rees, C.W.; Scriven,

E.F.V. Eds. *Comprehensive Heterocyclic Chemistry II*, Vol. 5 and Vol. 6. Elsevier Science 1996 and references therein). For example, (Scheme 46) alkylation of methyl glycolate with an alkyl halide containing ArI using a base such as sodium hydride in a solvent such as THF or DMSO provides the ester. Claisen condensation of this ester with *t*-butyl acetate at low temperature (typically below -15 °C) using a base such as LDA in THF provides the keto ester intermediate. This is reacted with formamidine acetate in the presence of a base such as sodium methoxide in methanol to give the pyrimidinone (Butters, M. J. *Heterocyclic Chem.*, **1992**, 29, 1369). This type of substituted aromatic system can be further functionalized to incorporate ArI as described elsewhere in the description of this invention.




Scheme 46

In certain cases, ArII (in Figure 8) can be assembled by ring transformation of another heterocycle, for example, treatment of the known 4-bromo-2-methoxy-furan (Scheme 47, Marini-Bettolo, R.; Flecker, P.; Tsai, T. Y. R.; Wiesner, K. *Can. J. Chem.* **1981**, 59, 1403) with an alkyl lithium at low temperature and reaction of this anion with an electrophile containing ArIII (such as a bromide, aldehyde, epoxide) provides the 4-substituted furan. Oxidative cleavage of this intermediate with dioxirane followed by treatment with hydrazine provides the pyridazinone which can be further modified to incorporate ArI as illustrated elsewhere in the description of this invention.




Scheme 47

A particularly useful protocol with regard to functionalization of heterocycles involves Mitsunobu etherification of hydroxyl substituted heterocycles (or keto-tautomers) such as outlined in Scheme 48. Treatment of the known 6-bromo-pyridin-2-one (Wibaut, J.P.; Waayman, P.W.; Vandijk, J. *Rec. Trav. Chim. Pays-Bas*, **1940**, *59*, 202) with an alcohol containing an ArI (or ArIII) under Mitsunobu's conditions provides the corresponding bromo-substituted pyridyl ether. (for typical procedures see Mitsunobu, O., *Synthesis*, **1981**, 1).



Scheme 48


The heterocyclic bromide, so formed, can be further functionalized in a number of ways. For example, coupling with a vinyl stannane can be effected under palladium (0) catalysis to provide systems with alkenyl linkers (Scheme 49).



Scheme 49

The choice of catalyst and reaction temperature for this transformation depends on the substrate employed but is most commonly tetrakis(triphenylphosphine) palladium, bis(triphenylphosphine) palladium chloride, 1,1'-bis(diphenylphosphino)ferrocene / bis-dibenzylideneacetone palladium or 1,2-bis(diphenylphosphino)ethane / bis(acetonitrile)dichloropalladium at a temperature between 50 and 150 °C. Suitable solvents include DMF, DMPU, HMPA, DMSO, toluene, and DME. (for examples see Farina, V. Krishnamurthy, V.; Scott, W.J. *Organic Reactions*, 1997, 50, 1). Reduction of the olefin using, for example, Wilkinson's catalyst in a solvent such as toluene, THF or an alcohol at a temperature between about 20 and 80 °C provides the corresponding alkane linked system.


In certain heterocyclic systems in which a bromide or chloride is situated ortho or para to a ring nitrogen, the halogen can be readily displaced by an alcohol (in the presence of base such as sodium hydride in a solvent such as Toluene, DMSO, THF, DMPU or HMPA) at or above room temperature (For examples see Kelly, T.R. et al. *J. Amer. Chem. Soc.*, 1994, 116, 3657 and Newcome, G.R. et al. *J. Org. Chem.*, 1977, 42, 1500). For example, alcoholysis of a 2,4-dichloro-pyrimidine (Scheme 50) using a controlled stoichiometric amount of an alcohol reagent containing ArI (or ArIII) provides the alkoxy substituted-bromo-pyrimidine. Subsequent reaction of this product (generally above room temperature) with a further equivalent of another alcohol containing ArIII (or ArI) provides the unsymmetrically dialkoxy-substituted heterocycle. Since the 4-position of the dichloro-pyrimidine is generally displaced first, the order in which the alkoxy substituents are introduced will dictate their orientation in the product.



Scheme 50

Similar procedures using 2,6-dibromo-pyridine or 2,6-dibromo-pyridazine provides the corresponding dialkoxy-substituted pyridines and pyridazines.


A simple alkoxy group positioned ortho to a nitrogen in these heterocyclic systems can be hydrolyzed to the corresponding hydroxy substituent using aqueous hydrochloric acid at a temperature normally between room temperature and reflux. (Scheme 51).



Scheme 51

For example (Scheme 51), treatment of the 2-methoxy-6-alkyl-substituted pyridine derivative with hydrochloric acid provides the 6-alkyl substituted pyridin-2-one. This intermediate, in turn, can be further derivatized to the corresponding 2-alkoxy or 2-alkenyl substituted systems as detailed elsewhere in the description of this invention.

A methyl, methylene or methine group positioned ortho to a ring nitrogen in these heterocyclic systems can be deprotonated with a base such as an alkyl lithium or LDA in a solvent such as THF, ether or HMPA, generally at low temperature (below 0°C) and the resulting anion reacted with electrophiles such as aldehydes, epoxides, alkyl halides or  $\alpha,\beta$ -unsaturated carbonyl compounds to provide a variety of functionalized heterocycles.



Scheme 52

For example (Scheme 52) the 2-alkoxy-4-methyl-pyrimidine is treated, sequentially, with LDA and an aldehyde at  $-78^{\circ}\text{C}$  to give the hydroxy substituted adduct. Subsequent dehydration of this intermediate with martin sulfurane in a solvent such as dichloromethane at ambient temperature followed by hydrogenation of the resulting olefin provides the 4-ArI-containing-alkyl-2-alkoxy-pyrimidine. Similar procedures applied to 2-chloro-6-methyl-pyrazine (Karmas, G.; Spoerri, P.E.; *J. Amer. Chem. Soc.*, 1952, 74, 1580) leads to the corresponding pyrazine.

In another embodiment of this invention, A can be an amide thus generating compounds of the formulas shown in Figure 9.






Figure 9

The preparation of an illustrative example within this series is shown in scheme 53. A hydroxy aldehyde can be reacted with a bromoalkyl-ester to provide an aldehyde-ester intermediate. Reductive amination of the aldehyde followed by acylation can provide the amide.




Scheme 53

Furthermore, compounds of the invention may be easily synthesized by solid phase methods, as illustrated in Schemes 54 and 55, using inputs (XII) - (XVII) listed in Table 1.










































Scheme 54











Scheme 55














82  
Table 1


 XII	$R_{15}NH_2$ XIII	 XIV	 XV	$R_{15}COCl$ XVI	$R_{15}NCO$ XVII	 XVIII
						
						
						
						
						

83


				
				

$\text{CH}_3\text{C}(\text{OH})(\text{Ad})\text{CH}_2\text{OH}$ <b>XII</b>	$\text{R}_{13}\text{NH}_2$ <b>XIII</b>	$\text{R}_1$ $\text{Ad}$ <b>XIV</b>	$\text{R}_2$ $\text{CH}_2\text{NH}_2$ <b>XV</b>	$\text{R}_{15}\text{COCl}$ <b>XVI</b>	$\text{R}_{15}\text{NCO}$ <b>XVII</b>	$\text{X}$ $\text{R}_7$ $\text{R}_8$ $\text{Ar}$ $\text{R}_9$ $\text{R}_{10}$ $\text{R}_{11}$ $\text{R}_{12}$ <b>XVIII</b>
						
						
						
						
						

An additional exemplification of the amide linker is shown in Scheme 56. Reaction of an activated carboxylic acid derivative such as, but not limited to, an acid chloride or anhydride with an amine of general formula (15) and a suitable base such as triethylamine provides the amide (16). More explicit exemplification is shown in Scheme 57. Carboxylic acid (17) is activated with oxalyl chloride to provide the acid chloride and then 2-amino-6-methylbenzoic acid (18), is added to provide the amide (19). Alternatively, 2-aminomethyl-6-methylbenzoic acid (20) can be used to provide the amide (21).



Scheme 56



Scheme 57

The preparation of 2-aminomethyl-6-methylbenzoic acid, (20), can be accomplished using the chemistry shown in Scheme 31 (X = Me, R' = H).

Compounds useful according to the invention may also be prepared by the application or adaptation of known methods, by which is meant methods used heretofore or described in the literature, for example those described by R. C. Larock in *Comprehensive Organic Transformations*, VCH publishers, 1989.

In the reactions described hereinafter, it may be necessary to protect reactive functional groups, for example hydroxy, amino, imino, thio or carboxy groups, where these are desired in the final product, to avoid their unwanted participation in the reactions. Conventional protecting groups may be used in accordance with standard practice, for examples see T.W. Green and P.G.M. Wuts in "Protective Groups in Organic Chemistry" John Wiley and Sons, 1991; J. F. W. McOmie in "Protective Groups in Organic Chemistry" Plenum Press, 1973.

According to a further feature of the present invention, compounds useful according to the invention may be prepared by interconversion of other compounds of the invention.

A compound of the invention including a group containing one or more nitrogen ring atoms, preferably imine (=N-), may be converted to the corresponding compound wherein one or more nitrogen ring atom of the group is oxidized to an N-oxide, preferably by reacting with a peracid, for example peracetic acid in acetic acid or m-chloroperoxybenzoic acid in an inert solvent such as dichloromethane, at a temperature from about room temperature to reflux, preferably at elevated temperature.

The products of this invention may be obtained as racemic mixtures of their dextro and levorotatory isomers since at least one asymmetric carbon atom may be present. When two asymmetric carbon atoms are present, the product may exist as a mixtures of diastereomers based on syn and anti configurations. These diastereomers may be separated by fractional crystallization. Each diastereomer may then be resolved into dextro and levorotatory optical isomers by conventional methods.

It will also be apparent to those skilled in the art that certain compounds of Formula I may exhibit geometrical isomerism. Geometrical isomers include the cis and trans forms of compounds of the invention having an alkenyl moiety. The present invention comprises the individual geometrical isomers and stereoisomers and mixtures thereof.

Such isomers can be separated from their mixtures, by the application or adaptation of known methods, for example chromatographic techniques and recrystallization techniques, or they are separately prepared from the appropriate isomers of their intermediates, for example by the application or adaptation of methods described herein.

Resolution may best be carried out in the intermediate stage where it is convenient to combine the racemic compound with an optically active compound by salt formation, ester formation, or amide formation to form two diastereomeric products. If an acid is added to an optically active base, then two diastereomeric salts are produced which possesses different properties and different solubilities and can be separated by fractional crystallization. When the salts have been completely separated by repeated crystallization, the base is split off by acid hydrolysis and enantiomerically purified acids are obtained.

Compounds useful according to the invention are useful in the form of the free base or acid or in the form of a pharmaceutically acceptable salt thereof. All forms are within the scope of the invention.

Where a compound useful according to the invention is substituted with a basic moiety, acid addition salts are formed and are simply a more convenient form for use; in practice, use of the salt form inherently amounts to use of the free base form. The acids which can be used to prepare the acid addition salts include preferably those which produce, when combined with the free base, pharmaceutically acceptable salts, that is, salts whose anions are non-toxic to the patient in pharmaceutical doses of the salts, so that the beneficial pharmaceutical effects of these compounds in the free base are not vitiated by side effects ascribable to the anions. Although pharmaceutically acceptable salts of said basic compounds are preferred, all acid addition salts are useful as sources of the free base form even if the particular salt, per se, is desired only as an intermediate product as, for example, when the salt is formed only for purposes of purification, and identification, or when it is used as an intermediate in preparing a pharmaceutically acceptable salt by ion exchange procedures. Pharmaceutically acceptable salts useful within the scope of the invention are those derived from the following acids: mineral acids such as hydrochloric acid, trifluoroacetic acid, sulfuric acid, phosphoric acid and sulfamic acid; and organic acids such as acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, cyclohexylsulfamic acid, quinic acid, and the like. The corresponding acid addition salts comprise the following: hydrohalides, e.g. hydrochloride and hydrobromide, trifluoroacetate, sulfate, phosphate, nitrate, sulfamate, acetate, citrate, lactate, tartrate, malonate, oxalate, salicylate, propionate, succinate, fumarate, maleate, methylene-bis- $\beta$ -hydroxynaphthoates, gentisates, mesylates, isothionates, di-p-toluoyletartrates, methanesulfonates, ethanesulfonates, benzenesulfonates, p-toluenesulfonates, cyclohexylsulfamate and quinate, respectively.

The acid addition salts of the compounds useful according to the invention are prepared by reaction of the free base with the appropriate acid, by the application or adaptation of known methods. For example, the acid addition salts of the compounds of this invention are prepared either by dissolving the free base in aqueous or aqueous-alcohol solution or other suitable solvents containing the appropriate

acid and isolating the salt by evaporating the solution, or by reacting the free base and acid in an organic solvent, in which case the salt separates directly or can be obtained by concentration of the solution.

The compounds useful according to the invention may be regenerated from the acid addition salts by the application or adaptation of known methods. For example, parent compounds useful according to the invention can be regenerated from their acid addition salts by treatment with an alkali, e.g., aqueous sodium bicarbonate solution or aqueous ammonia solution.

Where the compound useful according to the invention is substituted with an acidic moiety, base addition salts may be formed and are simply a more convenient form for use; in practice, use of the salt form inherently amounts to use of the free acid form. The bases which can be used to prepare the base addition salts include preferably those which produce, when combined with the free acid, pharmaceutically acceptable salts, that is, salts whose cations are non-toxic to the animal organism in pharmaceutical doses of the salts, so that the beneficial pharmaceutical effects on the activity of the compounds of the present invention in the free acid are not vitiated by side effects ascribable to the cations. Pharmaceutically acceptable salts useful according to the invention, include for example alkali and alkaline earth metal salts, including those derived from the following bases: sodium hydride, sodium hydroxide, potassium hydroxide, calcium hydroxide, aluminum hydroxide, lithium hydroxide, magnesium hydroxide, zinc hydroxide, ammonia, ethylenediamine, N-methyl-glucamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chloroprocaine, diethanolamine, procaine, diethylamine, N-benzylphenethylamine, piperazine, tris(hydroxymethyl)aminomethane, tetramethylammonium hydroxide, and the like.

Metal salts of compounds useful according to the present invention may be obtained by contacting a hydride, hydroxide, carbonate or similar reactive compound of the chosen metal in an aqueous or organic solvent with the free acid form of the compound. The aqueous solvent employed may be water or it may be a mixture of water with an organic solvent, preferably an alcohol such as methanol or ethanol, a ketone such as acetone, an aliphatic ether such as tetrahydrofuran, or an ester such as ethyl acetate. Such reactions are normally conducted at ambient temperature but they may, if desired, be conducted with heating.

Amine salts of compounds useful according to the present invention may be obtained by contacting an amine in an aqueous or organic solvent with the free acid form of the compound. Suitable aqueous solvents include water and mixtures of water with alcohols such as methanol or ethanol, ethers such as tetrahydrofuran, nitriles such as acetonitrile, or ketones such as acetone. Amino acid salts may be similarly prepared.

The base addition salts of the compounds useful according to the invention can be regenerated from the salts by the application or adaptation of known methods. For example, parent compounds useful according to the invention can be regenerated from their base addition salts by treatment with an acid, e.g. hydrochloric acid.

Salt forms useful according to the invention also include compounds having a quarternarized nitrogen. The quarternarized salts are formed by methods such as by alkylation of  $sp^3$  or  $sp^2$  hybridized nitrogen in the compounds.

As will be self-evident to those skilled in the art, some of the compounds useful according to the invention do not form stable salts. However, acid addition salts are most likely to be formed by compounds useful according to the invention having a nitrogen-containing heteroaryl group and/or wherein the compounds contain an amino group as a substituent. Preferable acid addition salts of the compounds useful according to the invention are those wherein there is not an acid labile group.

As well as being useful in themselves as active compounds, the salts of the compounds useful according to the invention are useful for the purposes of purification of the compounds, for example by exploitation of the solubility differences between the salts and the parent compounds, side products and/or starting materials by techniques well known to those skilled in the art.

Various substituents on the compounds useful according to the invention, e.g., as defined in  $R$ ,  $R_1$  and  $R_2$  can be present in the starting compounds, added to any one of the intermediates or added after formation of the final products by known methods of substitution or conversion reactions. If the substituents themselves are reactive, then the substituents can themselves be protected according to the techniques known in the art. A variety of protecting groups known in the art may be employed. Examples of many of these possible groups may be found in "Protective Groups in Organic Synthesis" by T. W. Green, John Wiley and Sons, 1981. For example, nitro groups can be added to the aromatic ring by nitration, and the nitro group then converted to other groups, such as amino, by reduction, and halo, by diazotization of the amino group and replacement of the diazo group. Acyl groups can be substituted onto the aryl groups by Friedel-Crafts acylation. The acyl groups then can be transformed to the corresponding alkyl groups by various methods, including the Wolff-Kishner reduction and Clemmensen reduction. Amino groups can be alkylated to form mono and dialkylamino groups; and mercapto and hydroxy groups can be alkylated to form corresponding ethers. Primary alcohols can be oxidized by oxidizing agents known in the art to form carboxylic acids or aldehydes, and secondary alcohols can be oxidized to



form ketones. Thus, substitution or alteration reactions can be employed to provide a variety of substituents throughout the molecule of the starting material, intermediates, or the final product.

The starting materials, intermediates and some compounds of the invention are prepared by the application or adaptation of known methods, for example methods as described in US Patent No.s 4,920,132; 4,920,131; and 5,059,610; publications Huang, Fu Chih et al , J. Med. Chem. (1991), 34(5), 1704-7; and Huang, Fu Chih et al , J. Med. Chem. (1990), 33(4), 1194-200; and the Reference Examples or their obvious chemical equivalents.

The present invention is further exemplified but not limited by the following examples, which illustrate the preparation of the compounds according to the invention.

## Example 1

Methyl 2,6-dimethyl-benzoate

To a cooled (0 °C) solution of 2,6-dimethylbenzoic acid (20.2g, 134 mmol) in dichloromethane (200 mL) is added DMF (1 mL) followed by oxalyl chloride (14 mL, 162 mmol). On completion of addition, the cold bath is removed and stirring continued for 3h. The resulting solution is concentrated under vacuum and the residue added slowly to a cooled (0 °C) solution comprising methanol (200 mL) and triethylamine (40 mL). On completion of addition, the reaction mixture is stirred for 30 min. then poured into hydrochloric acid solution (400 mL, 2N) which is then extracted with ether. The ether extract is washed with hydrochloric acid solution (1N), sodium bicarbonate solution and brine then dried over MgSO<sub>4</sub> and concentrated to give the title compound which is used without further purification. MS (EI) 164 (M)<sup>+</sup>.

## Example 2

Methyl 2-bromomethyl-6-methyl-benzoate

To a solution of methyl 2,6-dimethyl-benzoate (22.0 g, 134 mmol, example 1) in CCl<sub>4</sub> (250 mL) is added *N*-bromo-succinimide (19g, 107 mmol) followed by benzoyl peroxide (1.0 g, 4.0 mmol). The resulting solution is warmed to reflux and stirred at this temperature for 20 min. The reaction mixture is then allowed to cool before being diluted with ether (200 mL), filtered and concentrated. The residue is purified by flash chromatography (silica, 4% acetone in hexanes) to give the title compound. This product (approx. 85% purity, remainder is methyl 2,6dimethyl benzoate) is used without further purification. MS (EI) 242, 244 (M<sup>+</sup>, Br pattern).

## Example 3

3-(Quinolin-2-ylmethoxy)-phenol

2-Chloromethylquinoline hydrochloride (25.0 g, 117 mmol) and resorcinol monobenzoate (37.5 g, 175 mmol) are suspended in dimethylsulfoxide (180 mL) and are mixed with the aid of an overhead stirrer. The mixture is cooled to 15 °C and a 50% solution of sodium hydroxide (25 mL) is added slowly over 10 min with a slight exotherm. The reaction is allowed to come to room temperature and is allowed to stir overnight. The reaction is then heated to 95 °C and a

50% solution of sodium hydroxide (25 mL) is added over 10 min. After 20 min. hot water (300 mL) is added to the reaction and stirred 15 min. The reaction is hot filtered and the filtrate is cooled to provide a brick red solid which is dried *in vacuo* to provide the sodium salt pentahydrate. A portion of the salt (15.6 g, 43.0 mmol) is neutralized by heating in water (30 mL) with 1N HCl (43 mL) followed by cooling to provide a brown solid. The solid is dissolved in dichloromethane (550 mL) and methanol (14 mL), dried over magnesium sulfate, filtered and concentrated *in vacuo* to yield the title compound as a tan solid. A portion is recrystallized from ethyl acetate to provide an analytically pure sample; m.p. 152-153 °C, MS (ESI) 252 (M+H)<sup>+</sup>.

#### Example 4

##### Methyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxyethyl]-benzoate

3-(Quinolin-2-ylmethoxy)-phenol (5.2 g, 21 mmol, example 3), methyl 2-bromomethyl-6-methyl-benzoate (example 2) (5.0 g, 21 mmol) and potassium carbonate (4.4 g, 32 mmol) are combined in DMF (50 mL) and heated at 85 °C for 3h. The reaction is poured into ethyl acetate (500 mL) and washed with water (4 x 120 mL) and brine (100 mL). The solution is dried over magnesium sulfate, filtered and concentrated *in vacuo* to provide a residue which is purified by column chromatography (silica, 10 to 20% ethyl acetate in hexane) to provide the title compound. MS (ESI) 414 (M+H)<sup>+</sup>.

The following compounds are prepared using essentially the same procedure used in example 4 except using the cited phenol in place of 3-(quinolin-2-ylmethoxy)-phenol with either the methyl or isobutyl benzoate (example 2).

#### Example 4a

##### Methyl {2-methyl-6-[3-(2-quinolin-2-yl-vinyl)-phenoxyethyl]}-benzoate.

MS (ESI) 410 (M+H)<sup>+</sup>. Prepared from 3-(2-quinolin-2-yl-vinyl)-phenol (example 15).

#### Example 4b

##### Methyl (2-Methyl-6-{3-[2-(pyridin-2-yloxy)-ethoxy]-phenoxyethyl})-benzoate

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.15 (m, 1H), 7.60 (t, 1H), 7.1-7.38 (m, 4H), 6.89 (dd, 1H), 6.81 (d, 1H), 6.56 (m, 3H), 5.08 (s, 2H), 4.70 (t, 2H), 4.32 (t, 2H), 3.85 (s, 3H), 2.36 (s, 3H).

Prepared from 3-[2-(pyridin-2-yloxy)-ethoxy]-phenol (example 15a).

#### 5 Example 4c

Methyl 2-{3-[(Benzoxazol-2-yl-methyl-amino)-methyl]-phenoxy-methyl}-6-methyl-benzoate

MS (ESI) 417 (M+H)<sup>+</sup>. Prepared from 3-[(N-benzoxazol-2-yl-N-methyl-amino)-methyl]-phenol (example 10a).

10

#### Example 4d

Methyl 2-methyl-6-{3-[(methyl-quinolin-2-yl-amino)-methyl]-phenoxy-methyl}-benzoate

MS (ESI) 427 (M+H)<sup>+</sup>. Prepared from 3-[(N-methyl-N-quinolin-2-yl-amino)-methyl]-phenol (example 10b).

#### Example 4e

Isobutyl 2-methyl-6-[3-(quinolin-2-yloxy-methyl)-phenoxy-methyl]-benzoate

MS (ESI) 456 (M+H)<sup>+</sup>. Prepared from 3-(quinolin-2-yloxy-methyl)-phenol (example 15b).

#### Example 4f

Methyl 2-{3-[2-(5-ethyl-pyridin-2-yl)-ethoxy]-phenoxy-methyl}-6-methyl-benzoate

25 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.39 (bs, 1H), 7.46 (d, 1H), 7.28 (m, 2H), 7.16 (m, 3H), 6.52 (m, 3H), 5.06 (s, 2H), 4.31 (t, 2H), 3.82 (s, 3H), 3.22 (t, 2H), 2.63 (q, 2H), 2.38 (s, 3H), 1.24 (t, 3H).  
MS (ESI) 406 (M+H)<sup>+</sup>. Prepared from 3-[2-(5-ethyl-pyridin-2-yl)-ethoxy]-phenol (example 71).

#### Example 4g

Methyl 2-methyl-6-[3-(2-pyridin-2-yl-ethoxy)-phenoxy-methyl]-benzoate

30 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.56 (d, 1H), 7.62 (m, 1H), 7.28 (m, 2H), 7.16 (m, 3H), 6.52 (m, 4H), 5.06 (s, 2H), 4.34 (t, 2H), 3.82 (s, 3H), 3.25 (t, 2H), 2.38 (s, 3H). MS (ESI) 378 (M+H)<sup>+</sup>.  
Prepared from 3-(2-pyridin-2-yl-ethoxy)-phenol (example 71a).

## Example 4h

Methyl 2-[3-(Benzooxazol-2-ylaminomethyl)-phenoxy-methyl]-6-methyl-benzoate

5 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.54 (m, 1H), 7.38 (d, 1H), 7.26 (m, 3H), 7.18 (m, 2H), 7.05 (m, 1H), 6.98 (m, 2H), 6.88 (dd, 1H), 5.10 (s, 2H), 4.64 (bs, 2H), 3.80 (s, 3H), 2.37 (s, 3H). MS (ESI) 403 (M+H)<sup>+</sup>. Prepared from 3-(benzooxazol-2-ylaminomethyl)-phenol (example 10c).

## Example 4i

Methyl 2-methyl-6-[3-(pyridin-2-ylmethoxymethyl)-phenoxy-methyl]-benzoate

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.56 (d, 1H), 7.71 (m, 1H), 7.48 (d, 1H), 7.25 (m, 5H), 6.99 (m, 2H), 6.87 (dd, 1H), 5.11 (s, 2H), 4.69 (s, 2H), 4.63 (s, 2H), 3.82 (s, 3H), 2.38 (s, 3H). MS (ESI) 378 (M+H)<sup>+</sup>. Prepared from 3-(pyridin-2-ylmethoxymethyl)-phenol (example 74).

## Example 4j

Methyl 2-methyl-6-[3-(quinolin-2-ylmethoxymethyl)-phenoxy-methyl]-benzoate

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.19 (d, 1H), 8.06 (d, 1H), 7.82 (d, 1H), 7.69 (m, 2H), 7.53 (m, 1H), 7.24 (m, 4H), 7.01 (m, 2H), 6.88 (dd, 1H), 5.12 (s, 2H), 4.86 (s, 2H), 4.66 (s, 2H), 3.82 (s, 3H), 2.38 (s, 3H). MS (ESI) 428 (M+H)<sup>+</sup>. Prepared from 3-(quinolin-2-ylmethoxymethyl)-phenol (example 74a).

## Example 5

Methyl 2-methyl-6-[(3-hydroxy-phenoxy)-methyl]-benzoate

To a solution of 3-hydroxy-phenol (1.5g, 13.6 mmol) in acetonitrile (50 mL) is added methyl 2-(bromomethyl)-6-methyl-benzoate (0.82g, 3.4 mmol, example 2) followed by K<sub>2</sub>CO<sub>3</sub> (3.76g, 27.2 mmol). The resulting mixture is heated to 50 °C and stirred at this temperature for 90 min. then cooled, filtered and the filtrate concentrated under vacuum. The residue is purified by flash chromatography (silica, 5% ethyl acetate in dichloromethane) to give the title compound as a white solid. MS (EI) 272 (M<sup>+</sup>).

## Example 6

Methyl 2-methyl-6-[3-(2-phenyl-oxazol-4-ylmethoxy)-phenoxy-methyl]-benzoate

To a solution of 4-chloromethyl-2-phenyl-oxazole (100mg, 0.5 mmol, example 19) in DMF (2 mL) is added methyl 2-methyl-6-[(3-hydroxy-phenoxy)-methyl]-benzoate (136 mg, 0.5 mmol, example 5) followed by  $K_2CO_3$  (75 mg, 0.54 mmol). The resulting mixture is heated to 60 °C and stirred at this temperature for 8 h. This mixture is then cooled to room temperature, diluted with ether, washed with water and brine, dried over  $MgSO_4$  and concentrated. The residue is purified by flash chromatography (silica, 20% ethyl acetate in hexanes) to give the title compound. MS (ESI) 429 (M+H)<sup>+</sup>.

The following compounds are prepared using essentially the same procedure used in Example 6 except using the cited alkyl halide in place of 4-chloromethyl-2-phenyl-oxazole with either methyl (ethyl or isobutyl) 2-methyl-6-[(3-hydroxy-phenoxy)-methyl]-benzoate (example 5).

## Example 6a

Methyl 2-methyl-6-[3-(2-phenyl-thiazol-4-ylmethoxy)-phenoxy-methyl]-benzoate

MS (ESI) 446 (M+H)<sup>+</sup>. Prepared from 4-chloromethyl-2-phenyl-thiazole (example 20).

## Example 6b

Methyl 2-[3-(3,5-dimethyl-isoxazol-4-ylmethoxy)-phenoxy-methyl]-6-methyl-benzoate

MS (ESI) 382 (M+H)<sup>+</sup>. Prepared from 3,5-dimethyl-isoxazol-4-ylmethyl chloride.

## Example 6c

Methyl 2-methyl-6-[3-(5-phenyl-[1,2,4]oxadiazol-3-ylmethoxy)-phenoxy-methyl]-benzoate

MS (ESI) 431 (M+H)<sup>+</sup>. Prepared from 5-phenyl-[1,2,4]oxadiazol-3-ylmethyl chloride.

## Example 6d

Methyl 2-[3-(2,5-dimethyl-benzyl-oxo)-phenoxy-methyl]-6-methyl-benzoate

MS (ESI) 391 (M+H)<sup>+</sup>. Prepared from 2,5-dimethyl-benzyl chloride.

## Example 6e

Methyl 2-[3-(2,4-dichloro-benzyloxy)-phenoxy-methyl]-6-methyl-benzoate

5 MS (ESI) 431 (M+H)<sup>+</sup>. Prepared from 2,4-dichloro-benzyl chloride.

## Example 6f

Methyl 2-[3-(5-tert-butyl-[1,2,4]oxadiazol-3-ylmethoxy)-phenoxy-methyl]-6-methyl-benzoate

10 MS (ESI) 411 (M+H)<sup>+</sup>. Prepared from 5-tert-butyl-[1,2,4]oxadiazol-3-ylmethyl chloride.

## Example 6g

Methyl 2-[3-[(2,6-dichloro-phenyl)-5-methyl-isoxazol-4-ylmethoxy]-phenoxy-methyl]-6-methyl-benzoate

MS (ESI) 512 (M+H)<sup>+</sup>. Prepared from (3-(2,6-dichloro-phenyl)-5-methyl-isoxazol-4-yl)-methyl chloride.

## Example 6h

Methyl 2-methyl-6-[3-(2,4,5-trimethyl-benzyloxy)-phenoxy-methyl]-benzoate

MS (ESI) 405 (M+H)<sup>+</sup>. Prepared from 2,4,5-trimethyl-benzyl chloride.

## Example 6i

Methyl 2-methyl-6-[3-(3-methyl-naphthalen-2-ylmethoxy)-phenoxy-methyl]-benzoate

MS (ESI) 427 (M+H)<sup>+</sup>. Prepared from (3-methyl-naphthalen-2-yl)-methyl bromide.

## Example 6j

Methyl 2-[3-(5-acetyl-2-methoxy-benzyloxy)-phenoxy-methyl]-6-methyl-benzoate

MS (ESI) 435 (M+H)<sup>+</sup>. Prepared from 5-acetyl-2-methoxy-benzyl chloride.

## Example 6k

Methyl 2-[3-(6-fluoroquinolin-2-ylmethoxy)phenoxy)methyl]-6-methylbenzoate

MS (ESI) 432 (M+H)<sup>+</sup>. Prepared from 6-fluoroquinolin-2-ylmethyl bromide (example 27b).

5 Example 6l

Methyl 2-[3-(4-*tert*-butylbenzyloxy)phenoxy)methyl]-6-methylbenzoate

MS (ESI) 419 (M+H)<sup>+</sup>. Prepared from 4-(*tert*-butyl)benzyl bromide.

10 Example 6m

Methyl 2-[3-(4-isopropylbenzyloxy)phenoxy)methyl]-6-methylbenzoate

MS (ESI) 405 (M+H)<sup>+</sup>. Prepared from 4-isopropylbenzyl chloride.

15 Example 6n

Methyl 2-methyl-6-[3-(3-phenoxybenzyloxy)phenoxy)methyl]benzoate

MS (ESI) 455 (M+H)<sup>+</sup>. Prepared from 3-phenoxybenzyl chloride.

20 Example 6o

Methyl 2-[3-(4-*tert*-butylcyclohexylmethoxy)phenoxy)methyl]-6-methylbenzoate

MS 425 (M+H)<sup>+</sup>. Prepared from 4-*tert*-butylcyclohexylmethyl bromide (example 29a).

25 Example 6p

Methyl 2-methyl-6-[3-(quinoxalin-2-ylmethoxy)phenoxy)methyl]benzoate

MS 415 (M+H)<sup>+</sup>. Prepared from quinoxalin-2-ylmethyl chloride (See *Chem. Ber.* **1987**, 120, 649-651).

30

Example 6q

Methyl 2-methyl-6-[3-(2-methylbenzyloxy)phenoxy)methyl]benzoate

MS 377 (M+H)<sup>+</sup>. Prepared from  $\alpha$ -bromo-*o*-xylene.



## Example 6r

Methyl 2-methyl-6-[3-[2-(5-methylthiophen-2-yl)-oxazol-4-ylmethoxy]phenoxy]methyl]benzoate

MS (ESI) 450 (M+H)<sup>+</sup>. Prepared from 2-(5-methylthiophen-2-yl)oxazol-4-ylmethyl chloride (example 19a).

## Example 6s

Methyl 2-[3-(2-cyclohexyloxazol-4-ylmethoxy)phenoxy]methyl]-6-methylbenzoate

MS (ESI) 436 (M+H)<sup>+</sup>. Prepared from 2-cyclohexyloxazol-4-ylmethyl chloride (example 19b).

## Example 6t

Methyl 2-[3-[2-(3-fluorophenyl)oxazol-4-ylmethoxy]phenoxy]methyl]-6-methylbenzoate

MS (ESI) 448 (M+H)<sup>+</sup>. Prepared from 2-(3-fluorophenyl)oxazol-4-ylmethyl chloride (example 19c).

## Example 6u

Methyl 2-[3-[2-(4-fluorophenyl)oxazol-4-ylmethoxy]phenoxy]methyl]-6-methylbenzoate

MS (ESI) 448 (M+H)<sup>+</sup>. Prepared from 2-(4-fluorophenyl)oxazol-4-ylmethyl chloride (example 19d).

## Example 6v

Ethyl 2-[3-(6-chloropyridin-2-ylmethoxy)phenoxy]methyl]-6-methylbenzoate

MS (ESI) 412, 414 (M+H)<sup>+</sup>, Cl pattern. Prepared from 2-chloromethyl-6-chloropyridine (example 27c).

## Example 6w

Ethyl 2-methyl-6-[3-(5-methyl-2-phenyloxazol-4-ylmethoxy)phenoxy]methyl]benzoate

MS (ESI) 458 (M+H)<sup>+</sup>. Prepared from 4-chloromethyl-5-methyl-2-phenyloxazole.

Example 6x

5 Methyl 2-(3-benzyloxy-phenoxymethyl)-6-methyl-benzoate

MS (EI) 362 (M)<sup>+</sup>. Prepared from benzyl bromide.

Example 6y

10 Methyl 2-methyl-6-[3-(pyridin-2-ylmethoxy)-phenoxymethyl]-benzoate

Prepared from 2-chloromethyl-pyridine.

Example 6z

15 Methyl 2-[3-(7-chloroquinolin-2-ylmethoxy)phenoxymethyl]-6-methylbenzoate

MS (ESI) 447 (M+H)<sup>+</sup>, Cl pattern. Prepared from 7-chloroquinolin-2-ylmethyl bromide (example 46a).

Example 6aa

20 Methyl 2-[3-(6-methoxyquinolin-2-ylmethoxy)phenoxymethyl]-6-methylbenzoate

MS (ESI) 443 (M+H)<sup>+</sup>. Prepared from 6-methoxyquinolin-2-ylmethyl bromide (example 46b).

Example 6ab

25

Ethyl 2-[3-(2,4-diisopropyl-5-methyl-benzyloxy)-phenoxymethyl]-6-methyl-benzoate

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.14 (m, 6H), 6.62 (m, 3H), 5.10(s, 2H), 5.00 (s, 2H), 4.32 (m, 2H), 3.20 (m, 1H), 2.86 (m, 1H), 2.40 (bs, 6H), 1.28 (m, 15H). MS (EI) 484 (M)<sup>+</sup>. Prepared from 1-chloromethyl-2,4-diisopropyl-5-methyl-benzene.

30

Example 6ac

Ethyl 2-[3-(2,4-bis-trifluoromethyl-benzyloxy)-phenoxymethyl]-6-methyl-benzoate

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.92 (m, 2H), 7.82 (m, 1H), 7.30 (m, 2H), 7.18 (m, 2H), 6.57 (m, 3H), 5.29 (s, 2H), 5.10 (s, 2H), 4.31 (q, 2H), 2.39 (s, 3H), 1.29 (t, 3H). MS (EI) 512 ( $\text{M}^+$ ).

Prepared from 2,4-bis(trifluoromethyl)benzyl bromide.

## 5 Example 6ad

### Ethyl 2-[3-(biphenyl-4-ylmethoxy)-phenoxyethyl]-6-methyl-benzoate

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.60 (m, 4H), 7.40 (m, 7H), 7.18 (m, 2H), 6.58 (m, 3H), 5.09 (s, 2H), 5.07 (s, 2H), 4.30 (q, 2H), 2.39 (s, 3H), 1.28 (t, 3H). MS (EI) 452 ( $\text{M}^+$ ). Prepared from 4-phenylbenzyl chloride.

## Example 6ae

### Ethyl 2-methyl-6-[3-(naphthalen-1-ylmethoxy)-phenoxyethyl]-benzoate

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.04 (m, 1H), 7.88 (m, 2H), 7.52 (m, 4H), 7.24 (m, 4H), 6.67 (m, 2H), 6.59 (m, 1H), 5.46 (s, 2H), 5.10 (s, 2H), 4.30 (q, 2H), 2.39 (s, 3H), 1.28 (t, 3H). MS (EI) 426 ( $\text{M}^+$ ). Prepared from 1-chloromethyl-naphthalene.

## Example 6af

### Methyl 2-[3-(5-ethyl-pyridin-2-ylmethoxy)-phenoxyethyl]-6-methyl-benzoate

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.44 (bs, 1H), 7.54 (dd, 1H), 7.42 (d, 1H), 7.32 (m, 2H), 7.17 (m, 2H), 6.58 (m, 2H), 5.15 (s, 2H), 5.08 (s, 2H), 3.82 (s, 3H), 2.67 (q, 2H), 2.38 (s, 3H), 1.26 (t, 3H). MS (ESI) 392 ( $\text{M}+\text{H}^+$ ). Prepared from 5-ethyl-2-chloromethyl pyridine (example 68).

## Example 6ag

### Methyl 2-[3-(4-ethyl-benzoyloxy)-phenoxyethyl]-6-methyl-benzoate

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.28 (m, 8H), 6.58 (m, 3H), 5.08 (d, 2H), 5.00 (d, 2H), 3.81 (d, 3H), 2.68 (m, 2H), 2.38 (s, 3H), 1.24 (m, 3H). MS (EI) 390 ( $\text{M}^+$ ). Prepared from 1-chloromethyl-4-ethyl-benzene.

## Example 6ah

Methyl 2-[3-(3-bromo-benzyloxy)-phenoxy-methyl]-6-methyl-benzoate

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.58 (s, 1H), 7.44 (d, 1H), 7.25 (m, 6H), 6.57 (m, 3H), 5.08 (s, 2H), 5.00 (s, 2H), 3.81 (s, 3H), 2.38 (s, 3H). MS (EI) 440 (M)<sup>+</sup>. Prepared from 3-bromobenzyl bromide.

## Example 6ai

Ethyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenylethynyl]-benzoate

The title compound is prepared using essentially the same procedure used in Example 6 except using ethyl 2-(3-hydroxy-phenylethynyl)-6-methyl-benzoate (example 15c) in place of methyl 2-methyl-6-[(3-hydroxy-phenoxy)-methyl]-benzoate and 2-chloromethylquinoline in place of 4-chloromethyl-2-phenyl-oxazole. MS (ESI) 422 (M+H)<sup>+</sup>.

## Example 6aj

Methyl 2-methyl-6-[3-(5-phenylpyridin-2-ylmethoxy)phenoxy-methyl]benzoate

MS(ESI) 440 (M+H)<sup>+</sup>. Prepared from 5-phenylpyridin-2-ylmethyl chloride (example 27g).

## Example 6ak

Methyl 2-[3-(2-chloro-benzyloxy)-phenoxy-methyl]-6-methyl-benzoate

<sup>1</sup>H NMR (300MHz, CDCl<sub>3</sub>) δ 7.55 (m, 1H), 7.40 (m, 1H), 7.30 (m, 4H), 7.20 (m, 2H), 6.60 (m, 3H), 5.14 (s, 2H), 5.09 (s, 2H), 3.82 (s, 3H), 2.38 (s, 3H). MS (EI) 396 (M<sup>+</sup>), Cl pattern.

Prepared from 2-chlorobenzyl chloride.

## Example 6al

Methyl 2-[3-(4-chloro-benzyloxy)-phenoxy-methyl]-6-methyl-benzoate

<sup>1</sup>H NMR (300MHz, CDCl<sub>3</sub>) δ 7.32 (m, 6H), 7.18 (m, 2H), 6.56 (m, 3H), 5.08 (s, 2H), 5.00 (s, 2H), 3.81 (s, 3H), 2.38 (s, 3H). MS (EI) 396 (M<sup>+</sup>), Cl pattern. Prepared from 4-chlorobenzyl chloride.

## Example 6am

2-Methyl-6-[3-(3-methyl-quinoxalin-2-ylmethoxy)-phenoxyethyl]-benzoic acid methyl ester

MS (ESI) 429 (M+H)<sup>+</sup>. Prepared from 2-methyl-3-chloromethylquinoxaline (See Chem. Ber. **1987**, 120, 649).

## Example 6an

2-Methyl-6-[3-(naphthalen-2-ylmethoxy)-phenoxyethyl]-benzoic acid methyl ester

MS (EI) 412 (M<sup>+</sup>).

## Example 7

2-Methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxyethyl]-benzoic acid

A solution of methyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxyethyl]-benzoate (1.6 g, 3.8 mmol, example 4) in ethanol (25 mL) is heated with a 10N sodium hydroxide solution (4.0 mL, 40 mmol) at 70 °C for 14 h. The reaction is cooled, neutralized with a 2N HCl solution (20 mL) and concentrated to remove the ethanol. Ethyl acetate is added and washed with water. The aqueous layer is saturated with sodium chloride and back extracted with ethyl acetate. The organic layers are combined, dried over magnesium sulfate, filtered and concentrated to yield a crude solid. The solid is purified by column chromatography (silica, 5 to 10% methanol in dichloromethane) to provide the title compound. An analytically pure sample is prepared by recrystallization from methanol: m.p. 167-168 °C, <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) 8.15 (d, 2H), 7.80 (d, 1H), 7.71 (t, 1H), 7.61-7.51 (m, 2H), 7.26-7.10 (m, 3H), 7.00 (t, 1H), 6.66 (s, 1H), 6.52 (d, 1H), 6.46 (d, 1H), 5.26 (s, 2H), 5.15 (s, 2H), 2.44 (s, 3H); MS (ESI) 400 (M+H)<sup>+</sup>.

An alternate set of conditions that can be used for the hydrolysis of a benzoate ester is to heat a 0.1 M solution of the ester in THF/methanol (1:1) with 10 equivalents of a sodium hydroxide solution (10 N) at 60 °C for 3 h or until starting material disappears, as monitored by TLC analysis.

The following compounds are prepared using essentially the same procedure used in example 7 except using the cited ester in place of methyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxymethyl]-benzoate.

5 Example 7a

2-Methyl-6-[3-(2-quinolin-2-yl-vinyl)-phenoxymethyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.87 (bd, 1H), 8.14-8.36 (m, 4H), 8.00 (t, 1H), 7.81 (d, 1H), 7.71 (m, 1H), 7.34-7.48 (m, 5H), 7.29 (bd, 1H), 7.08 (m, 1H), 5.22 (s, 2H), 2.35 (s, 3H). MS (ESI) 396 (M+H)<sup>+</sup>. Prepared from methyl {2-methyl-6-[3-(2-quinolin-2-yl-vinyl)-phenoxymethyl]-benzoate (example 4a).

Example 7b

2-Methyl-6-{3-[2-(pyridin-2-yloxy)-ethoxy]-phenoxymethyl}-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.17 (d, 1H), 7.71 (m, 1H), 7.22 (d, 1H), 7.04-7.19 (m, 3H), 6.99 (dd, 1H), 6.86 (d, 1H), 6.55 (m, 3H), 5.13 (s, 2H), 4.53 (bs, 2H), 4.28 (bs, 2H), 2.25 (bs, 3H). MS (ESI) 380 (M+H)<sup>+</sup>. Prepared from methyl (2-methyl-6-{3-[2-(pyridin-2-yloxy)-ethoxy]-phenoxymethyl})-benzoate (example 4b).

Example 7c

2-{3-[(Benzoxazol-2-yl-methyl-amino)-methyl]-phenoxymethyl}-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.27 (bd, 1H), 7.07-7.3 (m, 5H), 7.03 (t, 1H), 6.80 (m, 3H), 5.10 (bs, 2H), 4.61 (bs, 2H), 3.03 (s, 3H), 2.38 (bs, 3H). MS (ESI) 403 (M+H)<sup>+</sup>. Prepared from methyl 2-{3-[(benzoxazol-2-yl-methyl-amino)-methyl]-phenoxymethyl}-6-methyl-benzoate (example 4c).

Example 7d

2-Methyl-6-{3-[(methyl-quinolin-2-yl-amino)-methyl]-phenoxymethyl}-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.09 (d, 1H), 7.74 (d, 1H), 7.50 (m, 3H), 7.20 (m, 4H), 6.85 (m, 3H), 6.65 (d, 1H), 5.20 (s, 2H), 4.75 (s, 2H), 3.29 (s, 3H), 2.05 (s, 3H). MS (ESI) 413 (M+H)<sup>+</sup>.

Prepared from methyl 2-methyl-6- {3-[(methyl-quinolin-2-yl-amino)-methyl]-phenoxy-methyl} -benzoate (example 4d).

#### Example 7e

5

##### 2-Methyl-6-[3-(2-phenyl-oxazol-4-ylmethoxy)-phenoxy-methyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.30 (s, 1H), 8.00 (m, 2H), 7.55 (m, 3H), 7.30 (m, 2H), 7.22 (m, 2H), 6.66 (m, 2H), 6.60 (d, 1H), 5.10 (s, 2H), 5.06 (s, 2H), 2.34 (s, 3H). MS (ESI) 416 (M+H)<sup>+</sup>.

Prepared from methyl 2-methyl-6-[3-(2-phenyl-oxazol-4-ylmethoxy)-phenoxy-methyl]-benzoate (example 6).

10

#### Example 7f

##### 2-Methyl-6-[3-(2-phenyl-thiazol-4-ylmethoxy)-phenoxy-methyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.95 (m, 2H), 7.43 (m, 3H), 7.32 (m, 2H), 7.24 (d, 1H), 7.20 (m, 1H), 7.14 (t, 1H), 6.66 (m, 1H), 6.56 (m, 1H), 5.20 (s, 2H), 5.15 (s, 2H), 2.41 (s, 3H). MS (ESI) 432 (M+H)<sup>+</sup>. Prepared from methyl 2-methyl-6-[3-(2-phenyl-thiazol-4-ylmethoxy)-phenoxy-methyl]-benzoate (example 6a).

15

20

#### Example 7g

##### 2-[3-(3,5-Dimethyl-isoxazol-4-ylmethoxy)-phenoxy-methyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.34 (m, 2H), 7.20 (m, 1H), 7.15 (t, 1H), 6.56 (m, 3H), 5.19 (s, 2H), 4.71 (s, 2H), 2.43 (s, 3H), 2.34 (s, 3H), 2.22 (s, 3H). MS (ESI) 368 (M+H)<sup>+</sup>. Prepared from methyl 2-[3-(3,5-dimethyl-isoxazol-4-ylmethoxy)-phenoxy-methyl]-6-methyl-benzoate (example 6b).

25

#### Example 7h

##### 2-Methyl-6-[3-(5-phenyl-[1,2,4]oxadiazol-3-ylmethoxy)-phenoxy-methyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.15 (m, 2H), 7.59 (m, 1H), 7.50 (m, 2H), 7.33 (m, 2H), 7.20 (m, 1H), 7.14 (t, 1H), 6.70 (m, 1H), 6.61 (m, 2H), 5.19 (s, 2H), 2.44 (s, 3H). MS (ESI) 417 (M+H)<sup>+</sup>.

30

Prepared from methyl 2-methyl-6-[3-(5-phenyl-[1,2,4]oxadiazol-3-ylmethoxy)-phenoxy-methyl]-benzoate (example 6c).

#### Example 7i

##### 2-[3-(2,5-Dimethyl-benzyloxy)-phenoxy-methyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.35 (m, 2H), 7.13-7.24 (m, 3H), 7.09 (d, 1H), 7.04 (d, 1H), 6.60 (m, 3H), 5.17 (s, 2H), 4.90 (s, 2H), 2.44 (s, 3H), 2.30 (s, 3H), 2.26 (s, 3H).

MS (ESI) 375 (M-H)<sup>-</sup>. Prepared from methyl 2-[3-(2,5-dimethyl-benzyloxy)-phenoxy-methyl]-6-methyl-benzoate (example 6d).

#### Example 7j

##### 2-[3-(2,4-Dichloro-benzyloxy)-phenoxy-methyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.40 (m, 2H), 7.34 (m, 2H), 7.24 (m, 2H), 7.17 (t, 1H), 6.59 (m, 3H), 5.19 (s, 2H), 5.03 (s, 2H), 2.45 (s, 3H). MS (ESI) 415 (M-H, Cl<sub>2</sub> pattern)<sup>-</sup>. Prepared from methyl 2-[3-(2,4-dichloro-benzyloxy)-phenoxy-methyl]-6-methyl-benzoate (example 6e).

#### Example 7k

##### 2-[3-(5-*tert*-Butyl-[1,2,4]oxadiazol-3-ylmethoxy)-phenoxy-methyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.32 (m, 2H), 7.19 (m, 1H), 7.15 (t, 1H), 6.66 (d, 1H), 6.60 (d, 1H), 6.59 (d, 1H), 5.17 (s, 2H), 5.10 (s, 2H), 1.45 (s, 9H). MS (ESI) 395 (M-H)<sup>-</sup>. Prepared from methyl 2-[3-(5-*tert*-butyl-[1,2,4]oxadiazol-3-ylmethoxy)-phenoxy-methyl]-6-methyl-benzoate (example 6f).

#### Example 7l

##### 2-{3-[3-(2,6-Dichloro-phenyl)-5-methyl-isoxazol-4-ylmethoxy]-phenoxy-methyl}-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.24-7.41 (m, 5H), 7.21 (m, 1H), 7.08 (t, 1H), 6.53 (m, 1H), 6.40 (m, 2H), 5.11 (s, 2H), 4.65 (s, 2H), 2.48 (s, 3H), 2.41 (s, 3H). MS (ESI) 496 (M-H)<sup>-</sup>. Prepared



from methyl 2-{3-[3-(2,6-dichloro-phenyl)-5-methyl-isoxazol-4-ylmethoxy]-phenoxy-methyl}-6-methyl-benzoate (example 6g).

#### Example 7m

##### 2-Methyl-6-[3-(2,4,5-trimethyl-benzyloxy)-phenoxy-methyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.35 (m, 2H), 7.20 (m, 1H), 7.15 (t, 1H), 7.10 (s, 1H), 6.97 (s, 1H), 6.60 (m, 3H), 5.16 (s, 2H), 4.87 (s, 2H), 2.42 (s, 3H), 2.25 (s, 3H), 2.21 (s, 3H), 2.20 (s, 3H). MS (ESI) 389 (M-H)<sup>-</sup>. Prepared from methyl 2-methyl-6-[3-(2,4,5-trimethyl-benzyloxy)-phenoxy-methyl]-benzoate (example 6h).

#### Example 7n

##### 2-Methyl-6-[3-(3-methyl-naphthalen-2-ylmethoxy)-phenoxy-methyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.77 (m, 3H), 7.64 (s, 1H), 7.41 (m, 2H), 7.33 (m, 2H), 7.19 (m, 2H), 6.61 (m, 3H), 5.17 (s, 2H), 5.09 (s, 2H), 2.47 (s, 3H), 2.43 (s, 3H). MS (ESI) 411 (M-H)<sup>-</sup>. Prepared from methyl 2-methyl-6-[3-(3-methyl-naphthalen-2-ylmethoxy)-phenoxy-methyl]-benzoate (example 6i).

#### Example 7o

##### 2-[3-(5-Acetyl-2-methoxy-benzyloxy)-phenoxy-methyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.05 (bs, 1H), 7.93 (bd, 1H), 7.33 (m, 2H), 7.20 (m, 1H), 7.13 (t, 1H), 6.91 (d, 1H), 6.60 (m, 3H), 5.16 (s, 2H), 5.03 (s, 2H), 3.89 (s, 3H), 2.53 (s, 3H), 2.43 (s, 3H). MS (ESI) 419 (M-H)<sup>-</sup>. Prepared from methyl 2-[3-(5-acetyl-2-methoxy-benzyloxy)-phenoxy-methyl]-6-methyl-benzoate (example 6j).

#### Example 7p

##### 2-[3-(6-Fluoroquinolin-2-ylmethoxy)phenoxy-methyl]-6-methylbenzoic acid

m.p. 153-154 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.12 (m, 2H), 7.61 (d, 1H), 7.43 (m, 2H), 7.28 (m, 2H), 7.17 (m, 1H), 7.05 (m, 1H), 6.66 (s, 1H), 6.51 (m, 2H), 5.26 (s, 2H), 5.14 (s, 2H), 2.45

(s, 3H). MS (ESI) 418 (M+H)<sup>+</sup>. Prepared from methyl 2-[3-(6-fluoroquinolin-2-ylmethoxy)phenoxyethyl]-6-methylbenzoate (example 6k).

#### Example 7q

##### 2-[3-(4-*tert*-Butylbenzyloxy)phenoxyethyl]-6-methylbenzoic acid

m.p. 122-123 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 7.41-7.28 (m, 5H), 7.23-7.12 (m, 3H), 6.61-6.55 (m, 3H), 5.16 (s, 2H), 4.95 (s, 2H), 2.45 (s, 3H), 1.32 (s, 9H). MS (ESI) 405 (M+H)<sup>+</sup>.

Prepared from methyl 2-[3-(4-*tert*-butylbenzyloxy)phenoxyethyl]-6-methylbenzoate (example 6l).

#### Example 7r

##### 2-[3-(4-Isopropylbenzyloxy)phenoxyethyl]-6-methylbenzoic acid

m.p. 132-133 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 7.35 (m, 5H), 7.22 (m, 2H), 7.17 (m, 1H), 6.58 (m, 3H), 5.15 (s, 2H), 4.97 (s, 2H), 2.92 (m, 1H), 2.46 (s, 3H), 1.25 (d, 6H). MS (ESI) 391 (M+H)<sup>+</sup>. Prepared from methyl 2-[3-(4-isopropylbenzyloxy)phenoxyethyl]-6-methylbenzoate (example 6m).

#### Example 7s

##### 2-Methyl-6-[3-(3-phenoxybenzyloxy)phenoxyethyl]benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 7.31 (m, 5H), 7.15 (m, 1H), 7.12-6.98 (m, 6H), 6.93 (m, 1H), 6.54 (m, 3H), 5.13 (s, 2H), 4.94 (s, 2H), 2.43 (s, 3H). MS (ESI) 441 (M+H)<sup>+</sup>. Prepared from methyl 2-methyl-6-[3-(3-phenoxybenzyloxy)phenoxyethyl]benzoate (example 6n).

#### Example 7t

##### 2-[3-(4-*tert*-Butylcyclohexylmethoxy)phenoxyethyl]-6-methylbenzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 7.34 (m, 2H), 7.21 (m, 1H), 7.12 (m, 1H), 6.50 (m, 3H), 5.16 (s, 2H), 3.67 (d, 2H), 2.45 (s, 3H), 1.92-1.75 (m, 4H), 1.64 (m, 2H), 0.98 (m, 4H), 0.84 (s, 9H). MS 411 (M+H)<sup>+</sup>. Prepared from methyl 2-[3-(4-*tert*-butylcyclohexylmethoxy)phenoxyethyl]-6-methylbenzoate (example 6o).

## Example 7u

2-Methyl-6-[3-(quinoxalin-2-ylmethoxy)phenoxyethyl]benzoic acid

- 5 m.p. 57-60 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.96 (s, 1H), 8.08 (m, 2H), 7.74 (m, 2H), 7.24 (m, 2H), 7.12 (m, 1H), 7.00 (m, 1H), 6.64 (s, 1H), 6.49 (m, 2H), 5.24 (s, 2H), 5.14 (s, 2H), 2.39 (s, 3H). MS 401 (M+H)<sup>+</sup>. Prepared from methyl 2-methyl-6-[3-(quinoxalin-2-ylmethoxy)phenoxyethyl]benzoate (example 6p).

## 10 Example 7v

2-Methyl-6-[3-(2-methylbenzyloxy)phenoxyethyl]benzoic acid

- 15 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 7.35 (m, 3H), 7.20 (m, 5H), 6.59 (m, 3H), 5.17 (s, 2H), 4.95 (s, 2H), 2.44 (s, 3H), 2.32 (s, 3H). MS(APCI) 385 (M+H+Na)<sup>+</sup>. Prepared from methyl 2-methyl-6-[3-(2-methylbenzyloxy)-phenoxyethyl]benzoate (example 6q).

## Example 7w

2-Methyl-6-{3-[2-(5-methylthiophen-2-yl)-oxazol-4-ylmethoxy]phenoxyethyl}benzoic acid

- 20 m.p. 129-130 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 7.54 (s, 1H), 7.48 (d, 1H), 7.27 (m, 2H), 7.11 (m, 2H), 6.74 (m, 1H), 6.66 (s, 1H), 6.53 (m, 2H), 5.12 (s, 2H), 4.95 (s, 2H), 2.51 (s, 3H), 2.39 (s, 3H). MS (ESI) 436 (M+H)<sup>+</sup>. Prepared from methyl 2-methyl-6-{3-[2-(5-methylthiophen-2-yl)-oxazol-4-ylmethoxy]phenoxyethyl}benzoate (example 6r).

## 25 Example 7x

2-[3-(2-Cyclohexyloxazol-4-ylmethoxy)phenoxyethyl]-6-methylbenzoic acid

- 30 m.p. 158-159 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 7.57 (s, 1H), 7.30 (m, 2H), 7.20 (m, 1H), 7.12 (m, 1H), 6.72 (m, 1H), 6.53 (m, 2H), 5.13 (s, 2H), 4.95 (s, 2H), 2.84 (m, 1H), 2.45 (s, 3H), 2.06 (m, 2H), 1.81 (m, 2H), 1.73-1.20 (m, 6H). Prepared from methyl 2-[3-(2-cyclohexyloxazol-4-ylmethoxy)-phenoxyethyl]-6-methylbenzoate (6s).

## Example 7y

2-{3-[2-(3-Fluorophenyl)oxazol-4-ylmethoxy]phenoxyethyl}-6-methylbenzoic acid

m.p. 152-154 °C. <sup>1</sup>H NMR (300 MHz, 5:1 CDCl<sub>3</sub>:CD<sub>3</sub>OD): δ 7.84 (d, 1H), 7.80 (s, 1H), 7.74 (d, 1H), 7.46 (m, 1H), 7.31 (m, 2H), 7.19 (m, 3H), 6.64 (m, 3H), 5.17 (s, 2H), 5.04 (s, 2H), 2.44 (s, 3H). MS (ESI) 434 (M+H)<sup>+</sup>. Prepared from methyl 2-{3-[2-(3-fluorophenyl)oxazol-4-ylmethoxy]phenoxyethyl}-6-methylbenzoate (example 6t).

## Example 7z

2-{3-[2-(4-Fluorophenyl)oxazol-4-ylmethoxy]phenoxyethyl}-6-methylbenzoic acid

m.p. 159-160 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.03 (m, 2H), 7.70 (s, 1H), 7.32 (d, 2H), 7.16 (m, 3H), 6.93 (m, 1H), 6.69 (m, 1H), 6.55 (m, 2H), 5.16 (s, 2H), 5.03 (s, 2H), 2.44 (s, 3H). MS (ESI) 434 (M+H)<sup>+</sup>. Prepared from methyl 2-{3-[2-(4-fluorophenyl)oxazol-4-ylmethoxy]phenoxyethyl}-6-methylbenzoate (example 6u).

## Example 7aa

2-[3-(6-Chloropyridin-2-ylmethoxy)phenoxyethyl]-6-methylbenzoic acid

m.p. 97-98 °C. <sup>1</sup>H NMR (300 MHz, 5:1 CDCl<sub>3</sub>:CD<sub>3</sub>OD): δ 7.73 (m, 1H), 7.47 (m, 1H), 7.28 (m, 3H), 7.16 (m, 2H), 6.60 (m, 3H), 5.16 (s, 2H), 5.12 (s, 2H), 2.42 (s, 3H). MS (ESI) 384, 386 (M+H)<sup>+</sup>, Cl pattern. Prepared from ethyl 2-[3-(6-chloropyridin-2-ylmethoxy)phenoxyethyl]-6-methylbenzoate (example 6v).

## Example 7ab

2-Methyl-6-[3-(5-methyl-2-phenyloxazol-4-ylmethoxy)phenoxyethyl]benzoic acid

m.p. 144-145 °C. <sup>1</sup>H NMR (300 MHz, 3:1 CDCl<sub>3</sub>:CD<sub>3</sub>OD): δ 7.99 (m, 2H), 7.42 (m, 3H), 7.30 (m, 2H), 7.19 (m, 2H), 6.63 (m, 3H), 5.17 (s, 2H), 4.95 (s, 2H), 2.45 (s, 3H), 2.43 (s, 3H). MS (ESI) 430 (M+H)<sup>+</sup>. Prepared from ethyl 2-methyl-6-[3-(5-methyl-2-phenyloxazol-4-ylmethoxy)-phenoxyethyl]benzoate (example 6w).

## Example 7ac

2-(3-Benzyloxy-phenoxy-methyl)-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 7.40-7.19 (m, 8H), 7.14 (t, 1H), 6.61-6.51 (m, 3H), 5.07 (s, 2H), 5.03 (s, 2H), 2.40 (s, 3H); MS (EI) 348 (M)<sup>+</sup>. Prepared from methyl 2-(3-benzyloxy-phenoxy-methyl)-6-methyl-benzoate (example 6x).

5

## Example 7ad

2-Methyl-6-[3-(pyridin-2-ylmethoxy)-phenoxy-methyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.53 (d, 1H), 7.87 (t, 1H), 7.60 (d, 1H), 7.37-7.13 (m, 5H), 6.64-6.59 (m, 3H), 5.15 (d, 4H), 2.40 (s, 3H); MS (ESI) 350 (M+H)<sup>+</sup>. Prepared from methyl 2-methyl-6-[3-(pyridin-2-ylmethoxy)-phenoxy-methyl]-benzoate (example 6y).

10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

## Example 7ae

2-[3-(7-Chloroquinolin-2-ylmethoxy)phenoxy-methyl]-6-methyl-benzoic acid

m.p. 188-193 °C; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 8.47 (d, 1H), 8.09 (s, 1H), 8.08 (d, 1H), 7.69 (dd, 2H), 7.29-7.14 (m, 4H), 6.68-6.56 (m, 3H), 5.34 (s, 2H), 5.10 (s, 2H), 2.31 (s, 3H); MS (ESI) 434, 436 (M+H;Cl)<sup>+</sup>. Prepared from methyl 2-[3-(7-chloroquinolin-2-ylmethoxy)phenoxy-methyl]-6-methylbenzoate (example 4bc).

## Example 7af

2-[3-(6-Methoxyquinolin-2-ylmethoxy)phenoxy-methyl]-6-methylbenzoic acid

m.p. 176-179 °C; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 8.29 (d, 1H), 7.91 (d, 1H), 7.60 (d, 1H), 7.42-7.39 (m, 2H), 7.28-7.14 (m, 4H), 6.67-6.55 (m, 3H), 5.27 (s, 2H), 5.09 (s, 2H), 3.90 (s, 3H), 2.31 (s, 3H); MS (ESI) 430 (M+H)<sup>+</sup>. Prepared from methyl 2-[3-(6-methoxyquinolin-2-ylmethoxy)phenoxy-methyl]-6-methylbenzoate (example 6aa).

25

## Example 7ag

2-Methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxy-methyl]-benzoic acid

m.p. 68-72 °C; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 8.25 (d, 1H), 7.88 (d, 1H), 7.78 (d, 1H), 7.67 (dd, 1H), 7.43 (dd, 1H), 7.30-7.05 (m, 7H), 6.89 (d, 1H), 5.45 (s, 2H), 5.11 (s, 2H), 2.30 (s, 3H);

30

MS (ESI) 400 (M+H)<sup>+</sup>. Prepared from isobutyl 2-methyl-6-[3-(quinolin-2-yloxy)methyl]-phenoxymethyl]-benzoate (example 4e).

#### Example 7ah

##### 2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxymethyl]-benzoic Acid

m.p. 39-65 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.44 (d, 1H), 8.26 (d, 1H), 7.85-7.75 (m, 3H), 7.59 (dd, 1H), 7.38 (s, 1H), 7.23 (obs, 2H), 7.15-7.10 (m, 2H), 6.88 (dd, 1H), 6.71 (d, 1H), 5.59 (s, 2H), 4.67 (s, 2H), 4.44 (s, 2H), 2.64 (s, 3H); MS (ESI) 414 (M+H)<sup>+</sup>. Prepared from methyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxymethyl]-benzoate (example 53).

#### Example 7ai

##### 2-[3-(Quinolin-2-ylmethoxy)-benzyloxy]-benzoic acid

m.p. 149-154 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.19 (d, 1H), 8.15 (dd, 1H), 8.11 (d, 1H), 7.83 (dd, 1H), 7.77-7.71 (m, 1H), 7.66 (d, 1H), 7.58-7.53 (m, 1H), 7.52-7.46 (m, 1H), 7.33 (t, 1H), 7.18-7.17 (m, 1H), 7.11 (t, 1H), 7.07-7.00 (m, 3H), 5.41 (s, 2H), 5.24 (s, 2H); MS (ESI) 386 (M+H)<sup>+</sup>. Prepared from methyl 2-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate (example 60).

#### Example 7aj

##### 3-Methoxy-2-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.20 (d, 1H), 8.10 (d, 1H), 7.84 (d, 1H), 7.74 (t, 1H), 7.69-7.65 (m, 2H), 7.56 (t, 1H), 7.30 (t, 1H), 7.20-7.12 (m, 3H), 7.02 (d, 1H), 5.41 (s, 2H), 5.22 (s, 2H), 3.93 (s, 3H); MS (ESI) 416 (M+H)<sup>+</sup>. Prepared from methyl 3-methoxy-2-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate (example 60a).

#### Example 7ak

##### 4-Methoxy-2-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoic acid

m.p. 117-118 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.20 (d, 1H), 8.14 (d, 1H), 8.09 (d, 1H), 7.83 (d, 1H), 7.74 (ddd, 1H), 7.65 (d, 1H), 7.56 (ddd, 1H), 7.34 (t, 1H), 7.14-7.13 (m, 1H), 7.06-7.01 (m, 2H), 6.64 (dd, 1H), 6.56 (d, 1H), 5.41 (s, 2H), 5.21 (s, 2H), 3.84 (s, 3H); MS (ESI) 416

(M+H)<sup>+</sup>. Prepared from methyl 4-methoxy-2-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate (example 60b).

#### Example 7al

##### 5-Methoxy-2-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoic acid

m.p. 248-249 °C; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 8.40 (d, 1H), 8.01 (t, 2H), 7.78 (t, 1H), 7.68 (d, 1H), 7.61 (t, 1H), 7.28-7.21 (m, 2H), 7.04 (d, 1H), 6.94 (dd, 1H), 6.78-6.71 (m, 2H), 6.56 (dd, 1H), 5.36 (s, 2H), 4.98 (s, 2H), 3.64 (s, 3H); MS (ESI) 416 (M+H)<sup>+</sup>. Prepared from methyl 5-methoxy-2-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate (example 60c).

#### Example 7am

##### 2-Methoxy-6-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoic acid

m.p. 149-152 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.34 (d, 1H), 8.16 (d, 1H), 7.80 (d, 1H), 7.75 (ddd, 1H), 7.69 (d, 1H), 7.55 (t, 1H), 7.40 (s, 1H), 7.27 (t, 1H), 7.18 (t, 1H), 6.91-87 (m, 2H), 6.60 (d, 2H), 5.45 (s, 2H), 5.08 (s, 2H), 3.89 (s, 3H); MS (ESI) 416 (M+H)<sup>+</sup>. Prepared from methyl 2-methoxy-6-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate (example 60d).

#### Example 7an

##### 2-Methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoic acid

m.p. 154-156 °C, <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.37 (d, 1H), 8.05 (d, 1H), 7.95-(d, 1H), 7.81-7.71 (m, 2H), 7.63-7.59 (t, 1H), 7.31-7.15 (m, 3H), 7.06-6.97 (m, 2H), 6.87 (d, 1H), 6.82 (d, 1H), 5.35 (s, 2H), 5.12 (s, 2H), 2.31 (s, 3H); MS (ESI) 400 (M+H)<sup>+</sup>. Prepared from ethyl-2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate (example 60e).

#### Example 7ao

##### 5-[3-(Quinolin-2-ylmethoxy)-benzyloxy]-nicotinic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.90 (s, 1H), 8.52 (s, 1H), 8.19 (d, 1H), 8.12 (d, 1H), 7.87-7.70 (m, 4H), 7.55 (t, 1H), 7.30 (t, 1H), 7.13 (s, 1H), 7.01 (t, 2H), 5.44 (s, 2H), 5.10 (s, 2H); MS (ESI)

387 (M+H)<sup>+</sup>. Prepared from methyl 5-[3-(quinolin-2-ylmethoxy)-benzyloxy]-nicotinate (example 62).

#### Example 7ap

##### 2-[3-(2,4-Diisopropyl-5-methyl-benzyloxy)-phenoxymethyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.35 (m, 2H); 7.08 (m, 4H); 6.60 (m, 3H); 5.18 (s, 2H); 4.98 (s, 2H); 3.19 (m, 1H); 2.86 (m, 1H); 2.44 (s, 3H); 2.35 (s, 3H); 1.22 (m, 12H). MS (EI) 484 (M)<sup>+</sup>.

Prepared from ethyl 2-[3-(2,4-diisopropyl-5-methyl-benzyloxy)-phenoxymethyl]-6-methyl-benzoate (example 6ab).

#### Example 7aq

##### 2-[3-(2,4-Bis-trifluoromethyl-benzyloxy)-phenoxymethyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.91 (s, 1H), 7.80 (m, 2H), 7.33 (d, 2H), 7.18 (m, 2H), 6.60 (m, 2H), 6.52 (dd, 1H), 5.24 (s, 2H), 5.17 (s, 2H), 2.45 (s, 3H). MS (EI) 484 (M)<sup>+</sup>. Prepared from ethyl 2-[3-(2,4-bis-trifluoromethyl-benzyloxy)-phenoxymethyl]-6-methyl-benzoate (example 6ac).

#### Example 7ar

##### 2-[3-(Biphenyl-4-ylmethoxy)-phenoxymethyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.58 (m, 4H), 7.44 (m, 4H), 7.35 (m, 3H), 7.18 (m, 2H), 6.60 (m, 3H), 5.17 (s, 2H), 5.02 (s, 2H), 2.44 (s, 3H). MS (EI) 424 (M)<sup>+</sup>. Prepared from ethyl 2-[3-(biphenyl-4-ylmethoxy)-phenoxymethyl]-6-methyl-benzoate (example 6ad).

#### Example 7as

##### 2-Methyl-6-[3-(naphthalen-1-ylmethoxy)-phenoxymethyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.00 (m, 1H), 7.85 (m, 2H), 7.47 (m, 4H), 7.34 (m, 2H), 7.17 (m, 2H), 6.62 (m, 3H), 5.40 (s, 2H), 5.16 (s, 2H), 2.43 (s, 3H). MS (EI) 398 (M)<sup>+</sup>. Prepared from ethyl 2-methyl-6-[3-(naphthalen-1-ylmethoxy)-phenoxymethyl]-benzoate (example 6ae).



## Example 7at

2-[3-(5-Ethyl-pyridin-2-ylmethoxy)-phenoxy-methyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.42 (bs, 1H), 7.65 (d, 1H), 7.39 (d, 1H), 6.63 (bs, 1H), 6.56 (m, 2H), 2.60 (q, 2H), 2.29 (s, 3H), 1.21 (t, 3H). MS (ESI) 378 (M+H)<sup>+</sup>. Prepared from methyl 2-[3-(5-ethyl-pyridin-2-ylmethoxy)-phenoxy-methyl]-6-methyl-benzoate (example 6af).

## Example 7au

10 2-[3-(4-Ethyl-benzyloxy)-phenoxy-methyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 7.2 (m, 8H), 6.6 (m, 3H), 5.13 (s, 2H), 5.05 (d, 2H), 2.64 (m, 2H), 2.28 (s, 3H), 1.17 (t, 3H). MS (ESI) 375 (M-H)<sup>-</sup>. Prepared from methyl 2-[3-(4-ethyl-benzyloxy)-phenoxy-methyl]-6-methyl-benzoate (example 6ag).

15 Example 7av2-[3-(3-Bromo-benzyloxy)-phenoxy-methyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 7.62 (bs, 1H), 7.50 (d, 1H), 7.42 (d, 1H), 7.33 (d, 1H), 7.08 (m, 4H), 6.67 (bs, 1H), 6.54 (m, 2H), 5.13 (s, 2H), 5.08 (s, 2H), 2.28 (s, 3H). MS (ESI) 425 (M-H)<sup>-</sup>. Prepared from methyl 2-[3-(3-bromo-benzyloxy)-phenoxy-methyl]-6-methyl-benzoate (example 6ah).

## Example 7aw

25 2-[3-[2-(5-Ethyl-pyridin-2-yl)-ethoxy]-phenoxy-methyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.36 (bs, 1H), 7.56 (d, 1H), 7.14 (m, 5H), 6.49 (m, 3H), 5.11 (bs, 2H), 4.27 (t, 2H), 3.09 (t, 2H), 2.56 (q, 2H), 2.29 (s, 3H), 1.17 (t, 3H). MS (ESI) 392 (M+H)<sup>+</sup>. Prepared from methyl 2-[3-[2-(5-ethyl-pyridin-2-yl)-ethoxy]-phenoxy-methyl]-6-methyl-benzoate (example 4f).

30

## Example 7ax

2-Methyl-6-[3-(2-quinolin-2-yl-ethoxy)-phenoxy-methyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.28 (d, 1H), 7.97 (d, 1H), 7.88 (d, 1H), 7.74 (t, 1H), 7.51 (m, 1H), 7.26 (d, 2H), 7.16 (m, 2H), 7.02 (m, 1H), 6.38 (m, 3H), 5.10 (s, 2H), 3.83 (t, 2H), 3.22 (t, 2H), 2.39 (s, 3H). Prepared from methyl 2-methyl-6-[3-(2-quinolin-2-yl-ethoxy)-phenoxy-methyl]-benzoate (example 21b).

5

#### Example 7ay

##### 2-Methyl-6-[3-(2-pyridin-2-yl-ethoxy)-phenoxy-methyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.49 (d, 1H), 7.69 (m, 1H), 7.26 (m, 3H), 7.04 (m, 2H), 6.38 (m, 4H), 5.03 (s, 2H), 3.67 (t, 2H), 2.94 (t, 2H), 2.31 (s, 3H). MS (ESI) 364 (M+H)<sup>+</sup>. Prepared from methyl 2-methyl-6-[3-(2-pyridin-2-yl-ethoxy)-phenoxy-methyl]-benzoate (example 4g).

10

15  
20

#### Example 7az

##### 2-[3-(Benzooxazol-2-ylaminomethyl)-phenoxy-methyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.61 (bt, 1H), 7.34 (d, 1H), 7.04 (m, 10H), 5.12 (s, 2H), 4.47 (bd, 2H), 2.29 (3H). MS (ESI) 389 (M+H)<sup>+</sup>. Prepared from methyl 2-[3-(benzooxazol-2-ylaminomethyl)-phenoxy-methyl]-6-methyl-benzoate (example 4h).

20

#### Example 7ba

##### 2-Methyl-6-[3-(pyridin-2-ylmethoxy-methyl)-phenoxy-methyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.52 (bd, 1H), 7.81 (m, 1H), 7.47 (d, 1H), 7.20 (m, 5H), 6.94 (m, 3H), 5.12 (s, 2H), 4.59 (s, 2H), 4.56 (s, 2H), 2.30 (s, 3H). MS (ESI) 364 (M+H)<sup>+</sup>. Prepared from methyl 2-methyl-6-[3-(pyridin-2-ylmethoxy-methyl)-phenoxy-methyl]-benzoate (example 4i).

25

#### Example 7bb

##### 2-Methyl-6-[3-(quinolin-2-ylmethoxy-methyl)-phenoxy-methyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.40 (d, 1H), 7.98 (d, 2H), 7.76 (m, 1H), 7.64 (m, 1H), 7.31 (m, 5H), 7.00 (m, 2H), 6.92 (dd, 1H), 5.12 (s, 2H), 4.79 (s, 2H), 4.62 (s, 2H), 2.32 (s, 3H). MS (ESI) 414 (M+H)<sup>+</sup>. Prepared from methyl 2-methyl-6-[3-(quinolin-2-ylmethoxy-methyl)-phenoxy-methyl]-benzoate (example 4j).

30

## Example 7bc

2-Methyl-6-{3-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxymethyl}benzoic acid

- 5  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.03 (m, 2H), 7.43 (m, 3H), 7.26 (m, 2H), 7.17 (m, 1H), 7.10 (m, 1H), 6.68 (s, 1H), 6.51 (m, 2H), 5.18 (s, 2H), 4.22 (m, 2H), 2.96 (m, 2H), 2.41 (s, 3H), 2.36 (s, 3H). MS (ESI) 444 ( $\text{M}+\text{H}$ ) $^+$ . Prepared from methyl 2-methyl-6-{3-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxymethyl}benzoate (example 21a).

## 10 Example 7bd

2-Methyl-6-[3-(6-phenylpyridin-2-ylmethoxy)phenoxymethyl]benzoic acid

- 15  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.88 (m, 2H), 7.63 (m, 1H), 7.52 (m, 1H), 7.37 (m, 4H), 7.18 (m, 2H), 6.99 (m, 2H), 6.60-6.36 (m, 3H), 5.14 (s, 2H), 5.00 (s, 2H), 2.23 (s, 3H). MS (ESI) 426 ( $\text{M}+\text{H}$ ) $^+$ . Prepared from ethyl 2-methyl-6-[3-(6-phenylpyridin-2-ylmethoxy)phenoxymethyl]-benzoate (example 33).

## Example 7be

2-Methyl-6-[3-(quinolin-2-ylmethoxy)-phenylsulfanylmethyl]-benzoic acid

- 20  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  8.40 (d, 1H), 7.99 (t, 2H), 7.78 (t, 1H), 7.65-7.57 (m, 2H), 7.20-7.01 (m, 6H), 6.85 (t, 1H), 5.33 (s, 2H), 4.25 (s, 2H), 2.26 (s, 3H); MS (ESI) 415 ( $\text{M}+\text{H}$ ) $^+$ . Prepared from isobutyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenylsulfanylmethyl]-benzoate (example 84).

25

## Example 7bf

2-Methyl-6-[3-(quinolin-2-ylmethoxy)-phenylsulfinylmethyl]-benzoic acid

- 30  $^1\text{H}$  NMR (300 MHz,  $\text{DMSO}-d_6$ )  $\delta$  8.43 (d, 1H), 8.01 (t, 2H), 7.78 (t, 1H), 7.68 (d, 1H), 7.62 (t, 1H), 7.48 (t, 1H), 7.25-7.22 (m, 4H), 7.10 (d, 1H), 6.94 (t, 1H), 5.41 (s, 2H), 4.22 (d, 1H), 4.11 (d, 1H), 2.33 (s, 3H); MS (ESI) 432 ( $\text{M}+\text{H}$ ) $^+$ . Prepared from isobutyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenylsulfinylmethyl]-benzoate (example 85).

## Example 7bg

2-Methyl-6-[3-(quinolin-2-ylmethoxy)-phenylsulfonylmethyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 8.43 (d, 1H), 8.05-7.98 (m, 2H), 7.77 (t, 1H), 7.67 (d, 1H), 7.60 (t, 1H), 7.51 (t, 1H), 7.43-7.35 (m, 2H), 7.26-7.20 (m, 3H), 6.95-6.92 (m, 1H), 5.42 (s, 2H), 4.81 (s, 2H), 2.32 (s, 3H); MS (ESI) 448 (M+H)<sup>+</sup>. Prepared from isobutyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenylsulfonylmethyl]-benzoate (example 86).

## Example 7bh

2-Methyl-6-[3-(quinolin-2-ylmethoxy)-phenylethynyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.19 (d, 1H), 8.04 (d, 1H), 7.89 (d, 1H), 7.82 (d, 1H), 7.73 (dt, 2H), 7.64 (t, 2H), 7.54 (t, 1H), 7.26 (t, 1H), 6.94 (dd, 1H), 6.81-6.85 (m, 2H), 5.34 (s, 2H), 2.76 (s, 3H); MS (ESI) 394 (M+H)<sup>+</sup>. Prepared from ethyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenylethynyl]-benzoate (example 6ai).

## Example 7bi

2-Methyl-6-[3-(5-phenylpyridin-2-ylmethoxy)phenoxyethyl]benzoic acid mpt 80-83°C. <sup>1</sup>H

NMR (300 MHz, 5:1 CDCl<sub>3</sub>:CD<sub>3</sub>OD): δ 8.76 (s, 1H), 7.97 (d, 1H), 7.62 (m, 3H), 7.48 (m, 3H), 7.30 (m, 2H), 7.19 (m, 2H), 6.68 (s, 1H), 6.62 (d, 2H), 5.21 (s, 2H), 5.17 (s, 2H), 2.44 (s, 3H). MS (ESI) 426 (M+H)<sup>+</sup>. Prepared from methyl 2-methyl-6-[3-(5-phenylpyridin-2-ylmethoxy)phenoxyethyl]benzoate (example 6aj).

## Example 7bj

2-[3-(2-Chloro-benzoyloxy)-phenoxyethyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300MHz, DMSO) δ 7.56 (m, 1H), 7.50 (m, 1H), 7.38 (m, 2H), 7.14 (m, 2H), 7.06 (m, 2H), 6.67 (s, 1H), 6.58 (m, 2H), 5.13 (bd, 4H), 2.28 (s, 3H). MS (EI) 382 (M<sup>+</sup>), Cl pattern. Prepared from 2-[3-(2-chloro-benzoyloxy)-phenoxyethyl]-6-methyl-benzoate (example 6ak).

## Example 7bk

2-[3-(4-Chloro-benzyloxy)-phenoxymethyl]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300MHz, DMSO) δ 7.42 (m, 3H), 7.08 (m, 5H), 6.66 (s, 1H), 6.54 (m, 2H), 5.13 (s, 2H), 5.07 (s, 2H), 2.28 (s, 3H). MS (EI) 382 (M<sup>+</sup>), Cl pattern. Prepared from 2-[3-(4-chloro-benzyloxy)-phenoxymethyl]-6-methyl-benzoate (example 6al).

5

## Example 7bl

2-Methyl-6-[3-(3-methyl-quinoxalin-2-ylmethoxy)-phenoxymethyl]-benzoic acid

<sup>1</sup>H NMR (300MHz, DMSO) δ 8.05 (m, 2H), 7.82 (m, 2H), 7.14 (m, 4H), 6.75 (s, 1H), 6.61 (m, 2H), 5.41 (s, 2H), 5.11 (s, 2H), 2.76 (s, 3H), 2.27 (s, 3H). MS (ESI) 415 (M+H)<sup>+</sup>. Prepared from 2-Methyl-6-[3-(3-methyl-quinoxalin-2-ylmethoxy)-phenoxymethyl]-benzoic acid methyl ester (example 6am).

10

## Example 7bm

2-Methyl-6-[3-(naphthalen-2-ylmethoxy)-phenoxymethyl]-benzoic acid

<sup>1</sup>H NMR (300MHz, DMSO) δ 7.94 (m, 4H), 7.54 (m, 3H), 7.16 (m, 4H), 6.69 (s, 1H), 6.58 (m, 2H), 5.24 (s, 2H), 5.11 (s, 2H), 2.29 (s, 3H). Prepared from 2-Methyl-6-[3-(naphthalen-2-ylmethoxy)-phenoxymethyl]-benzoic acid methyl ester (example 6an).

15

20

## Example 8

3-[(2-Methoxyethoxy)-methoxy]-benzonitrile

To a cooled (0 °C) suspension of sodium hydride (840mg, 60% dispersion in mineral oil, 21 mmol) in THF (20 mL) is added a solution comprising 3-hydroxy-benzonitrile (2.4g, 20 mmol), MEM Chloride (2.25 mL, 20 mmol) and DMPU (2 mL) in THF (20 mL). On complete addition, the cold bath is removed and stirring continued for 3h. The reaction mixture is then diluted with ether, washed with water and brine, dried over MgSO<sub>4</sub> and concentrated. The residue is purified by flash chromatography (silica, 30% ethyl acetate 10 % dichloromethane in hexane) to give the title compound as an oil. MS (ESI) 207 (M+H)<sup>+</sup>.

25

30

## Example 8a

3-[(2-Methoxyethoxy)-methoxy]-benzaldehyde

The title compound is prepared using essentially the same procedure used in example 8 except using 3-hydroxy-benzaldehyde in place of 3-hydroxy-benzonitrile. MS (EI) 210 (M)<sup>+</sup>.

## 5 Example 9

3-[(2-Methoxyethoxy)-methoxy]-benzylamine

To a cooled (0°C) solution of 3-[(2-methoxyethoxy)-methoxy]-benzonitrile (3.9g, 18.8 mmol, example 8) in THF (40 mL) is added LAH (40 mL, 1M in THF). The resulting solution is stirred for 10 min. then the cold bath removed and stirring continued for 2 h. The resulting mixture is cooled to 0 °C then water (1.5 mL) added dropwise followed by NaOH solution (1.5 mL, 5M) and water (1.5 mL). The resulting suspension is diluted with ether then filtered through celite. The filtrate is concentrated to give the title compound which is used without further purification. MS (ESI) 211 (M+H)<sup>+</sup>.

## 15 Example 10

3-(Quinolin-2-ylaminomethyl)-phenol

To a solution of 3-[(2-methoxyethoxy)-methoxy]-benzylamine (422 mg, 2mmol, example 9) in DMSO (4 mL) is added 2-chloroquinoline (328 mg, 2 mmol). The resulting solution is warmed to 140 °C and stirred at this temperature for 3h. The resulting mixture is cooled, diluted with water, then extracted with ethyl acetate. The organic extract is washed with brine, dried over MgSO<sub>4</sub> and concentrated. The residue is taken up in methanol (10 mL) then p-toluene sulphonic acid monohydrate (190 mg, 1 mmol) is added. This mixture is warmed to 60 °C and stirred at this temperature for 2h. The reaction mixture is then cooled, concentrated under reduced pressure and the residue purified by flash chromatography (silica, 30% ethyl acetate in dichloromethane) to give the title compound. MS (ESI) 251 (M+H)<sup>+</sup>.

The following compounds are prepared using essentially the same procedure used in example 10 except using the cited chloride and amine in place of 2-chloro-quinoline and 3-[(2-methoxyethoxy)-methoxy]-benzylamine.

## 20 Example 10a

3-[(N-Benzoxazol-2-yl-N-methyl-amino)-methyl]-phenol

MS (ESI) 255 (M+H)<sup>+</sup>. Prepared from 3-[(methylamino)-methyl]-(2-methoxyethoxy-methoxy)-benzene (example 28) and 2-chloro-benzoxazole. Also, omit heating to 140 °C. Reaction stirred at room temperature.

## Example 10b

3-[(N-Methyl-N-quinolin-2-yl-amino)-methyl]-phenol

MS (ESI) 265 (M+H)<sup>+</sup>. Prepared from 3-[(methylamino)-methyl]-(2-methoxyethoxy-methoxy)-benzene (example 28) and 2-chloro-quinoline.

## Example 10c

3-(Benzoxazol-2-ylaminomethyl)-phenol

The title compound is prepared using essentially the same procedure used in example 10 except 2-chloro-benzoxazole is used in place of 2-chloro-quinoline. Also, omit heating to 140 °C. Reaction stirred at room temperature. MS (ESI) 241 (M+H)<sup>+</sup>.

## Example 11

2-(3-([2-Methoxyethoxy]-methoxy)-phenoxy)-ethanol

To a cooled (0 °C) solution of t-butyl (3-([2-methoxyethoxy]-methoxy)-phenoxy)-acetate (1.2g, 3.8 mmol, example 12) in THF (10 mL) is added a solution of lithium aluminum hydride (5 mL, 1M in THF). The resulting solution is stirred for 10 min. then water (0.2 mL) is added dropwise, followed by NaOH solution (0.2 mL, 5M) and water (0.2 mL). The resulting mixture is diluted with ether, filtered through celite and the filtrate concentrated to give the title compound as an oil which is used without further purification. MS (EI) 242 (M)<sup>+</sup>.

## Example 11a

2-(5-Methyl-2-phenyloxazol-4-yl)ethanol

The title compound is prepared using essentially the same procedure used in Example 11 except using methyl 2-(5-methyl-2-phenyloxazol-4-yl)acetate (example 32) in place of *t*-butyl 3-([2-methoxyethoxy]-methoxy)-phenoxy]-acetate. MS (ESI) 204 (M+H)<sup>+</sup>.

## 5 Example 12

### *t*-Butyl 3-([2-methoxyethoxy]-methoxy)-phenoxy]-acetate

The title compound is prepared using essentially the same procedure used in Example 4 except using 3-([2-methoxyethoxy]-methoxy)-phenol (example 13) in place of 3-(quinolin-2-ylmethoxy)-phenol and *t*-butyl bromoacetate in place of methyl 2-bromomethyl-6-methylbenzoate. MS (EI) 312 (M)<sup>+</sup>.

## Example 13

### 3-([2-Methoxyethoxy)-methoxy]-phenol

To a cooled (0 °C) suspension of NaH (440 mg, 60% dispersion in oil, 11 mmol) in THF (10 mL) is slowly added a solution comprising 3-benzoyl-phenol (2.14g, 10 mmol), MEM chloride (1.28 mL, 10.5 mmol) and DMPU (3 mL) in THF (20 mL). On complete addition, the cold bath is removed and stirring continued for 2.5 h. Sat. NH<sub>4</sub>Cl solution is added and the mixture diluted with ether, washed with water and brine, dried over MgSO<sub>4</sub> and concentrated. The residue is taken up in methanol (10 mL) and THF (10 mL) then sodium hydroxide solution (10 mL, 2N) added. This mixture is stirred for 20 min. then hydrochloric acid (10 mL, 2N) added. The mixture is then diluted with ether, washed with sat. NaHCO<sub>3</sub> solution and brine, dried over MgSO<sub>4</sub> and concentrated. The residue is purified by flash chromatography (silica, 30% ethyl acetate in hexanes) to give the title compound as an oil. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.00 (t, 1H), 6.52 (bd, 1H), 6.48 (bs, 1H), 6.38 (bd, 1H), 5.14 (s, 2H), 3.71 (m, 2H), 3.47 (m, 2H), 3.30 (s, 3H).

## Example 14

### [2-Methoxyethoxy)-methoxy]-3-[2-(pyridin-2-yloxy)-ethoxy]-benzene

To a solution of 2-(3-([2-methoxyethoxy]-methoxy)-phenoxy]-ethanol (242 mg, 1 mmol, example 11) in DMSO (1.5 mL) is added sodium hydride (44mg, 60% dispersion in mineral oil,



1.1 mmol) followed by 2-fluoro-pyridine (176 mL, 2 mmol). The resulting solution is warmed to 60 °C and stirred at this temperature for 3 h. cooled, diluted with ether, washed with water and brine, dried over MgSO<sub>4</sub> and concentrated. The residue is purified by flash chromatography (silica, 30% ethyl acetate in hexanes) to give the title compound. MS (ESI) 320 (M+H)<sup>+</sup>.

#### Example 15

##### 3-(2-Quinolin-2-yl-vinyl)-phenol

To a solution of 1-(2-methoxyethoxy)-methoxy-3-(2-quinolin-2-yl-vinyl)-benzene (120mg, 0.35 mmol, example 16) is added p-toluene sulphonic acid monohydrate. (74 mg, 0.39 mmol). The resulting solution is heated to 60 °C and stirred at this temperature for 4 h. The reaction mixture is then cooled, concentrated and the residue taken up in dichloromethane. This solution is washed with sat. NaHCO<sub>3</sub> solution, dried over MgSO<sub>4</sub> then concentrated to give the title compound as a solid. MS (ESI) 248 (M+H)<sup>+</sup>.

The following compounds are prepared using essentially the same procedure used in example 15 except using the cited MEM ether in place of 1-(2-methoxyethoxy)-methoxy-3-(2-quinolin-2-yl-vinyl)-benzene.

#### Example 15a

##### 3-[2-(pyridin-2-yloxy)-ethoxy]-phenol

MS (ESI) 232 (M+H)<sup>+</sup>. Prepared from [(2-methoxyethoxy)-methoxy]-3-[2-(pyridin-2-yloxy)-ethoxy]-benzene (example 14).

#### Example 15b

##### 3-(Quinolin-2-yloxymethyl)-phenol

MS (ESI) 252 (M+H)<sup>+</sup>. Prepared from 2-[3-(2-methoxy-ethoxymethoxy)-benzyloxy]-quinoline (example 81).

#### Example 15c

##### Ethyl 2-(3-hydroxy-phenylethynyl)-6-methyl-benzoate

MS (EI) 280 (M)<sup>+</sup>. Prepared from ethyl-2-[3-(2-methoxy-ethoxymethoxy)-phenylethynyl]-6-methyl-benzoate (example 98).

#### Example 16

##### 1-(2-Methoxyethoxy)-methoxy-3-(2-quinolin-2-yl-vinyl)-benzene

To a cooled (-78 °C) suspension of triphenyl-(quinolin-2-yl-methyl)-phosphonium chloride (1.76g, 4 mmol, example 17) in THF (24 mL) is added, dropwise, n-butyl lithium solution (1.7 mL, 2.5 M in hexanes). The resulting mixture is stirred for 30 min. then a solution of 3-[(2-methoxyethoxy)-methoxy]-benzaldehyde (756 mg, 3.6 mmol, example 8a) in THF (3 mL) is added. This mixture is stirred for 30 min then the cold bath removed and stirring continued for 2h. The reaction mixture is then diluted with ethyl acetate, washed with sat. ammonium acetate solution and brine, dried over MgSO<sub>4</sub> and concentrated. The residue is purified by flash chromatography (silica, 40% ethyl acetate in hexanes) to give the title compound as an oil. MS (ESI) 336 (M+H)<sup>+</sup>.

#### Example 17

##### Triphenyl-(quinolin-2-yl-methyl)-phosphonium chloride

To a solution of 2-chloromethyl-quinoline (2.9g, 20 mmol) in acetonitrile (32 mL) is added triphenylphosphine (4.49g, 17 mmol). The resulting mixture is warmed to 60 °C and stirred at this temperature for 15 h. This mixture is cooled, diluted with ether, then filtered. The solid is washed with ether, then dried under high vacuum to give the title compound as a solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.20 (d, 1H), 8.06 (d, 1H), 7.95 (m, 6H), 7.42-7.8 (m, 13H), 6.10 (d, 2H).

#### Example 18

##### {2-Methyl-6-[3-(2-quinolin-2-yl-ethyl)-phenoxy-methyl]-phenoxy}-acetic acid

To a solution of {2-methyl-6-[3-(2-quinolin-2-yl-vinyl)-phenoxy-methyl]-phenoxy}-acetic acid (94 mg, 0.23 mmol, example 41a) in DMF (1.5 mL) is added tris(triphenylphosphine)rhodium chloride (25mg, 0.027 mmol). The resulting solution is placed under an atmosphere of hydrogen, heated to 60 °C and stirred at this temperature for 5h. The reaction mixture is cooled to room temperature and the system is then flushed with nitrogen and concentrated under vacuum. The

residue is purified by reverse phase HPLC to give the title compound as a trifluoroacetate salt.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.57 (d, 1H), 8.40 (d, 1H), 8.0 (m, 2H), 7.80 (t, 1H), 7.52 (d, 1H), 7.24 (bd, 1H), 7.14 (m, 2H), 7.04 (t, 1H), 6.93 (bs, 1H), 6.83 (d, 1H), 6.74 (d, 1H), 5.11 (s, 2H), 4.50 (s, 2H), 3.68 (t, 2H), 3.20 (t, 2H), 2.28 (s, 3H). MS (ESI) 428 (M+H)<sup>+</sup>.

5

The following compound is prepared using essentially the same procedure used in example 18 except using the cited acid in place of {2-methyl-6-[3-(2-quinolin-2-yl-vinyl)-phenoxymethyl]-phenoxy}-acetic acid.

#### 10 Example 18a

##### 2-Methyl-6-[3-(2-quinolin-2-yl-ethyl)-phenoxymethyl]-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.62 (d, 1H), 8.44 (d, 1H), 8.0 (m, 2H), 7.80 (t, 1H), 7.61 (d, 1H), 7.21 (d, 1H), 7.11 (d, 1H), 7.07 (t, 1H), 6.94 (bs, 1H), 6.76 (bd, 1H), 6.68 (d, 1H), 5.09 (s, 2H), 3.70 (t, 2H), 3.18 (t, 2H), 2.40 (s, 3H). MS (ESI) 398 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(2-quinolin-2-yl-vinyl)-phenoxymethyl]-benzoic acid (example 4a).

15

#### Example 19

##### 4-Chloromethyl-2-phenyl-oxazole

Benzamide (1.21g, 10 mmol) is mixed with 1,3-dichloroacetone (1.26g, 10 mmol) and the mixture heated to 130 °C and stirred at this temperature for 1 h. The resulting mixture is then cooled, diluted with ethyl acetate, washed with K<sub>2</sub>CO<sub>3</sub> solution (sat.), then brine, dried over MgSO<sub>4</sub> and concentrated to give the title compound as a solid, which is used without further purification. MS (ESI) 194 (M+H, Cl pattern)<sup>+</sup>.

25

The following compounds are prepared using essentially the same procedure used in Example 19 except using the cited amide in place of benzamide.

#### 30 Example 19a

##### 2-(5-Methylthiophen-2-yl)oxazol-4-ylmethyl chloride

MS (ESI) 214, 216 (M+H)<sup>+</sup>, Cl pattern. Prepared from 5-methylthiophene-2-carboxamide.

## Example 19b

2-Cyclohexyloxazol-4-ylmethyl chloride

5 MS (ESI) 200, 202 (M+H)<sup>+</sup>, Cl pattern. Prepared from cyclohexanecarboxamide.

## Example 19c

2-(3-Fluorophenyl)oxazol-4-ylmethyl chloride

10 MS (ESI) 212, 214 (M+H)<sup>+</sup>, Cl pattern. Prepared from 3-fluorobenzamide.

## Example 19d

2-(4-Fluorophenyl)oxazol-4-ylmethyl chloride

15 MS (ESI) 212, 214 (M+H)<sup>+</sup>, Cl pattern. Prepared from 4-fluorobenzamide

## Example 20

4-Chloromethyl-2-phenyl-thiazole

20 A solution of thiobenzamide (1.37g, 10 mmol) and 1,3-dichloro-acetone (1.27g, 10 mmol) in ethanol (25 mL) is warmed to 75 °C and stirred at this temperature for 1h. The resulting solution is cooled, poured into ice then brought to pH 8 with K<sub>2</sub>CO<sub>3</sub> solution (sat.). This mixture is extracted with ethyl acetate, dried over MgSO<sub>4</sub> and concentrated to give the title compound. This product is used without further purification. MS (ESI) 210 (M+H)<sup>+</sup>.

25

## Example 21

{2-Methyl-6-[3-(2-pyridin-2-yl-ethoxy)-phenoxy-methyl]-phenoxy}-acetonitrile

30 To a solution of [2-methyl-6-(3-hydroxy-phenoxy-methyl)-phenoxy]-acetonitrile (135 mg, 0.5 mmol, example 25) and 2-(pyridin-2-yl)-ethanol (126 mL, 0.94 mmol) in THF (2 mL) is added triphenylphosphine (262 mg, 1 mmol) followed by DEAD (118 mL, 0.75 mmol). The resulting solution is stirred for 2h, then concentrated and the residue purified by flash chromatography (silica, 50% ethyl acetate in hexanes) to give the title compound as an oil. MS (ESI) 375 (M+H)<sup>+</sup>.

The following compound is prepared using essentially the same procedure used in Example 21 except using the cited alcohol and phenol in place of 2-(pyridin-2-yl)-ethanol and [2-methyl-6-(3-hydroxy-phenoxy-methyl)-phenoxy]-acetonitrile respectively.

#### Example 21a

##### Methyl 2-methyl-6-{3-[2-(5-methyl-2-phenyloxazol-4-yl)ethoxy]phenoxy-methyl}benzoate

MS (ESI) 458 (M+H)<sup>+</sup>. Prepared from 2-(5-methyl-2-phenyloxazol-4-yl)ethanol (example 11a) and methyl 2-(3-hydroxyphenoxy-methyl)-6-methylbenzoate (example 5).

#### Example 21b

##### Methyl 2-methyl-6-[3-(2-quinolin-2-yl-ethoxy)-phenoxy-methyl]-benzoate

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.08 (m, 2H), 7.80 (d, 1H), 7.70 (t, 1H), 7.51 (t, 1H), 7.42 (d, 1H), 7.29 (m, 2H), 7.16 (m, 2H), 6.53 (m, 3H), 5.06 (s, 2H), 4.46 (t, 2H), 3.81 (s, 3H), 3.45 (t, 2H), 2.37 (s, 3H). MS (ESI) 428 (M+H)<sup>+</sup>. Prepared from 2-quinolin-2-yl-ethanol (example 69a) and 2-(3-hydroxy-phenoxy-methyl)-6-methyl-benzoate (example 5).

#### Example 22

##### 2-Cyanomethoxy-3-methylbenzaldehyde

A mixture of 2-hydroxy-3-methylbenzaldehyde (10.2 g, 75.0 mmoles, Aldrich), bromoacetonitrile (5.70 mL, 82.5 mmoles), and potassium carbonate (11.4 g, 82.5 mmoles) in DMF (150 mL) is heated to 55°C for 3 hours, cooled, then diluted with ether. The mixture is washed with distilled water, saturated NaCl solution, then the organic layer dried over MgSO<sub>4</sub> and concentrated to give the title compound as a yellow solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 10.20 (s, 1H), 7.70 (d, 1H), 7.53 (d, 1H), 7.29 (m, 1H), 4.81 (s, 2H), 2.42 (s, 3H).

The following compounds are prepared using essentially the same procedure used in example 22 except using the cited phenol in place of 2-hydroxy-3-methylbenzaldehyde.

#### Example 22a

2-Cyanomethoxy-3,5-dichloro-benzaldehyde

MS (EI) 229, 231 (M)<sup>+</sup>. Prepared from 3,5-dichlorosalicylaldehyde.

## 5 Example 22b

2-Cyanomethoxy-5-chloro-3-methoxy-benzaldehyde

MS (EI) 225 (M)<sup>+</sup>. Prepared from 5-chloro-2-hydroxy-3-methoxy-benzaldehyde (example 44).

## 10 Example 22c

Methyl 2-(2-formyl-6-methyl-phenoxy)-propionate

The title compound is prepared using essentially the same procedure used in example 22 except using methyl 2-bromopropionate in place of bromoacetonitrile.

## Example 23

(2-Hydroxymethyl-6-methyl-phenoxy)-acetonitrile

A 2M triglyme solution of sodium borohydride (16.0 mL, 32.1 mmoles) is slowly added to a cooled (-78°C) solution of 2-cyanomethoxy-3-methylbenzaldehyde (11.25g, 64.2 mmoles, example 22) in THF (180 mL). After stirring for one hour, the reaction is warmed to 0°C for two hours, then quenched with 2N HCl (16.8 mL) and diluted with ether. The organic layer is isolated and washed with twice with distilled water and brine, then dried over MgSO<sub>4</sub>. The organic solution is concentrated to give the title compound as a yellow oil.

## 25

The following compounds are prepared using essentially the same procedure used in example 23 except using the cited aldehyde in place of 2-cyanomethoxy-3-methylbenzaldehyde.

## Example 23a

## 30

(2,4-Dichloro-6-hydroxymethyl-phenoxy)-acetonitrile

Prepared using 2-cyanomethoxy-3,5-dichloro-benzaldehyde (example 22a)

## Example 23b

(4-Chloro-2-hydroxymethyl-6-methoxy-phenoxy)-acetonitrile

MS (EI) 227 (M)<sup>+</sup>. Prepared using 2-cyanomethoxy-5-chloro-3-methoxy-benzaldehyde (example 22b)

5

Example 23c

2-(2-Hydroxymethyl-6-methyl-phenoxy)-propionic acid methyl ester

MS (EI) 194 (M)<sup>+</sup>. Prepared using methyl 2-(2-formyl-6-methyl-phenoxy)-propionate (example 22c).

10

Example 24

(2-Bromomethyl-6-methyl-phenoxy)-acetonitrile

Triphenylphosphine (15.2g, 57.8 mmoles) is added to 2-cyanomethoxy-3-methylbenzylalcohol (9.3g, 52.5 mmoles, example 23) in THF (175 mL). The mixture is stirred until homogeneous and cooled to 0°C, followed by addition, in three portions, of *N*-bromosuccinimide (10.3g, 57.8 mmoles). After 90 minutes the reaction is concentrated and the residue purified by column chromatography (silica, 5:1 hex: EtOAc) to yield the title compound as a pale yellow crystalline solid. MS (EI) 239, 241 (M)<sup>+</sup>, Br pattern.

The following compounds are prepared using essentially the same procedure used in example 24 except using the cited alcohol in place of 2-cyanomethoxy-3-methylbenzylalcohol.

25 Example 24a

(2-Bromomethyl-4,6-dichloro-phenoxy)-acetonitrile

MS (EI) 277 (M-16)<sup>+</sup>. Prepared from (2,4-dichloro-6-hydroxymethyl-phenoxy)-acetonitrile (example 23a).

30

Example 24b

(2-Bromomethyl-4-chloro-6-methoxy-phenoxy)-acetonitrile

MS (EI) 289 (M)<sup>+</sup>. Prepared from (4-chloro-2-hydroxymethyl-6-methoxy-phenoxy)-acetonitrile (example 23b).

Example 24c

5

Methyl 2-(2-bromomethyl-6-methyl-phenoxy)-propionate

MS (EI) 286 (M)<sup>+</sup>, bromine pattern. Prepared from methyl 2-(2-hydroxymethyl-6-methyl-phenoxy)-propionate (example 23c).

10 Example 25

20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

(2-[3-Hydroxyphenoxy)methyl]-6-methylphenoxy)acetonitrile

Heated (60°C) a mixture of 2-cyanomethoxy-3-methylbenzyl bromide (10.2g, 42.7 mmoles, example 24), resorcinol (18.8g, 171 mmoles), and potassium carbonate (47.2g, 342 mmoles) in acetonitrile (140 mL) for two hours. The reaction is diluted with ether and washed three times with distilled water, once with brine, and dried over MgSO<sub>4</sub>. The organic layer is isolated and concentrated, and the resulting residue is purified by column chromatography (silica, 5% EtOAc/CH<sub>2</sub>Cl<sub>2</sub>) to yield the title compound as a white crystalline solid. MS(EI) 269 (M)<sup>+</sup>.

The following compounds are prepared using essentially the same procedure used in example 25 except using the cited bromide in place of 2-cyanomethoxy-3-methylbenzyl bromide.

Example 25a

25 [4-Chloro-2-(3-hydroxy-phenoxy)methyl]-6-methyl-phenoxy]-acetonitrile

Prepared from (2-bromomethyl-4-chloro-6-methyl-phenoxy)-acetonitrile (example 43).

Example 25b

30 [4,6-Dichloro-2-(3-hydroxy-phenoxy)methyl]-phenoxy]-acetonitrile

Prepared from (2-bromomethyl-4,6-dichloro-phenoxy)-acetonitrile (example 24a).

Example 26



### 2-Methyl-6,7-difluoroquinoline

To a refluxing solution of 3,4-difluoroaniline (2.30 ml, 23.2 mmoles), tetrachloro-1,4-benzoquinone (5.70 g, 23.2 mmoles), and concentrated hydrochloric acid (6 ml) in 2-butanol (40 ml) is added crotonaldehyde (1.92 ml, 23.2 mmoles). After 2.5 hours the reaction mixture is concentrated and the resulting residue is stirred in warm (50 °C) THF (15 ml). This mixture is cooled (0 °C) and the solid collected by filtration and washed with cold THF. The solid is stirred in distilled water (200 ml), and the resulting solution made basic with  $K_2CO_3$  and extracted with EtOAc (3 x 100 ml). The organic extracts are combined and dried over sodium sulfate, and then concentrated to give the title compound. MS (ESI) 180 (M+H)<sup>+</sup>.

The following compounds are prepared using essentially the same procedure used in example 26 except using the cited aniline in place of 3,4-difluoroaniline.

#### Example 26a

### 2-Methyl-6,8-difluoroquinoline

MS (ESI) 180 (M+H)<sup>+</sup>. Prepared using 2,4-difluoro-aniline.

#### Example 27

### 6,8-Difluoroquinolin-2-ylmethyl bromide

A solution of 2-methyl-6,8-difluoroquinoline (0.147 g, 0.820 mmoles, Example 26a), benzoyl peroxide (9.93 mg, 0.0410 mmoles), and *N*-bromosuccinamide (0.168 g, 0.943 mmoles) in carbon tetrachloride (20 mL) is heated to reflux for 18 hours. The reaction is concentrated and the resulting residue purified by column chromatography (silica, 3:1  $CH_2Cl_2$ :hexane, then  $CH_2Cl_2$ ) to yield the title compound as a white solid. MS (ESI) 258, 260 (M+H)<sup>+</sup>, Br pattern.

The following compounds are prepared using essentially the same procedure used in example 27 except using the cited methyl compound in place of 6,8-difluoro-2-methyl quinoline.

#### Example 27a

### 6,7-Difluoroquinolin-2-ylmethyl chloride

MS (ESI) 214, 216 (M+H)<sup>+</sup>, Cl pattern. Prepared from 6,7-difluoro-quinoline (example 26) and NCS in place of NBS.

Example 27b

5

6-Fluoroquinolin-2-ylmethyl bromide

MS (ESI) 240, 242 (M+H)<sup>+</sup>, Br pattern. Prepared from 6-fluoro-2-methylquinoline.

Example 27c

10

2-Chloromethyl-6-chloropyridine

MS (ESI) 162, 164, 166 (M+H)<sup>+</sup>, Cl<sub>2</sub> pattern. Prepared from 6-chloro-2-picoline and NCS in place of NBS.

Example 27d

15

2-Bromomethyl-benzonitrile

MS (EI) 195 (M)<sup>+</sup>, Br pattern. Prepared from *o*-tolunitrile.

Example 27e

20

Methyl 3-bromomethyl-thiophene-2-carboxylate

MS (EI) 234 (M)<sup>+</sup>, Br pattern. Prepared from methyl 3-methyl-thiophene-2-carboxylate.

Example 27f

25

6,7-Dichloro-2-chloromethyl-quinoline

MS (ESI) 246 (M+H)<sup>+</sup>. Prepared from 6,7-dichloro-quinoline and NCS in place of NBS.

Example 27g

30

5-Phenylpyridin-2-ylmethyl chloride

MS (ESI) 204, 206 (M+H)<sup>+</sup>, Cl pattern. Prepared from 5-phenyl-2-methylpyridine (example 104) and NCS in place of NBS.

## Example 28

3-[(Methylamino)-methyl]-(2-methoxyethoxy-methoxy)-benzene

- 5 To a solution of 3-(2-methoxyethoxy-methoxy)-benzaldehyde (2.10g, 10 mmol, example 8a) in THF (60 mL) is added methylamine (20 mL, 2M in THF) followed by palladium on carbon (210 mg, 10% Pd ). The resulting mixture is stirred for 24 h under an atmosphere of hydrogen gas, then purged with nitrogen, filtered through celite and the filtrate concentrated. The residue is purified by flash chromatography (silica, 10% methanol in dichloromethane) to give the title compound as an oil.
- 10

## Example 29

1-Methyl-4-oxo-1,4-dihydroquinolin-2-ylmethyl bromide

- 15 A solution of 1-methyl-4-oxo-1,4-dihydroquinolin-2-ylmethanol (112 mg, 0.592 mmoles, Coppola, G.M. *J. Heterocyclic Chem.*, **1986**, 23, 1717) and phosphorous tribromide (56.2 uL, 0.592 mmoles) in 3:1 CH<sub>2</sub>Cl<sub>2</sub>:DMF (20 mL) is stirred 18 hours and another portion (20 uL) of phosphorous tribromide is added. After 24 hours distilled water (10 mL) is added and extracted with EtOAc. The organic layer is concentrated and the resulting residue purified by column chromatography (silica, 20:1 CH<sub>2</sub>Cl<sub>2</sub>:MeOH) to yield the title compound as a white solid. MS (ESI) 252, 254 (M+H)<sup>+</sup>, Br pattern.
- 20

The following compound is prepared using essentially the same procedure used in example 29 except using the cited alcohol in place of 1-methyl-4-oxo-1,4-dihydroquinolin-2-ylmethanol.

## Example 29a

4-tert-Butylcyclohexylmethyl bromide

- 30 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 3.27 (d, 2H), 1.93 (m, 2H), 1.81 (m, 2H), 1.54 (m, 2H), 0.98 (m, 4H), 0.84 (s, 9H). Prepared from 4-tert-butylcyclohexylmethanol (example 30).

## Example 30

4-tert-Butylcyclohexylmethanol

To a cooled (0°C) solution of 4-*tert*-butylcyclohexane-carboxylic acid (3.00g, 16.3 mmol) in THF (30 mL) is slowly added a THF solution of borane-THF complex (1.0M, 21.2 mL, 21.2 mmol). The solution is stirred at room temperature for 18 hours, then quenched with 2N HCl solution (30 mL), and extracted with EtOAc. The organic layer is isolated, washed with 1N NaOH, dried over sodium sulfate, and concentrated to yield the title compound as a clear oil. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 2.06 (d, 2H), 1.82 (m, 4H), 1.52 (m, 2H), 0.88 (m, 4H), 0.83 (s, 9H).

### Example 31

#### Methyl 4-bromo-3-oxopentanoate

To a cooled (0°C) solution of methyl 3-oxopentanoate (9.62 mL, 76.8 mmol, Acros) in carbon tetrachloride (60 mL) is added dropwise over a period of 45 minutes a solution of bromine (3.96 mL, 76.8 mmol) in carbon tetrachloride (10 mL). After 30 minutes, let stir at room temperature for one hour. Bubbled N<sub>2</sub> through reaction mixture for twenty minutes. Concentrated to yield the title compound as a brown oil. MS (EI) 208, 210 (M)<sup>+</sup>, Br pattern.

### Example 32

#### Methyl 2-(5-methyl-2-phenyloxazol-4-yl)acetate

A solution of benzamide (0.606g, 5.00 mmol) and methyl 4-bromo-3-oxopentanoate (1.05g, 5.00 mmol, example 31) are heated in toluene (6 mL) to 120°C for 18 hours. The reaction is then purified by column chromatography (silica, 4:1 hex: EtOAc) to give the title compound as a clear oil. MS (APCI) 232 (M+H)<sup>+</sup>.

### Example 33

#### Ethyl 2-methyl-6-[3-(6-phenylpyridin-2-ylmethoxy)phenoxyethyl]benzoate

A solution of phenylboronic acid (74.0mg, 0.607 mmol), ethyl 2-[3-(6-chloropyridin-2-ylmethoxy)-phenoxyethyl]-6-methylbenzoate (250mg, 0.607 mmol, example 6v), and sodium carbonate (77.8mg, 1.21 mmol) in 1:1 H<sub>2</sub>O:AcCN (8 mL) is stirred under vacuum for five minutes. The reaction is placed under nitrogen, and tetrakis(triphenylphosphine)-palladium(0) (60.7mg) is added followed by heating to 90°C. After two hours, another portion

(15mg) of phenylboronic acid is added. After another hour heating is stopped. Distilled water (10 mL) is added, followed by extraction with methylene chloride (twice with 20 mL). The organic extracts are combined and concentrated, and the resulting residue is purified by column chromatography (silica, 6:1 hex:EtOAc) to yield the title compound. MS (ESI) 454 (M+H)<sup>+</sup>.

5

#### Example 34

##### 1-Oxyquinolin-2-ylmethyl chloride

10



Partitioned 2-(chloromethyl)quinoline hydrochloride (1.00 g, 4.67 mmoles) between methylene chloride (15 mL) and sodium hydroxide solution (1M, 15 mL) to form the free base. The organic layer is isolated and cooled (0°C), followed by addition of 3-chloroperbenzoic acid (57-86%, 1.13 g, ~4.67 mmoles). After stirring at room temperature 18 hours the reaction mixture is washed with dilute sodium hydroxide. The organic layer is isolated and concentrated. The resulting residue is purified by column chromatography (silica, 1:1 hex:EtOAc) to yield the title compound as a white solid. MS (ESI) 194, 196 (M+H)<sup>+</sup>, Cl pattern.

20



#### Example 35

##### {2-[3-(Quinolin-2-ylmethoxy)phenoxy]methyl-6-methylphenoxy} acetonitrile

25

3-(Quinolin-2-ylmethoxy)-phenol (1.3g, 5.4 mmol, example 3), (2-bromomethyl-6-methylphenoxy)-acetonitrile (1.56 g, 6.5 mmol, example 24), tetrabutylammonium iodide (99 mg, 0.27 mmol) and potassium carbonate (0.45 g, 3.3 mmol) are refluxed in acetone (20 mL) for 16 h. The reaction is filtered, washed with dichloromethane, concentrated and purified by column chromatography (silica, 1% ether in dichloromethane) to provide the title compound. MS (ESI) 411 (M+H)<sup>+</sup>.

30

#### Example 35a

##### {2-Methyl-6-[3-(quinolin-2-ylaminomethyl)-phenoxy]methyl]-phenoxy}-acetonitrile

MS (ESI) 409 (M+H)<sup>+</sup>. Prepared from 3-(quinolin-2-ylaminomethyl)-phenol (example 10).

## Example 35b

{2-Methyl-6-[3-(2-quinolin-2-yl-vinyl)-phenoxymethyl]-phenoxy}-acetonitrile

- 5 MS (ESI) 407 (M+H)<sup>+</sup>. Prepared from 3-(2-quinolin-2-yl-vinyl)-phenol (example 15).

## Example 35c

(2-Methyl-6-[3-[2-(pyridin-2-yloxy)-ethoxy]-phenoxymethyl]-phenoxy)-acetonitrile

- 10 MS (ESI) 391 (M+H)<sup>+</sup>. Prepared from 3-[2-(pyridin-2-yloxy)-ethoxy]-phenol (example 15a).

## Example 35d

{2-[3-(Benzooxazol-2-ylaminomethyl)-phenoxymethyl]-6-methyl-phenoxy}-acetonitrile

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.15 (m, 1H), 5.36 (bs, 1H), 5.08 (s, 2H), 4.67 (s, 4H), 2.37 (s, 3H). MS (ESI) 400 (M+H)<sup>+</sup>. Prepared from 3-(benzooxazol-2-ylaminomethyl)-phenol (example 10c).

## Example 35e

{2-[3-(4-Chloro-quinolin-2-ylmethoxymethyl)-phenoxymethyl]-6-methyl-phenoxy}-acetonitrile

- <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.22 (d, 1H), 8.06 (d, 1H), 7.76 (m, 2H), 7.63 (m, 1H), 7.32 (m, 2H), 7.23 (m, 1H), 7.13 (m, 1H), 7.04 (m, 2H), 6.94 (m, 1H), 5.09 (s, 2H), 4.83 (s, 2H), 4.72 (s, 2H), 4.68 (s, 2H), 2.39 (s, 3H). MS (ESI) 459 (M+H)<sup>+</sup>. Prepared from 3-(4-chloro-quinolin-2-ylmethoxymethyl)-phenol (example 74b).
- 25

## Example 35f

{2-[3-(6-Methoxy-quinolin-2-ylmethoxymethyl)-phenoxymethyl]-6-methyl-phenoxy}-acetonitrile

- <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.08 (d, 1H), 7.96 (d, 1H), 7.60 (d, 1H), 7.19 (m, 8H), 6.92 (dd, 1H), 5.07 (s, 2H), 4.84 (s, 2H), 4.71 (s, 2H), 4.66 (s, 2H), 3.93 (s, 3H), 2.39 (s, 3H). MS (ESI) 455 (M+H)<sup>+</sup>. Prepared from 3-(6-methoxy-quinolin-2-ylmethoxymethyl)-phenol (example 74c).
- 30

## Example 35g

2-Methyl-6-[3-(quinolin-2-ylmethoxymethyl)-phenoxy]-acetonitrile

Prepared from 3-(quinolin-2-ylmethoxymethyl)-phenol (example 74a).

5

The following compounds are prepared using essentially the same procedure used in example 35 except using the cited bromide in place of (2-bromomethyl-6-methyl-phenoxy)-acetonitrile.

## Example 36a

10

2-{2-Methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxy]-propionic acid methyl ester

MS (ESI) 457 (M+H)<sup>+</sup>. Prepared using methyl 2-(2-bromomethyl-6-methyl-phenoxy)-propionate (example 24c).

15

## Example 36b

{2,4-Dichloro-6-[3-(quinolin-2-ylmethoxy)-phenoxy]-acetonitrile

MS (ESI) 465 (M+H)<sup>+</sup>, Cl<sub>2</sub> pattern. Prepared from (2-bromomethyl-4,6-dichloro-phenoxy)-acetonitrile (example 24a)

20

## Example 36c

{4-Chloro-2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxy]-acetonitrile

25

MS (ESI) 445 (M+H)<sup>+</sup>. Prepared from (2-bromomethyl-4-chloro-6-methyl-phenoxy)-acetonitrile (example 43).

## Example 36d

30

{2-tert-Butyl-6-[3-(quinolin-2-ylmethoxy)-phenoxy]-acetonitrile

MS (ESI) 453 (M+H)<sup>+</sup>. Prepared from (2-bromomethyl-6-tert-butyl-phenoxy)-acetonitrile (example 43a).

## Example 36e

{4-Chloro-2-methoxy-6-[3-(quinolin-2-ylmethoxy)-phenoxymethyl]-phenoxy}-acetonitrile

MS (ESI) 461 (M+H)<sup>+</sup>, Cl pattern. Prepared from (2-bromomethyl-4-chloro-6-methoxy-phenoxy)-acetonitrile (example 24b).

5

## Example 36f

2-[3-(Quinolin-2-ylmethoxy)-phenoxymethyl]-benzonitrile

MS (ESI) 386 (M+H)<sup>+</sup>. Prepared from 2-bromomethyl-benzonitrile (example 27d).

10

## Example 36g

Methyl 2-[3-(Quinolin-2-ylmethoxy)-phenoxymethyl]-thiophene-2-carboxylate

MS (ESI) 406 (M+H)<sup>+</sup>. Prepared from methyl 3-bromomethyl-thiophene-2-carboxylate (example 27e).

15

## Example 36h

Ethyl {2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxymethyl]-phenoxy}-acetate

MS (ESI) 457 (M+H)<sup>+</sup>. Prepared from ethyl (2-bromomethyl-6-methyl-phenoxy)-acetate (example 43b).

20

## Example 36i

Ethyl 7-[3-(quinolin-2-ylmethoxy)-phenoxymethyl]-benzofuran-2-carboxylate

MS (ESI) 354 (M+H)<sup>+</sup>. Prepared from ethyl 7-bromomethyl-benzofuran-2-carboxylate (example 94).

25

## Example 36j

30

Ethyl {2-methyl-6-[3-methyl-5-(quinolin-2-ylmethoxy)-phenoxymethyl]-phenoxy}-acetate

The title compound is prepared using essentially the same procedure used in example 35 except using 3-methyl-5-(quinolin-2-ylmethoxy)-phenol (example 55) in place of



3-(quinolin-2-ylmethoxy)-phenol and ethyl (2-bromomethyl-6-methyl-phenoxy)-acetate (example 43b) in place of (2-bromomethyl-6-methyl-phenoxy)-acetonitrile.

The following compounds are prepared using essentially the same procedure used in example 35 except using the cited phenol in place of 3-(quinolin-2-ylmethoxy)-phenol and (2-bromomethyl-4-chloro-6-methyl-phenoxy)-acetonitrile (example 43) in place of (2-bromomethyl-6-methyl-phenoxy)-acetonitrile.

#### Example 36k

##### {4-Chloro-2-methyl-6-[3-(2-pyridin-2-yl-ethoxy)-phenoxy-methyl]-phenoxy}-acetonitrile

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.56 (d, 1H), 7.62 (m, 1H), 7.30 (m, 1H), 7.18 (m, 3H), 6.57 (m, 4H), 4.99 (s, 2H), 4.67 (s, 2H), 4.36 (t, 2H), 3.27 (t, 2H), 2.36 (s, 3H). MS (ESI) 409 (M+H)<sup>+</sup>.

Prepared from 3-(2-pyridin-2-yl-ethoxy)-phenol (example 71a).

#### Example 36l

##### {2-[3-(Benzoxazol-2-ylaminomethyl)-phenoxy-methyl]-4-chloro-6-methyl-phenoxy}-acetonitrile

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.14 (m, 10H), 5.35 (bs, 1H), 5.04 (s, 2H), 4.67 (d, 2H), 4.63 (s, 2H), 2.34 (s, 3H). MS (ESI) 434 (M+H)<sup>+</sup>. Prepared from 3-(benzoxazol-2-ylaminomethyl)-phenol (example 10c).

#### Example 37

##### 2-[3-(2-Chloromethyl-benzyloxy)-phenoxy-methyl]-quinoline

The title compound is prepared using essentially the same procedure used in example 35 except using excess 1,2-bis-chloromethyl-benzene in place of (2-bromomethyl-6-methyl-phenoxy)-acetonitrile and without using tetrabutylammonium iodide. MS (ESI) 390 (M+H)<sup>+</sup>, CI pattern.

#### Example 38

##### {2-[3-(Quinolin-2-ylmethoxy)-phenoxy-methyl]-phenyl}-acetonitrile

Sodium cyanide (14 mg, 0.28 mmol) is added to a solution of 2-[3-(2-chloromethyl-benzyloxy)-phenoxymethyl]-quinoline (110 mg, 0.28 mmol, example 37) in DMSO (5 mL) and the reaction is stirred 5 h. The reaction is partitioned between water and ethyl acetate, the organic phase is washed with water, dried and concentrated to provide the title compound which is used without further purification. MS (ESI) 381 (M+H)<sup>+</sup>.

### Example 39

#### {2-[3-(Quinoxalin-2-ylmethoxy)phenoxymethyl]-6-methylphenoxy} acetonitrile

A solution of (2-[3-hydroxyphenoxymethyl]-6-methylphenoxy)acetonitrile (100 mg, 0.37 mmol, example 25), quinoxalin-2-ylmethyl chloride [72 mg, 0.40 mmol (See *Chem. Ber.* **1987**, 120, 649-651)] in DMF (1 mL) is heated with potassium carbonate (105 mg, 0.75 mmol) at 60 °C for 16 h. The reaction is filtered and partitioned between ethyl acetate and water. The organic phase is washed with water, dried over magnesium sulfate, concentrated and purified by column chromatography (silica, 30% ethyl acetate in hexanes) to provide the title compound. MS (ESI) 412 (M+H)<sup>+</sup>.

The following compounds are prepared using essentially the same procedure used in example 39 except using the cited halide in place of quinoxalin-2-ylmethyl chloride.

### Example 39a

#### {2-[3-(7-Chloro-isoquinolin-3-ylmethoxy)-phenoxymethyl]-6-methyl-phenoxy}-acetonitrile

MS (ESI) 445 (M+H, Cl pattern)<sup>+</sup>. Prepared from {2-methyl-6-[(3-hydroxy-phenoxy)-methyl]-phenoxy}-acetonitrile (example 25) and (7-chloro-isoquinolin-3-yl) methyl bromide (see Ewing, William R.; Becker, Michael R.; Choi-Sledeski, Yong Mi; Pauls, Heinz W.; He, Wei; Condon, Stephen M.; Davis, Roderick S.; Hanney, Barbara A.; Spada, Alfred P.; Burns, Christopher J.; Jiang, John Z.; Li, Aiwen; Myers, Michael R.; Lau, Wan F.; Poli, Gregory B; *Substituted piperazinone derivatives and other oxoazaheterocyclyl compounds useful as factor Xa inhibitors*. PCT Int. Appl., (1999), WO 9937304.

### Example 39b

{2-Methyl-6-[3-(naphthalen-2-ylmethoxy)-phenoxy-methyl]-phenoxy}-acetonitrile

MS (ESI) 410 (M+H)<sup>+</sup>. Prepared from naphthalen-2-ylmethyl chloride.

Example 39c

5

{2-[3-(4-tert-Butyl-benzyloxy)-phenoxy-methyl]-6-methyl-phenoxy}-acetonitrile

MS (ESI) 416 (M+H)<sup>+</sup>. Prepared from 4-tert-butylbenzyl chloride.

Example 39d

10

{2-Methyl-6-[3-(2-phenoxy-ethoxy)-phenoxy-methyl]-phenoxy}-acetonitrile

MS (ESI) 390 (M+H)<sup>+</sup>. Prepared from 2-phenoxy-ethyl-bromide.

Example 39e

15

{2-Methyl-6-[3-(3-phenyl-propoxy)-phenoxy-methyl]-phenoxy}-acetonitrile

MS (ESI) 388 (M+H)<sup>+</sup>. Prepared from 3-phenyl-propyl bromide.

Example 39f

20

{2-Methyl-6-[3-(3-phenoxy-benzyloxy)-phenoxy-methyl]-phenoxy}-acetonitrile

MS (ESI) 452 (M+H)<sup>+</sup>. Prepared from 3-phenoxy-benzyl chloride.

Example 39g

25

{2-[3-(3-Methoxy-benzyloxy)-phenoxy-methyl]-6-methyl-phenoxy}-acetonitrile

MS (ESI) 390 (M+H)<sup>+</sup>. Prepared from 3-methoxy-benzyl chloride.

Example 39h

30

{2-[3-(3,4-Dichloro-benzyloxy)-phenoxy-methyl]-6-methyl-phenoxy}-acetonitrile

MS (ESI) 428 (M+H)<sup>+</sup>. Prepared from 3,4-dichloro-benzyl chloride.

Example 39i

{2-[3-(6,7-Difluoroquinolin-2-ylmethoxy)phenoxy)methyl]-6-methyl-phenoxy}acetonitrile  
MS (ESI) 446 (M+H)<sup>+</sup>. Prepared from 6,7-difluoroquinolin-2-ylmethyl chloride (example 27).

5 Example 39j

{2-[3-(6,8-Difluoroquinolin-2-ylmethoxy)phenoxy)methyl]-6-methylphenoxy}acetonitrile  
MS (ESI) 446 (M+H)<sup>+</sup>. Prepared from 6,8-difluoroquinolin-2-ylmethyl bromide (example 27a).

10 Example 39k

{2-Methyl-6-[3-(1-oxyquinolin-2-ylmethoxy)phenoxy)methyl]phenoxy}acetonitrile  
MS (ESI) 427 (M+H)<sup>+</sup>. Prepared from 1-oxyquinolin-2-ylmethyl chloride (example 34).

Example 39l

{2-[3-(6-Fluoroquinolin-2-ylmethoxy)phenoxy)methyl]-6-methylphenoxy}acetonitrile  
MS (ESI) 429 (M+H)<sup>+</sup>. Prepared from 6-fluoroquinolin-2-ylmethyl bromide (example 27b).

Example 39m

{2-Methyl-6-[3-(1-methyl-4-oxo-1,4-dihydroquinolin-2-ylmethoxy)phenoxy)methyl]-phenoxy}acetonitrile  
MS (ESI) 441 (M+H)<sup>+</sup>. Prepared from 1-methyl-4-oxo-1,4-dihydroquinolin-2-ylmethyl bromide (example 29).

Example 39n

{2-[3-(4-Chloroquinolin-2-ylmethoxy)phenoxy)methyl]-6-methylphenoxy}acetonitrile  
MS (ESI) 445 (M+H)<sup>+</sup>. Prepared from 4-chloroquinolin-2-ylmethyl chloride (example 46).

Example 39o

{2-[3-(7-Chloroquinolin-2-ylmethoxy)phenoxy)methyl]-6-methylphenoxy}acetonitrile

MS (ESI) 445 (M+H)<sup>+</sup>. Prepared from 7-chloroquinolin-2-ylmethyl chloride (example 46a).

Example 39p

5 {2-[3-(6-Methoxyquinolin-2-ylmethoxy)phenoxyethyl]-6-methylphenoxy}-acetonitrile

MS (ESI) 441 (M+H)<sup>+</sup>. Prepared from 6-methoxyquinolin-2-ylmethyl chloride (example 46b).

Example 39q

10 {2-Methyl-6-[3-(pyridin-4-ylmethoxy)-phenoxyethyl]-phenoxy}-acetonitrile

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.62 (bd, 2H), 7.24 (m, 6H), 6.62 (m, 3H), 5.08 (s, 2H), 5.06 (s, 2H), 4.70 (s, 2H), 2.39 (s, 3H). MS (ESI) 361 (M+H)<sup>+</sup>. Prepared from 4-chloromethyl-pyridine hydrochloride.

15 {2-Methyl-6-[3-(pyridin-2-ylmethoxy)-phenoxyethyl]-phenoxy}-acetonitrile

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.60 (d, 1H), 7.72 (m, 1H), 7.52 (d, 1H), 7.21 (m, 5H), 6.63 (m, 3H), 5.20 (s, 2H), 5.04 (s, 2H), 4.69 (s, 2H), 2.39 (s, 3H). MS (ESI) 361 (M+H)<sup>+</sup>. Prepared from 2-chloromethyl-pyridine hydrochloride.

20

Example 39s

{2-Methyl-6-[3-(pyridin-3-ylmethoxy)-phenoxyethyl]-phenoxy}-acetonitrile

25 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.68 (bs, 1H), 8.59 (bd, 1H), 7.78 (m, 1H), 7.24 (m, 5H), 6.63 (m, 3H), 5.07 (s, 2H), 5.06 (s, 2H), 4.70 (s, 2H), 2.39 (s, 3H). MS (ESI) 361 (M+H)<sup>+</sup>. Prepared from 3-chloromethyl-pyridine hydrochloride.

Example 39t

30

{2-[3-(6,7-Dichloro-quinolin-2-ylmethoxy)-phenoxyethyl]-6-methyl-phenoxy}-acetonitrile

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.20 (s, 1H), 8.10 (d, 1H), 7.94 (s, 1H), 7.70 (d, 1H), 7.20 (m, 4H), 6.65 (m, 3H), 5.34 (s, 2H), 5.05 (s, 2H), 4.69 (s, 2H), 2.38 (s, 3H). MS (ESI) 479 (M+H)<sup>+</sup>.

Prepared from 6,7-dichloro-2-chloromethyl-quinoline (example 27f).

#### Example 39u

5 {2-Methyl-6-[3-(2-phenyl-thiazol-4-ylmethoxy)-phenoxymethyl]-phenoxy}-acetonitrile  
MS (ESI) 443 (M+H)<sup>+</sup>. Prepared from 4-chloromethyl-2-phenyl-thiazole (example 20).

The following compounds are prepared using essentially the same procedure used in example 39 except using the cited phenol in place of (2-[3-hydroxyphenoxymethyl]-6-methylphenoxy)acetonitrile and 6-fluoroquinolin-2-ylmethyl bromide (example 27b) in place of quinoxalin-2-ylmethyl chloride.

15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995

#### Example 40a

5 {4-Chloro-2-[3-(6-fluoroquinolin-2-ylmethoxy)phenoxymethyl]-6-methylphenoxy} acetonitrile  
MS (ESI) 463, 465 (M+H)<sup>+</sup>, Cl<sub>2</sub> pattern. Prepared from (2-[3-hydroxyphenoxymethyl]-4-chloro-6-methylphenoxy)acetonitrile (example 25a).

#### Example 40b

{2,4-Dichloro-6-[3-(6-fluoroquinolin-2-ylmethoxy)phenoxymethyl]phenoxy} acetonitrile  
MS (ESI) 483, 485, 487 (M+H)<sup>+</sup>, Cl<sub>2</sub> pattern. Prepared from [4,6-dichloro-2-(3-hydroxy-phenoxymethyl)-phenoxy]-acetonitrile (example 25b)

#### 25 Example 41

{2-Methyl-6-[3-(quinolin-2-ylaminomethyl)-phenoxymethyl]-phenoxy}-acetic acid

To a solution of {2-methyl-6-[3-(quinolin-2-ylaminomethyl)-phenoxymethyl]-phenoxy}-acetonitrile (134 mg, 0.31 mmol, example 35a) in methanol (1 mL) is added THF (1 mL) followed by sodium hydroxide solution (0.2 mL, 10 N). The resulting mixture is warmed to 60 °C and stirred at this temperature for 3h. The reaction mixture is then cooled to room temperature and acidified to ca. pH 5 with hydrochloric acid (1 mL, 2N), then extracted with ethyl acetate, washed with brine, dried over MgSO<sub>4</sub> and concentrated. The residue is purified by flash chromatography (silica, 10% methanol in dichloromethane) to give the title compound. <sup>1</sup>H NMR

(300 MHz, CDCl<sub>3</sub>) δ 8.10 (d, 1H), 7.80 (d, 1H), 7.68 (t, 2H), 7.40 (t, 1H), 7.25 (m, 2H), 7.04 (m, 3H), 6.90 (m, 3H), 6.6 (bs, 1H), 5.15 (s, 2H), 4.60 (d, 2H), 4.50 (s, 2H), 2.27 (s, 3H). MS (ESI) 429 (M+H)<sup>+</sup>.

- 5 The following compounds are prepared using essentially the same procedure used in example 41 except using the cited nitrile or ester in place of {2-methyl-6-[3-(quinolin-2-ylaminomethyl)-phenoxy]methyl}-phenoxy}-acetonitrile.

#### Example 41a

##### {2-Methyl-6-[3-(2-quinolin-2-yl-vinyl)-phenoxy]methyl}-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.38 (d, 1H), 8.01 (d, 1H), 7.97 (d, 1H), 7.89 (d, 1H), 7.83 (d, 1H), 7.78 (dt, 1H), 7.59 (dt, 1H), 7.53 (d, 1H), 7.44 (bd, 1H), 7.36 (m, 3H), 7.25 (bd, 1H), 7.11 (t, 1H), 7.02 (dt, 1H), 5.25 (s, 2H), 4.54 (s, 2H), 2.32 (s, 3H). MS (ESI) 426 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(2-quinolin-2-yl-vinyl)-phenoxy]methyl}-phenoxy}-acetonitrile (example 35b).

#### Example 41b

##### (2-Methyl-6-{3-[2-(pyridin-2-yloxy)-ethoxy]-phenoxy]methyl}-phenoxy)-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.17 (dd, 1H), 7.72 (m, 1H), 7.29 (dd, 1H), 7.20 (m, 2H), 7.07 (t, 1H), 6.99 (m, 1H), 6.85 (d, 1H), 6.60 (m, 3H), 5.13 (s, 2H), 4.57 (t, 2H), 4.49 (s, 2H), 4.30 (t, 2H), 2.28 (s, 3H). MS (ESI) 410 (M+H)<sup>+</sup>. Prepared from (2-methyl-6-{3-[2-(pyridin-2-yloxy)-ethoxy]-phenoxy]methyl}-phenoxy)-acetonitrile (example 35c).

#### Example 41c

##### {2-[3-(7-Chloro-isoquinolin-3-ylmethoxy)-phenoxy]methyl}-6-methyl-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 9.13 (d, 1H), 7.96 (s, 1H), 7.79 (s, 1H), 7.74 (d, 1H), 7.65 (d, 1H), 7.30 (d, 1H), 7.20 (d, 1H), 7.10 (m, 2H), 6.81 (s, 1H), 6.62 (d, 1H), 6.49 (d, 1H), 5.25 (s, 2H), 5.16 (s, 2H), 4.59 (s, 2H), 2.37 (s, 3H). MS (ESI) 464 (M+H, Cl pattern)<sup>+</sup>. Prepared from {2-[3-(7-chloro-isoquinolin-3-ylmethoxy)-phenoxy]methyl}-6-methyl-phenoxy}-acetonitrile (example 39a).

## Example 41d

{2-Methyl-6-[3-(naphthalen-2-ylmethoxy)-phenoxy]methyl]-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.00 (m, 3H), 7.48 (dd, 1H), 7.43 (m, 2H), 7.23 (m, 2H), 7.16 (m, 2H), 7.05 (t, 1H), 6.62 (m, 1H), 6.55 (m, 2H). MS (ESI) 429 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(naphthalen-2-ylmethoxy)-phenoxy]methyl]-phenoxy}-acetonitrile (example 39b).

## Example 41e

{2-[3-(4-tert-Butyl-benzyloxy)-phenoxy]methyl]-6-methyl-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.38 (m, 3H), 7.27 (m, 2H), 7.24 (d, 1H), 7.17 (d, 1H), 7.10 (d, 1H), 6.59 (m, 3H), 5.07 (s, 2H), 4.98 (s, 2H), 4.57 (s, 2H), 2.33 (s, 3H), 1.33 (s, 9H). MS (ESI) 435 (M+H)<sup>+</sup>. Prepared from {2-[3-(4-tert-butyl-benzyloxy)-phenoxy]methyl]-6-methyl-phenoxy}-acetonitrile (example 39c).

## Example 41f

{2-Methyl-6-[3-(2-phenoxy-ethoxy)-phenoxy]methyl]-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.03-7.33 (m, 6H), 6.95 (m, 3H), 6.57 (m, 3H), 5.07 (s, 2H), 4.55 (s, 2H), 4.29 (m, 4H), 2.33 (s, 3H). MS (ESI) 409 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(2-phenoxy-ethoxy)-phenoxy]methyl]-phenoxy}-acetonitrile (example 39d).

## Example 41g

{2-Methyl-6-[3-(3-phenyl-propoxy)-phenoxy]methyl]-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.07-7.41 (m, 9H), 6.54 (m, 3H), 5.08 (s, 2H), 4.58 (s, 2H), 3.94 (t, 2H), 2.80 (t, 2H), 2.34 (s, 3H), 2.09 (qn, 2H). MS (ESI) 407 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(3-phenyl-propoxy)-phenoxy]methyl]-phenoxy}-acetonitrile (example 39e).

## Example 41h

{2-Methyl-6-[3-(3-phenoxy-benzyloxy)-phenoxy]methyl]-phenoxy}-acetic acid



<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.31 (m, 4H), 7.06-7.25 (m, 6H), 7.01 (m, 2H), 6.94 (m, 1H), 6.57 (m, 3H), 5.06 (s, 2H), 4.99 (s, 2H), 4.56 (s, 2H), 2.33 (s, 3H). MS (ESI) 471 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(3-phenoxy-benzyloxy)-phenoxymethyl]-phenoxy}-acetonitrile (example 39f).

5

#### Example 41i

##### {2-[3-(3-Methoxy-benzyloxy)-phenoxymethyl]-6-methyl-phenoxy}-acetic acid

MS (ESI) 409 (M+H)<sup>+</sup>. Prepared from {2-[3-(3-methoxy-benzyloxy)-phenoxymethyl]-6-methyl-phenoxy}-acetonitrile (example 39g).

10

#### Example 41j

##### {2-[3-(3,4-Dichloro-benzyloxy)-phenoxymethyl]-6-methyl-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.60 (d, 1H), 7.51 (d, 1H), 7.30-7.38 (m, 2H), 7.28 (d, 1H), 7.23 (d, 1H), 7.17 (d, 1H), 6.66 (m, 3H), 5.16 (s, 2H), 5.05 (s, 2H), 4.65 (s, 2H), 2.42 (s, 3H). MS (ESI) 447 (M+H)<sup>+</sup>. Prepared from {2-[3-(3,4-dichloro-benzyloxy)-phenoxymethyl]-6-methyl-phenoxy}-acetonitrile (example 39h).

15

20

25

30

35

40

45

50

55

60

65

70

75

80

85

90

95

100

105

110

115

120

125

130

135

140

145

150

155

160

165

170

175

180

185

190

#### Example 41k

##### {2-[3-(6,7-Difluoroquinolin-2-ylmethoxy)phenoxymethyl]-6-methylphenoxy}acetic acid

m.p. 94-95 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.10 (d, 1H), 7.73 (m, 1H), 7.56 (d, 1H), 7.44 (d, 1H), 7.29 (s, 1H), 7.18 (m, 1H), 7.08 (m, 2H), 6.79 (s, 1H), 6.61 (d, 1H), 6.51 (m, 1H), 5.31 (s, 2H), 5.15 (s, 2H), 4.61 (s, 2H), 2.35 (s, 3H). MS (ESI) 466 (M+H)<sup>+</sup>. Prepared from {2-[3-(6,7-difluoroquinolin-2-ylmethoxy)phenoxymethyl]-6-methyl-phenoxy}-acetonitrile (example 39i).

25

#### Example 41l

##### {2-[3-(6,8-Difluoroquinolin-2-ylmethoxy)phenoxymethyl]-6-methylphenoxy}acetic acid

m.p. 137-141 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.16 (d, 1H), 7.79 (d, 1H), 7.29-7.05 (m, 6H), 6.71 (s, 1H), 6.61 (m, 2H), 5.53 (s, 2H), 5.10 (s, 2H), 4.57 (s, 2H), 2.33 (s, 3H). MS (ESI) 466

30

(M+H)<sup>+</sup>. Prepared from {2-[3-(6,8-difluoroquinolin-2-ylmethoxy)phenoxy]methyl}-6-methylphenoxy}acetonitrile (example 39j).

#### Example 41m

##### {2-Methyl-6-[3-(1-oxyquinolin-2-ylmethoxy)phenoxy]methyl}phenoxy}acetic acid

m.p. 146-147 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.73 (d, 1H), 8.01 (m, 1H), 7.93 (m, 1H), 7.89 (m, 1H), 7.70 (m, 1H), 7.55 (d, 1H), 7.26 (m, 2H), 7.01 (m, 2H), 6.75 (m, 3H), 5.69 (s, 2H), 5.26 (s, 2H), 4.47 (s, 2H), 2.24 (s, 3H). MS (ESI) 446 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(1-oxyquinolin-2-ylmethoxy)phenoxy]methyl}phenoxy}-acetonitrile (example 39k).

#### Example 41n

##### {2-[3-(6-Fluoroquinolin-2-ylmethoxy)phenoxy]methyl]-6-methylphenoxy}acetic acid

m.p. 160-161 °C. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.17 (m, 2H), 7.72 (d, 1H), 7.48 (m, 2H), 7.27 (m, 1H), 7.19 (d, 1H), 7.08 (m, 2H), 6.63 (s, 1H), 6.62 (d, 1H), 6.53 (d, 1H), 5.35 (s, 2H), 5.14 (s, 2H), 4.61 (s, 2H), 2.34 (s, 3H). MS (ESI) 448 (M+H)<sup>+</sup>. Prepared from {2-[3-(6-fluoroquinolin-2-ylmethoxy)phenoxy]methyl]-6-methylphenoxy}acetonitrile (example 39l).

#### Example 41o

##### {2-Methyl-6-[3-(1-methyl-4-oxo-1,4-dihydroquinolin-2-ylmethoxy)phenoxy]methyl}phenoxy}acetic acid

m.p. 192-194°C. <sup>1</sup>H NMR (300 MHz, 1:1 CDCl<sub>3</sub>: CD<sub>3</sub>OD): δ 8.29 (m, 1H), 7.76 (m, 2H), 7.41 (m, 1H), 7.25-6.92 (m, 4H), 6.81-6.41 (m, 4H), 5.11 (m, 4H), 4.39 (s, 2H), 3.88 (s, 3H), 2.27 (s, 3H). MS (ESI) 460 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(1-methyl-4-oxo-1,4-dihydroquinolin-2-ylmethoxy)phenoxy]methyl}phenoxy}acetonitrile (example 39m).

#### Example 41p

##### {4-Chloro-2-[3-(6-fluoroquinolin-2-ylmethoxy)phenoxy]methyl]-6-methylphenoxy}acetic acid

m.p. 140-141°C. <sup>1</sup>H NMR (300 MHz, 5:1 CDCl<sub>3</sub>: CD<sub>3</sub>OD): δ 8.20 (d, 1H), 8.07 (m, 1H), 7.69 (d, 1H), 7.50 (m, 2H), 7.25 (s, 1H), 7.16 (m, 1H), 7.11 (s, 1H), 6.67 (s, 1H), 6.60 (m, 2H), 5.30

(s, 2H), 5.07 (s, 2H), 4.24 (s, 2H), 2.25 (s, 3H). MS (ESI) 482, 484 (M+H)<sup>+</sup>, Cl pattern. Prepared from {4-chloro-2-[3-(6-fluoroquinolin-2-ylmethoxy)phenoxy]methyl}-6-methylphenoxy} acetonitrile (example 40a).

## 5 Example 41q

{2,4-Dichloro-6-[3-(6-fluoroquinolin-2-ylmethoxy)phenoxy]methyl}phenoxy} acetic acid  
m.p. 189-190°C. <sup>1</sup>H NMR (300 MHz, 5:1 CDCl<sub>3</sub>: CD<sub>3</sub>OD): δ 8.22 (d, 1H), 8.07 (m, 1H), 7.73 (d, 1H), 7.52 (m, 2H), 7.35 (m, 2H), 7.21 (m, 1H), 6.72 (s, 1H), 6.65 (m, 2H), 5.34 (s, 2H), 5.22 (s, 2H), 4.68 (s, 2H). MS (ESI) 502, 504, 506 (M+H)<sup>+</sup>, Cl<sub>2</sub> pattern. Prepared from {2,4-dichloro-6-[3-(6-fluoroquinolin-2-ylmethoxy)phenoxy]methyl}phenoxy} acetonitrile (example 40b).

## Example 41r

{2-Methyl-6-[3-(2-pyridin-2-yl-ethoxy)-phenoxy]methyl}-phenoxy}-acetic acid  
<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.50 (d, 1H), 7.73 (dt, 1H), 7.36 (d, 1H), 7.20 (m, 4H), 7.04 (m, 1H), 6.58 (m, 2H), 6.50 (d, 1H), 5.77 (s, 2H), 5.15 (s, 2H), 4.35 (m, 4H), 3.20 (t, 2H), 2.25 (s, 3H). MS (ESI) 394 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(2-pyridin-2-yl-ethoxy)-phenoxy]methyl}-phenoxy}-acetonitrile (example 21).

## Example 41s

{2-[3-(Quinolin-2-ylmethoxy)phenoxy]methyl}-6-methylphenoxy}-acetic acid  
m.p. 154-157 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.25 (d, 1H), 8.19 (d, 1H), 7.81 (d, 1H), 7.77-7.70 (m, 2H), 7.60-7.55 (m, 1H), 7.27 (dd, 1H), 7.18 (d, 1H), 7.13-7.04 (m, 2H), 6.85 (t, 1H), 6.61 (dd, 1H), 6.53 (dd, 1H), 5.40 (s, 2H), 5.18 (s, 2H), 4.62 (s, 2H), 2.35 (s, 3H); MS (ESI) 430 (M+H)<sup>+</sup>. Prepared from {2-[3-(quinolin-2-ylmethoxy)phenoxy]methyl}-6-methylphenoxy} acetonitrile (example 35).

## Example 41t

{2-Methyl-6-[3-(quinoxalin-2-ylmethoxy)-phenoxy]methyl}-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 9.07 (s, 1H), 8.14–8.05 (m, 2H), 7.87–7.82 (m, 2H), 7.19–7.05 (m, 3H), 6.97–6.92 (m, 1H), 6.77 (s, 1H), 6.67–6.57 (m, 2H), 5.41 (s, 2H), 5.19 (s, 2H), 4.10 (s, 2H), 2.21 (s, 3H); MS (ESI) 431 (M+H)<sup>+</sup>. Prepared from {2-[3-(quinoxalin-2-ylmethoxy)-phenoxy-methyl]-6-methylphenoxy} acetonitrile (example 39).

5

#### Example 41u

##### 2-{2-Methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxy-methyl]-phenoxy}-propionic acid

10

<sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 8.40 (d, 1H), 7.99 (dd, 2H), 7.77 (dd, 1H), 7.67–7.58 (m, 2H), 7.22–7.10 (m, 3H), 6.98 (dd, 1H), 6.73 (s, 1H), 6.60 (dd, 2H), 5.33 (s, 2H), 5.19 (dd, 2H), 4.40–4.34 (m, 1H), 2.24 (s, 3H), 1.35 (d, 3H); MS (ESI) 444 (M+H)<sup>+</sup>. Prepared from methyl 2-{2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxy-methyl]-phenoxy}-propionate (example 36a).

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

#### Example 41v

##### {2,4-Dichloro-6-[3-(quinolin-2-ylmethoxy)-phenoxy-methyl]-phenoxy}-acetic acid

m.p. 201–203 °C; <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>) δ 8.38 (d, 1H), 7.98 (dd, 2H), 7.76 (dd, 1H), 7.66–7.57 (m, 2H), 7.54 (d, 1H), 7.39 (d, 1H), 7.18 (dd, 1H), 6.75 (s, 1H), 6.66–6.61 (m, 2H), 5.33 (s, 2H), 5.28 (s, 2H), 4.48 (s, 2H); MS (ESI) 484, 486 (M+H; Cl<sub>2</sub>)<sup>+</sup>. Prepared from {2,4-dichloro-6-[3-(quinolin-2-ylmethoxy)-phenoxy-methyl]-phenoxy}-acetonitrile (example 36b).

#### Example 41w

##### {4-Chloro-2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxy-methyl]-phenoxy}acetic acid

25

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.37 (d, 1H), 8.02 (d, 1H), 7.93 (d, 1H), 7.78–7.70 (m, 2H), 7.60 (t, 1H), 7.24–7.15 (m, 3H), 6.73 (s, 1H), 6.66–6.61 (m, 2H), 5.34 (s, 2H), 5.14 (s, 2H), 4.27 (s, 2H), 2.29 (s, 3H); MS (ESI) 464 (M+H)<sup>+</sup>. Prepared from {4-chloro-2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxy-methyl]-phenoxy}acetonitrile (example 36c).

30

#### Example 41x

##### {2-tert-Butyl-6-[3-(quinolin-2-ylmethoxy)-phenoxy-methyl]-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.33 (d, 1H), 8.01 (d, 1H), 7.90 (d, 1H), 7.80-7.55 (m, 3H), 7.35-7.28 (m, 2H), 7.15-7.01 (m, 2H), 6.71 (s, 1H), 6.60 (d, 2H), 5.40 (s, 2H), 5.10 (s, 2H), 4.40 (s, 2H), 1.41 (s, 9H); MS (ESI) 472 (M+H)<sup>+</sup>. Prepared from {2-tert-butyl-6-[3-(quinolin-2-ylmethoxy)-phenoxy]methyl}-phenoxymethyl}-acetonitrile (example 36d).

5

## Example 41y

{4-Chloro-2-methoxy-6-[3-(quinolin-2-ylmethoxy)-phenoxy]methyl}-phenoxymethyl}-acetic acid

10 M.P. 185-190 °C, <sup>1</sup>H NMR (300 MHz, DMSO) δ 8.45 (d, 1H), 8.04 (t, 2H), 7.82 (t, 1H), 7.71-7.76 (m, 2H), 7.23 (t, 1H), 7.13 (d, 1H), 7.02 (d, 1H), 6.77 (d, 1H), 6.71-6.67 (m, 2H), 5.38 (s, 2H), 5.28 (s, 2H), 4.62 (s, 2H), 3.86 (s, 3H); MS (ESI) 480 (M+H)<sup>+</sup>. Prepared from {4-chloro-2-methoxy-6-[3-(quinolin-2-ylmethoxy)-phenoxy]methyl}-phenoxymethyl}-acetonitrile (example 36e).

15

## Example 41z

2-[3-(Quinolin-2-ylmethoxy)-phenoxy]methyl}-benzoic acid

20 <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.13 (d, 1H), 8.03 (d, 1H), 7.79 (d, 1H), 7.70-7.41 (m, 5H), 7.10 (t, 1H), 6.67 (s, 1H), 6.58 (d, 2H), 5.51 (s, 2H), 5.36 (s, 2H); MS (ESI) 386 (M+H)<sup>+</sup>. Prepared from 2-[3-(quinolin-2-ylmethoxy)-phenoxy]methyl}-benzonitrile (example 36f).

20

## Example 41aa

2-[3-(Quinolin-2-ylmethoxy)-phenoxy]methyl}-thiophene-2-carboxylic acid

25 <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.36 (d, 1H), 8.04 (d, 1H), 7.94 (d, 1H), 7.78 (t, 1H), 7.70 (d, 1H), 7.61 (t, 1H), 7.48 (d, 1H), 7.13-7.19 (m, 2H), 6.70 (s, 1H), 6.61 (dt, 2H), 5.46 (s, 2H), 5.32 (s, 2H); MS (ESI) 392 (M+H)<sup>+</sup>. Prepared from methyl 2-[3-(quinolin-2-ylmethoxy)-phenoxy]methyl}-thiophene-2-carboxylate (example 36g).

30

## Example 41ab

{2-[3-(Quinolin-2-ylmethoxy)-phenoxy]methyl}-phenyl}-acetic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.20-8.11 (m, 2H), 7.83-7.52 (m, 4H), 7.40-7.27 (m, 4H), 7.18-7.10 (m, 1H), 6.67-6.46 (m, 3H), 5.35 (s, 2H), 5.12 (s, 2H), 3.76 (s, 2H); MS 400 (M+H)<sup>+</sup>. Prepared from {2-[3-(quinolin-2-ylmethoxy)-phenoxy-methyl]-phenyl}-acetonitrile (example 38).

## 5 Example 41ac

### {4-Chloro-2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxymethyl]-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.19 (d, 1H), 8.09 (d, 1H), 7.82 (d, 1H), 7.78-7.65 (m, 2H), 7.75 (dd, 1H), 7.31-26 (dd, 1H), 7.18 (dd, 1H), 7.05-6.92 (m, 4H), 5.20 (s, 2H), 4.26 (s, 2H), 4.23 (s, 2H), 4.15 (s, 2H), 2.12 (s, 3H); MS (ESI) 478, 480 (M+H;Cl)<sup>+</sup>. Prepared from (example 52).

## Example 41ad

### {2-[3-(4-Chloro-quinolin-2-ylmethoxy)-phenoxy-methyl]-6-methyl-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.20 (dd, 1H), 8.08 (d, 1H), 7.91-7.74 (m, 3H), 7.24-7.15 (m, 3H), 7.01 (t, 1H), 6.74 (t, 1H), 6.66-6.59 (m, 2H), 5.32 (s, 2H), 5.12 (s, 3H), 4.43 (s, 2H), 2.24 (s, 3H); MS (ESI) 464 (M+H)<sup>+</sup>. Prepared from {2-[3-(4-chloro-quinolin-2-ylmethoxy)-phenoxy-methyl]-6-methyl-phenoxy}-acetonitrile (example 39n).

## Example 41ae

### {2-[3-(7-Chloro-quinolin-2-ylmethoxy)-phenoxy-methyl]-6-methyl-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.42 (d, 1H), 8.03-8.00 (m, 2H), 7.68-7.60 (m, 2H), 7.27-7.07 (m, 3H), 6.94 (t, 1H), 6.70 (d, 1H), 6.59 (dd, 2H), 5.31 (s, 2H), 5.18 (s, 2H), 2.20 (s, 3H); MS (ESI) 464 (M+H)<sup>+</sup>. Prepared from {2-[3-(7-chloro-quinolin-2-ylmethoxy)-phenoxy-methyl]-6-methyl-phenoxy}-acetonitrile (example 39o).

## Example 41af

### {2-[3-(6-Methoxy-quinolin-2-ylmethoxy)-phenoxy-methyl]-6-methyl-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.07 (t, 2H), 7.60 (d, 1H), 7.36 (dd, 1H), 7.27-7.24 (m, 1H), 7.15 (d, 1H), 7.05-7.00 (m, 3H), 6.79 (s, 1H), 6.58 (d, 1H), 6.49 (dd, 1H), 5.31 (s, 2H), 5.14 (s, 2H), 4.54

(s, 2H), 3.90 (s, 3H), 2.30 (s, 3H); MS (ESI) 460 (M+H)<sup>+</sup>. Prepared from {2-[3-(6-methoxy-quinolin-2-ylmethoxy)-phenoxymethyl]-6-methyl-phenoxy}-acetonitrile (example 39p).

#### Example 41ag

##### {2-[4-Bromo-3-(quinolin-2-ylmethoxy)-phenoxymethyl]-6-methyl-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.42 (d, 1H), 8.00-7.97 (m, 2H), 7.79-7.70 (m, 2H), 7.63-7.58 (m, 1H), 7.46 (d, 1H), 7.20-7.13 (m, 2H), 6.99-6.95 (m, 2H), 6.59 (dd, 1H), 5.42 (s, 2H), 5.17 (s, 2H), 4.30 (s, 2H), 2.22 (s, 3H); MS (ion spray) 508 (M+H)<sup>+</sup>. Prepared from ethyl {2-[4-bromo-3-(quinolin-2-ylmethoxy)-phenoxymethyl]-6-methyl-phenoxy}-acetate (example 54).

#### Example 41ah

##### {2-[2-Bromo-5-(quinolin-2-ylmethoxy)-phenoxymethyl]-6-methyl-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.18-8.15 (m, 2H), 7.81-7.72 (m, 2H), 7.63-7.53 (m, 2H), 7.37 (d, 1H), 7.28 (d, 1H), 7.17-7.13 (m, 1H), 7.09-7.03 (m, 1H), 6.91 (d, 1H), 6.42 (dd, 2H), 5.36 (s, 2H), 5.32 (s, 2H), 4.63 (s, 2H), 2.32 (s, 3H); MS (ion spray) 508 (M+H)<sup>+</sup>. Prepared from ethyl {2-[2-bromo-5-(quinolin-2-ylmethoxy)-phenoxymethyl]-6-methyl-phenoxy}-acetate (example 54).

#### Example 41ai

##### {2-Methyl-6-[3-methyl-5-(quinolin-2-ylmethoxy)-phenoxymethyl]-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.21-8.14 (m, 2H), 7.82-7.66 (m, 3H), 7.58 (t, 1H), 7.27-7.24 (m, 1H), 7.17 (d, 1H), 7.04 (t, 1H), 6.60 (s, 1H), 6.43 (s, 1H), 6.37 (s, 1H), 5.24 (s, 2H), 5.13 (s, 2H), 4.60 (s, 2H), 2.32 (s, 3H), 2.19 (s, 3H); MS (ion spray) 444 (M+H)<sup>+</sup>. Prepared from ethyl {2-methyl-6-[3-methyl-5-(quinolin-2-ylmethoxy)-phenoxymethyl]-phenoxy}-acetate (example 36j).

#### Example 41aj

##### {2-[2-Acetyl-5-(quinolin-2-ylmethoxy)-phenoxymethyl]-6-methyl-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.39 (d, 1H), 8.00-7.96 (m, 2H), 7.76 (t, 1H), 7.67-7.57 (m, 3H), 7.28 (d, 1H), 7.14 (d, 1H), 7.02-7.00 (m, 2H), 6.69 (d, 1H), 5.43 (s, 2H), 5.35 (s, 2H), 4.27 (s,

2H), 2.39 (s, 3H), 2.23 (s, 3H); MS (ion spray) 472 (M+H)<sup>+</sup>. Prepared from ethyl {2-[2-acetyl-5-(quinolin-2-ylmethoxy)-phenoxyethyl]-6-methyl-phenoxy}-acetate (example 59).

#### Example 41ak

##### {4-Chloro-2-methyl-6-[3-(2-pyridin-2-yl-ethoxy)-phenoxyethyl]-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.52 (d, 1H), 7.73 (m, 1H), 7.26 (m, 5H), 6.56 (m, 3H), 5.12 (s, 2H), 4.48 (s, 2H), 4.34 (t, 2H), 3.17 (t, 2H), 2.26 (s, 3H). MS (ESI) 428 (M+H)<sup>+</sup>. Prepared from {4-chloro-2-methyl-6-[3-(2-pyridin-2-yl-ethoxy)-phenoxyethyl]-phenoxy}-acetonitrile (example 36k).

#### Example 41al

##### {2-[3-(Benzooxazol-2-ylaminomethyl)-phenoxyethyl]-6-methyl-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.48 (bs, 1H), 7.12 (m, 11H), 5.14 (s, 2H), 4.46 (m, 4H), 2.26 (s, 3H). MS (ESI) 419 (M+H)<sup>+</sup>. Prepared from {2-[3-(benzooxazol-2-ylaminomethyl)-phenoxyethyl]-6-methyl-phenoxy}-acetonitrile (example 35d).

#### Example 41am

##### {2-[3-(Benzooxazol-2-ylaminomethyl)-phenoxyethyl]-4-chloro-6-methyl-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.47 (bt, 1H), 7.28 (m, 5H), 7.02 (m, 5H), 5.14 (s, 2H), 4.49 (d, 2H), 4.46 (s, 2H), 2.26 (s, 3H). MS (ESI) 453 (M+H)<sup>+</sup>. Prepared from {2-[3-(benzooxazol-2-ylaminomethyl)-phenoxyethyl]-4-chloro-6-methyl-phenoxy}-acetonitrile (example 36l).

#### Example 41an

##### {2-[3-(4-Chloro-quinolin-2-ylmethoxymethyl)-phenoxyethyl]-6-methyl-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.20 (d, 1H), 8.05 (d, 1H), 7.81 (m, 3H), 7.25 (m, 2H), 7.12 (m, 1H), 7.00 (m, 4H), 5.21 (s, 2H), 4.77 (s, 2H), 4.62 (s, 2H), 4.15 (s, 2H), 2.24 (s, 3H). MS (ESI) 478 (M+H)<sup>+</sup>. Prepared from {2-[3-(4-chloro-quinolin-2-ylmethoxymethyl)-phenoxyethyl]-6-methyl-phenoxy}-acetonitrile (example 35e).



## Example 41ao

{2-[3-(6-Methoxy-quinolin-2-ylmethoxymethyl)-phenoxy]methyl]-6-methyl-phenoxy}-acetic acid  
<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.18 (d, 1H), 8.10 (d, 1H), 7.64 (d, 1H), 7.36 (m, 2H), 7.26 (m, 3H), 7.12 (m, 2H), 6.98 (m, 2H), 5.24 (s, 2H), 4.73 (s, 2H), 4.69 (s, 2H), 4.62 (s, 2H), 3.93 (s, 3H), 2.38 (s, 3H). MS (ESI) 474 (M+H)<sup>+</sup>. Prepared from {2-[3-(6-methoxy-quinolin-2-ylmethoxymethyl)-phenoxy]methyl]-6-methyl-phenoxy}-acetonitrile (example 35f).

## Example 41ap

{2-Methyl-6-[3-(quinolin-2-ylmethoxymethyl)-phenoxy]methyl]-phenoxy}-acetic acid  
<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.30 (d, 1H), 8.18 (d, 1H), 7.85 (d, 1H), 7.74 (m, 2H), 7.58 (m, 1H), 7.38 (d, 1H), 7.28 (m, 3H), 7.12 (m, 1H), 6.99 (m, 2H), 5.24 (s, 2H), 4.78 (s, 2H), 4.71 (s, 2H), 4.63 (s, 2H), 2.39 (s, 3H). MS (ESI) 444 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(quinolin-2-ylmethoxymethyl)-phenoxy]methyl]-phenoxy}-acetonitrile (example 35g).

## Example 41aq

{2-Methyl-6-[3-(pyridin-4-ylmethoxy)-phenoxy]methyl]-phenoxy}-acetic acid  
<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.57 (bs, 2H), 7.44 (m, 2H), 7.22 (m, 3H), 7.07 (m, 1H), 6.69 (m, 1H), 6.61 (m, 2H), 5.15 (s, 2H), 5.13 (s, 2H), 4.47 (s, 2H), 2.27 (s, 3H). MS (ESI) 380 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(pyridin-4-ylmethoxy)-phenoxy]methyl]-phenoxy}-acetonitrile (example 39q).

## Example 41ar

{2-Methyl-6-[3-(pyridin-2-ylmethoxy)-phenoxy]methyl]-phenoxy}-acetic acid  
<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.58 (bd, 1H), 7.83 (m, 1H), 7.50 (d, 1H), 7.34 (m, 1H), 7.21 (m, 3H), 7.05 (m, 1H), 6.64 (m, 3H), 5.14 (s, 4H), 4.38 (s, 2H), 2.26 (s, 3H). MS (ESI) 380 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(pyridin-2-ylmethoxy)-phenoxy]methyl]-phenoxy}-acetonitrile (example 39r).

## Example 41as

{2-Methyl-6-[3-(pyridin-3-ylmethoxy)-phenoxy-methyl]-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.67 (s, 1H), 8.55 (bd, 1H), 7.86 (d, 1H), 7.43 (m, 1H), 7.28 (m, 3H), 7.07 (m, 1H), 6.70 (s, 1H), 6.62 (m, 2H), 5.13 (s, 4H), 4.47 (s, 2H), 2.27 (s, 3H). MS (ESI) 380 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(pyridin-3-ylmethoxy)-phenoxy-methyl]-phenoxy}-acetonitrile (example 39s).

## Example 41at

{2-[3-(6,7-Dichloro-quinolin-2-ylmethoxy)-phenoxy-methyl]-6-methyl-phenoxy}-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.44 (m, 2H), 8.30 (s, 1H), 7.76 (d, 1H), 7.22 (m, 3H), 7.04 (m, 1H), 6.72 (m, 1H), 6.63 (m, 2H), 5.35 (s, 2H), 5.13 (s, 2H), 4.46 (s, 2H), 2.26 (s, 3H). MS (ESI) 498 (M+H)<sup>+</sup>. Prepared from {2-[3-(6,7-dichloro-quinolin-2-ylmethoxy)-phenoxy-methyl]-6-methyl-phenoxy}-acetonitrile (example 39t).

## Example 41au

Ethyl 4-benzyloxy-2-[3-(2-carboxymethoxy-3-methyl-benzyloxy)-benzyloxy]-6-methyl-

benzoate <sup>1</sup>H NMR (300 MHz, DMSO) δ 7.31 (m, 8H), 7.06 (m, 2H), 6.96 (m, 2H), 6.65 (d, 1H), 6.54 (d, 1H), 5.12 (d, 6H), 4.48 (s, 2H), 4.22 (q, 2H), 2.28 (s, 3H), 2.19 (s, 3H), 1.21 (t, 3H). MS (ESI) 571 (M+H)<sup>+</sup>. Prepared from ethyl 4-benzyloxy-2-[3-(2-cyanomethoxy-3-methyl-benzyloxy)-benzyloxy]-6-methyl-benzoate (example 64b).

## Example 41av

4-Benzyloxy-2-[3-(2-carboxymethoxy-3-methyl-benzyloxy)-benzyloxy]-6-methyl-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 7.32 (m, 8H), 7.02 (m, 4H), 6.64 (d, 1H), 6.52 (d, 1H), 5.13 (m, 6H), 4.48 (s, 2H), 2.28 (s, 3H), 2.22 (s, 3H). MS (ESI) 543 (M+H)<sup>+</sup>. Prepared from ethyl 4-benzyloxy-2-[3-(2-carboxymethoxy-3-methyl-benzyloxy)-benzyloxy]-6-methyl-benzoate (example 41au).

## Example 41aw

{2-Methyl-6-[3-(1,3,3-trimethyl-2-oxo-2,3-dihydro-1H-indol-6-yloxymethyl)-phenoxy]-acetic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 7.21 (m, 5H), 6.99 (m, 3H), 6.75 (d, 1H), 6.64 (dd, 1H), 5.24 (s, 2H), 5.07 (s, 2H), 4.12 (s, 2H), 3.10 (s, 3H), 2.25 (s, 3H), 1.21 (s, 6H). MS (ESI) 476 (M+H)<sup>+</sup>.

- 5 Prepared from {2-methyl-6-[3-(1,3,3-trimethyl-2-oxo-2,3-dihydro-1H-indol-6-yloxymethyl)-phenoxy]-acetonitrile (example 64c).

Example 41ax

10 7-[3-(quinolin-2-ylmethoxy)-phenoxy]-benzofuran-2-carboxylic acid

<sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.36 (dd, 1H), 8.03 (dd, 1H), 7.94 (dd, 1H), 7.61-7.78 (m, 4H), 7.45-7.49 (m, 2H), 7.16-7.29 (m, 2H), 6.77 (s, 1H), 6.67 (dt, 2H), 5.42 (s, 2H), 5.32 (s, 2H); MS (ESI) 426 (M+H)<sup>+</sup>. Prepared from ethyl 7-[3-(quinolin-2-ylmethoxy)-phenoxy]-benzofuran-2-carboxylate (example 36i).

Example 41ay

{2-Methyl-6-[3-(2-phenyl-thiazol-4-ylmethoxy)-phenoxy]-acetic acid.

<sup>1</sup>H NMR (300 MHz, DMSO): δ 7.92 (m, 2H), 7.77 (s, 1H), 7.47 (m, 3H), 7.20 (m, 1H), 7.15 (m, 2H), 6.98 (t, 1H), 6.72 (t, 1H), 6.60 (m, 2H), 5.15 (s, 2H), 5.14 (s, 2H), 4.15 (s, 2H), 2.23 (s, 3H); MS (ESI) 462 (M+H)<sup>+</sup>. Prepared from {2-methyl-6-[3-(2-phenyl-thiazol-4-ylmethoxy)-phenoxy]-acetonitrile (example 39u).

25

Example 42

(4-Chloro-2,6-dimethyl-phenoxy)-acetonitrile

- 4-Chloro-2,6-dimethylphenol (5.0 g, 32 mmol), bromoacetonitrile (2.2 mL, 32 mmol) and potassium carbonate (6.6 g, 48 mmol) are combined with acetone (50 mL) and heated at reflux for 18 h. The reaction is filtered, concentrated and the residue partitioned between dichloromethane and water. The organic phase is washed with 1N HCl and water and is then

30

dried over magnesium sulfate, concentrated and purified by column chromatography (silica, 10% ethyl acetate in hexanes) to provide the title compound. MS (EI) 195 (M)<sup>+</sup>, Cl pattern.

#### Example 42a

##### (2-tert-Butyl-6-methylphenoxy)-acetonitrile

The title compound is prepared using essentially the same procedure used in example 42 except using 2-tert-butyl-6-methylphenol in place of 4-chloro-2,6-dimethylphenol. MS (EI) 203 (M)<sup>+</sup>.

#### Example 42b

##### Ethyl (2,6-dimethyl-phenoxy)-acetate

The title compound is prepared using essentially the same procedure used in example 42 except using 2,6-dimethylphenol in place of 4-chloro-2,6-dimethylphenol and ethyl bromoacetate in place of bromoacetonitrile.

#### Example 43

##### (2-Bromomethyl-4-chloro-6-methyl-phenoxy)-acetonitrile

(4-Chloro-2,6-dimethyl-phenoxy)-acetonitrile (700 mg, 3.6 mmol, example 42), N-bromosuccinimide (510 mg, 2.9 mmol) and benzoyl peroxide (72 mg, 0.29 mmol) are heated at reflux in carbon tetrachloride (10 mL) for 16 h. The reaction is cooled, filtered and the filtrate is concentrated and purified by column chromatography (silica, 5% ethyl acetate in hexanes) to provide the title compound. MS (EI) 273, 275(M)<sup>+</sup>, Br pattern.

The following compounds are prepared using essentially the same procedure used in example 43 except using the cited methyl analog in place of (4-chloro-2,6-dimethyl-phenoxy)-acetonitrile.

#### Example 43a

##### (2-Bromomethyl-6-tert-butyl phenoxy)-acetonitrile

MS (EI) 281 (M)<sup>+</sup>. Prepared from (2-tert-butyl-6-methylphenoxy)-acetonitrile (example 42a).

#### Example 43b

##### Ethyl (2-bromomethyl-6-methyl-phenoxy)-acetate

Prepared from ethyl (2,6-dimethyl-phenoxy)-acetate (example 42b).

#### Example 44

#### 5 5-Chloro-2-hydroxy-3-methoxy-benzaldehyde

A solution of sulfuryl chloride (15 mL, 190 mmol) in toluene (20 mL) is added dropwise over 1.5 h to a solution of o-vanillin (25.0 g, 164 mmol) in toluene (90 mL) and the reaction is then stirred 16 h. Water (30 mL) is added over 10 minutes with ice-bath cooling. The solid is filtered, washed with water and dried to provide the title compound. MS (EI) 186 (M)<sup>+</sup>.

#### Example 45

#### 4-Chloro-2-methyl-1-oxo-quinoline

mCPBA 70% pure (6.9 g, 29 mmol) is added to a solution of 4-chloroquinaldine (5.1 g, 29 mmol) in dichloroethane and heated to 50 °C for 4 h. The reaction is concentrated and partitioned between ethyl acetate and aqueous potassium carbonate. The organic phase is washed with additional aqueous potassium carbonate, water and is then dried over magnesium sulfate. The solution is filtered and concentrated to yield the title compound which is used without further purification. MS (ESI) 194 (M+H)<sup>+</sup>.

The following compounds are prepared using essentially the same procedure used in example 45 except using the cited quinaldine in place of 4-chloroquinaldine.

#### Example 45a

#### 7-Chloro-2-methyl-1-oxo-quinoline

MS (ESI) 194 (M+H)<sup>+</sup>. Prepared from 7-chloroquinaldine.

#### Example 45b

#### 6-Methoxy-2-methyl-1-oxo-quinoline

MS (ESI) 190 (M+H)<sup>+</sup>. Prepared from 6-methoxyquinaldine.

#### Example 45c

5-ethyl-2-methyl-pyridine 1-oxide

MS (ESI) 138 (M+H)<sup>+</sup>. Prepared from 5-ethyl-2-methylpyridine.

## 5 Example 46

4-Chloroquinolin-2-ylmethyl chloride

4-Chloro-2-methyl-1-oxo-quinoline (4.3 g, 22 mmol) is dissolved in chloroform (200 mL) and p-toluenesulfonyl chloride (3.7 g, 20 mmol) is added and the reaction is heated at 65 °C for 24 h.

10 The reaction is allowed to cool and then concentrated and partitioned between ethyl acetate and 10% aqueous potassium carbonate. The organic phase is dried over magnesium sulfate, concentrated and purified by column chromatography (silica, 60% dichloromethane in hexanes) to provide the title compound. MS (ESI) 212 (M+H)<sup>+</sup>.

15 The following compounds are prepared using essentially the same procedure used in example 45 except using the cited quinaldine in place of 4-chloroquinaldine.

## Example 46a

20 7-Chloroquinolin-2-ylmethyl chloride

MS (ESI) 212 (M+H)<sup>+</sup>. Prepared from 7-chloro-2-methyl-1-oxo-quinoline (example 45a).

## Example 46b

25 6-Methoxyquinolin-2-ylmethyl chloride

MS (ESI) 208 (M+H)<sup>+</sup>. Prepared from 6-methoxy-2-methyl-1-oxo-quinoline (example 45b).

## Example 47

30 2-{3-[5-Chloro-3-methyl-2-(1H-tetrazol-5-ylmethoxy)-benzyloxy]-phenoxy-methyl}-quinoline

Sodium azide (395 mg, 6.1 mmol) and ammonium chloride (325 mg, 6.1 mmol) are added to a solution of {4-chloro-2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxy-methyl]-phenoxy}-acetonitrile (300 mg, 0.68 mmol, example 36c) in DMF (2 mL) and heated at 110 °C for 2 h.

The reaction is then cooled and poured into a 1 N sodium hydroxide solution (20 mL) with the

formation of a solid. This mixture is then washed with ether (4x) and the ether is discarded. The remaining aqueous solution contains a solid which is filtered. This solid is dissolved in 10% ethanol water (250 mL) and the pH is lowered to about 5 with 2N HCl. A solid precipitates which is filtered to provide the title compound. m.p. 181-184 °C; <sup>1</sup>H NMR (300 MHz, DMSO-  
d<sub>6</sub>) δ 8.40 (d, 1H), 8.01-7.97 (m, 2H), 7.77 (dd, 1H), 7.66-7.60 (m, 2H), 7.33 (d, 2H), 7.18 (dd, 1H), 6.72 (dd, 1H), 6.65 (dd, 1H), 6.59 (dd, 1H), 5.33 (s, 2H), 5.27 (s, 2H), 5.07 (s, 2H), 2.24 (s, 3H); MS (ESI) 488, 490 (M+H;Cl)<sup>+</sup>.

#### Example 48

##### [3-(Quinolin-2-ylmethoxy)-phenyl]-methanol

2-Chloromethylquinoline hydrochloride (11.6 g, 54 mmol), 3-hydroxybenzyl alcohol (6.7 g, 54 mmol) and potassium carbonate (16 g, 116 mmol) are heated in DMF (45 mL) at 50 °C for 14 h. The temperature is increased to 80 °C and heated an additional 24 h. The reaction is cooled and added to water, filtered and the solid is washed with water to yield a semi-pure product. The residue is dissolved in ethyl acetate, dried over magnesium sulfate, filtered and concentrated. The sample is then recrystallized from ethyl acetate and hexanes to provide the title compound. MS (ESI) 266 (M+H)<sup>+</sup>.

#### Example 49

##### 2-(3-Chloromethyl-phenoxy)-quinoline hydrochloride

Thionyl chloride (0.95 mL, 13 mmol) is added to a solution of [3-(quinolin-2-ylmethoxy)-phenyl]-methanol (2.9 g, 11 mmol, example 48) in dichloromethane (30 mL) and allowed to stir 18 h. The reaction is concentrated *in vacuo* and azeotroped twice from chloroform to yield the title compound which is used without further purification.

#### Example 50

##### 2-[3-(Quinolin-2-ylmethoxy)-benzyloxy]-6-trifluoromethyl-benzaldehyde

[3-(Quinolin-2-ylmethoxy)-phenyl]-methanol (300 mg, 1.13 mmol, example 48) is dissolved in DMF (6 mL) and sodium hydride (60%, 60 mg, 1.5 mmol) is added and allowed to stir 20 min. 2-Fluoro-6-(trifluoromethyl)benzaldehyde (0.30 mL, 2.2 mmol) is added and the reaction is

heated at 90 °C for 5 h. The reaction is partitioned between ethyl acetate (200 mL) and water (200 mL), dried over magnesium sulfate, filtered, concentrated *in vacuo* and purified by column chromatography (silica, 25% ethyl acetate in hexanes) to provide the title compound; MS (ESI) 438 (M+H)<sup>+</sup>.

#### Example 51

##### 2-[3-(Quinolin-2-ylmethoxy)-benzyloxy]-6-trifluoromethyl-benzoic acid

A solution of 2-[3-(quinolin-2-ylmethoxy)-benzyloxy]-6-trifluoromethyl-benzaldehyde (46 mg, 0.1 mmol, example 50) in 2-methyl-2-butene (1 mL), t-butanol (2 mL) and water (2 mL) is treated with sodium dihydrogenphosphate dihydrate (153 mg, 1.1 mmol) and sodium chlorite (198 mg, 2.2 mmol). After 45 min. the reaction is partitioned between dichloromethane (50 mL) and water (50 mL). The organic layer is dried over magnesium sulfate, filtered and concentrated *in vacuo* to provide the title compound: m.p. 184-185 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.28 (d, 1H), 8.12 (d, 1H), 7.82-7.75 (m, 2H), 7.66-7.55 (m, 2H), 7.42 (dd, 1H), 7.30-7.27 (m, 2H), 7.16 (d, 1H), 7.07 (dd, 1H), 6.88 (d, 1H), 6.77 (dd, 1H), 5.44 (s, 2H), 5.07 (s, 2H); MS (ESI) 454 (M+H)<sup>+</sup>.

#### Example 52

##### {4-Chloro-2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxymethyl]-phenoxy}-acetonitrile

[3-(Quinolin-2-ylmethoxy)-phenyl]-methanol (190 mg, 0.72 mmol, example 48) is dissolved in DMF (6 mL) and sodium hydride (60%, 30 mg, 0.75 mmol) is added and allowed to stir for 10 min. (2-Bromomethyl-4-chloro-6-methyl-phenoxy)-acetonitrile (210 mg, 0.78 mmol, example 43) is added and the reaction is allowed to stir 6 h. The reaction is partitioned between ethyl acetate and water and the organic phase is washed with additional water. The organic phase is dried, concentrated and purified by column chromatography (silica, 25 % ethyl acetate in hexanes) to provide the title compound. MS (ESI) 458 (M+H)<sup>+</sup>, Cl<sup>-</sup> pattern.

#### Example 53

##### Methyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxymethyl]-benzoate



The title compound is prepared using essentially the same procedure used in example 52 except using methyl 2-bromomethyl-6-methyl-benzoate (example 2) in place of (2-bromomethyl-4-chloro-6-methyl-phenoxy)-acetonitrile. MS (ESI) 427 (M+H)<sup>+</sup>.

#### 5 Example 54

Ethyl {2-[4-bromo-3-(quinolin-2-ylmethoxy)-phenoxy]methyl}-6-methyl-phenoxy}-acetate and  
Ethyl {2-[2-bromo-5-(quinolin-2-ylmethoxy)-phenoxy]methyl}-6-methyl-phenoxy}-acetate

Ethyl {2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenoxy]methyl}-phenoxy}-acetate (350 mg, 0.76 mmol, example 36h), NBS (150 mg, 0.84) and benzoyl peroxide (20 mg, 0.08 mmol) are dissolved/suspended in chloroform (7 mL) and heated to reflux for 2 hrs. The reaction is cooled to r.t., filtered, preadsorbed onto silica gel by rotary evaporation. The crude material is purified by flash chromatography (silica, 15% ethyl acetate in hexanes) to give the title compounds in approximately a 1:1 ratio. Regioisomers are determined by NMR NOE data. MS (ion spray) 537 (M+H)<sup>+</sup> for both compounds.

#### Example 55

3-Methyl-5-(quinolin-2-ylmethoxy)-phenol

2-(Chloromethyl)-quinoline hydrochloride (1.28 g, 6.0 mmol), orcinol (568 mg, 4.0 mmol), K<sub>2</sub>CO<sub>3</sub> (1.68 g, 12.0 mmol) and a catalytic amount of tetrabutylammonium iodide (~ 10 mg) are dissolved/suspended in anhyd. DMF (10 mL) and heated at 50 °C overnight. The reaction is cooled to r.t. and partitioned between water (100 mL) and ethyl ether (100 mL). The pH of the aqueous layer is adjusted to ~ 5 and further extracted with ethyl ether (100 mL). The organic fractions are pooled and washed with brine (2 x 100 mL), dried over MgSO<sub>4</sub>, filtered and preadsorbed onto silica gel. The crude preadsorbed material is purified by flash chromatography (silica, 20% ethyl acetate in hexanes) to give the title compound. This product (approx. 80% purity, remainder is 2-methylquinoline) is used without further purification. MS (ESI) 266 (M+H)<sup>+</sup>.

#### Example 56

2-[3-(Quinolin-2-ylmethoxy)-benzyloxy]-benzaldehyde

2-(3-Chloromethyl-phenoxy)methyl-quinoline (371 mg, 1.3 mmol, example 49) and salicylaldehyde (133  $\mu$ L, 1.25 mmol) are dissolved in acetone (10 mL).  $K_2CO_3$  (525 mg, 3.75 mmol) is added and the contents are heated to reflux for 16 hrs. The reaction is cooled to r.t., poured into water (100 mL) and extracted with ethyl ether (3 x 50 mL). The ether layers are combined and washed with Brine (3 x 75 mL) and dried over  $MgSO_4$ . The crude material is preadsorbed onto silica gel and purified by flash chromatography (silica, 20 to 25% ethyl acetate in hexanes) to give the title compound. MS (ion spray) 370 (M+H)<sup>+</sup>.

#### Example 57

##### 3-{2-[3-(Quinolin-2-ylmethoxy)-benzyloxy]-phenyl}-acrylic acid

2-[3-(Quinolin-2-ylmethoxy)-benzyloxy]-benzaldehyde (110 mg, 0.3 mmol, example 56) is dissolved in pyridine (1 mL). Malonic acid (63 mg, 0.6 mmol) and piperidine (10  $\mu$ L, 0.1 mmol) are added and the contents are heated to 85 °C for 2 hrs, then at 110 °C for another 2 hrs. The reaction is cooled and placed under a nitrogen stream at 40 °C to remove the pyridine. A small amount of toluene is added and the contents are again placed under a nitrogen stream at 40 °C (repeat). The crude material is purified on silica gel by flash chromatography (silica, 2.5% methanol in dichloromethane) to give the title compound. <sup>1</sup>H NMR (300 MHz,  $CDCl_3$ )  $\delta$  8.33 (d, 1H), 8.21-8.16 (m, 2H), 7.81-7.70 (m, 3H), 7.56-7.50 (m, 2H), 7.37-7.25 (m, 3H), 7.03-6.93 (m, 4H), 6.50 (d, 1H), 5.48 (s, 2H), 5.14 (s, 2H); MS (ion spray) 412 (M+H)<sup>+</sup>.

#### Example 58

##### 1-[2-Hydroxy-4-(quinolin-2-ylmethoxy)-phenyl]-ethanone

2',4'-Dihydroxy-acetophenone (912 mg, 6 mmol) and 2-chloromethyl-quinoline hydrochloride (856 mg, 4.0 mmol, example 49) are dissolved in acetonitrile (20 mL).  $K_2CO_3$  (1.12 g, 8.0 mmol) is added and the contents are heated to 50 °C for 16 hrs. The reaction is cooled to room temperature, and the solvent is removed by rotary evaporation. The contents are partitioned between ethyl acetate (100 mL) and water (100 mL), the aqueous layer is acidified with 2 N HCl to ~ pH 2 and further extracted with ethyl acetate (2 x 50 mL). All organic fractions are combined and washed with brine (3 x 150 mL), dried over  $MgSO_4$  and concentrated. The crude material is preadsorbed onto silica gel and purified by flash chromatography (silica, 15% ethyl acetate in hexanes) to give the title compound; MS (ion spray) 294 (M+H)<sup>+</sup>.

## Example 59

Ethyl {2-[2-acetyl-5-(quinolin-2-ylmethoxy)-phenoxyethyl]-6-methyl-phenoxy}-acetate

1-[2-Hydroxy-4-(quinolin-2-ylmethoxy)-phenyl]-ethanone (185 mg, 0.63 mmol, example 58) is dissolved in 2:1 DMF/acetonitrile (6 mL). Ethyl (2-bromomethyl-6-methyl-phenoxy)-acetate (272 mg, 0.95 mmol, example 43b) and  $K_2CO_3$  (177 mg, 1.26 mmol) are added and the contents are heated to 50 °C for 2 days. The reaction is cooled to r.t. and the volume reduced under a nitrogen stream at 40 °C. The contents are partitioned between ethyl acetate (50 mL) and water (50 mL). The aqueous layer is further extracted with ethyl acetate (2 x 50 mL). The organic fractions are combined and washed with brine (3 x 75 mL), dried over  $MgSO_4$  and concentrated. The crude material is dissolved in 1:1 dichloromethane/methanol, preadsorbed onto silica gel, and purified by flash chromatography (silica, 20% ethyl acetate in hexanes) to give the title compound. MS (ion spray) 500 (M+H)<sup>+</sup>.

## Example 60

Methyl 2-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate

The free base of 2-(3-chloromethyl-phenoxyethyl)-quinoline hydrochloride (540 mg, 1.7 mmol, example 49) is prepared by partitioning the material between ethyl ether and sodium bicarbonate and drying the organic phase with magnesium sulfate. This material is then dissolved with methyl salicylate (260 mg, 1.7 mmol) in DMF (10 mL) at 0 °C and sodium hydride (60%, 65 mg 1.7 mmol) is added. The reaction is brought to room temperature for 15 min. and is then heated at 60 °C for 6 h. The reaction is cooled and partitioned between ethyl acetate and a saturated ammonium chloride solution. The organic phase is dried over magnesium sulfate, filtered, concentrated *in vacuo* and purified by column chromatography (silica, 50 to 80 % ether in hexanes) to provide the title compound; MS (ESI) 400 (M+H)<sup>+</sup>.

The following compounds are prepared using essentially the same procedure used in example 60 except using the cited substituted salicylate in place of methyl salicylate.

## Example 60a

Methyl 3-methoxy-2-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate

Prepared from methyl 3-methoxysalicylate.

Example 60b

Methyl 4-methoxy-2-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate

Prepared from methyl 4-methoxysalicylate.

Example 60c

Methyl 5-methoxy-2-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate

Prepared from methyl 5-methoxysalicylate.

Example 60d

Methyl 2-methoxy-6-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate

Prepared from methyl 6-methoxysalicylate (example 61).

Example 60e

Ethyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate

MS (ESI) 428 (M+H)<sup>+</sup>. Prepared from ethyl 6-methylsalicylate (See, Hauser, Frank M., *Synthesis* **1980**, 10, 814-15.

Example 61

Methyl 6-methoxysalicylate

A mixture of 6-methoxysalicylic acid (10.0 g, 59.5 mmol) in methanol (40 mL) and sulfuric acid (2 mL) is heated at reflux 48 h. Although some acid remains the reaction is concentrated to remove the methanol and partitioned between ethyl acetate and saturated sodium carbonate solution. The organic phase is separated and washed with sodium carbonate until no acid remains by TLC analysis. The organic phase is dried and concentrated to provide the title compound as a low melting solid.

## Example 62

Methyl 5-[3-(quinolin-2-ylmethoxy)-benzyloxy]-nicotinate

To a solution of 5-hydroxy nicotinic acid methyl ester (200 mg, 1.3 mmol) in DMF (3 mL) is added 60% sodium hydride emulsion (50 mg, 1.2 mmol) and this mixture is stirred 30 minutes. The free base of 2-(3-chloromethyl-phenoxymethyl)-quinoline hydrochloride (350 mg, 1.2 mmol, example 49) is prepared by partitioning the material between ethyl ether and sodium bicarbonate and drying the organic phase with magnesium sulfate. A solution of this free base in DMF (2 mL) is added to the alcohol and this mixture is stirred at 25 °C for 16 hours. The solvent is removed *in vacuo*, dichloromethane (10 mL) and water (5 mL) is added, and this mixture is acidified to pH6 with acetic acid. The organic layer is dried over magnesium sulfate and the solvent removed *in vacuo*. The residue is purified by flash chromatography (silica, 4% methanol in dichloromethane) to give the title compound. MS (ESI) 401 (M+H)<sup>+</sup>.

## Example 63

Ethyl 4-benzyloxy-2-hydroxy-6-methyl-benzoate

To a solution of ethyl-2,4-dihydroxy-6-methyl benzoate (4.22 g, 22 mmol) in acetone (80 mL) is added potassium carbonate (3.0 g, 22 mmol) and benzyl bromide (2.6 mL, 22 mmol) and this mixture is heated under reflux overnight. The cooled reaction is diluted with ethyl acetate (100 mL) and water (100 mL) and the organic layer washed with water (2 x 80 mL) and brine (2 x 80 mL). The organic layer is dried over magnesium sulfate and the solvent removed to provide the title compound without further purification. MS (EI) 286 (M)<sup>+</sup>.

## Example 63a

Ethyl 2-hydroxy-4-methoxy-6-methyl-benzoate

The title compound is prepared using essentially the same procedure used in example 63 except using iodomethane in place of benzyl bromide.

## Example 64

Ethyl 4-benzyloxy-2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate

To a solution of ethyl 4-benzyloxy-2-hydroxy-6-methyl-benzoate (5.1 g, 16 mmol, example 63) in DMF (100 mL), with 25 °C water bath cooling, is added 60% sodium hydride emulsion (1.3 g, 32 mmol) over 2 minutes. This mixture is stirred 30 minutes with the cooling bath removed. A solution of 3-(quinolin-2-ylmethoxy)-benzyl chloride hydrochloride (5.1g, 16 mmol, example 49), in DMF (55 mL) is added and the reaction heated at 60 °C for 6 hours. The solvent is removed *in vacuo* and the residue purified by flash chromatography (silica, 0.5 to 2% methanol in dichloromethane) to give the title compound. MS (ESI) 534 (M+H)<sup>+</sup>.

#### Example 64a

##### Ethyl 4-methoxy-2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate

The title compound is prepared using essentially the same procedure used in example 64 except using ethyl 2-hydroxy-4-methoxy-6-methyl-benzoate (example 63a) in place of ethyl 4-benzyloxy-2-hydroxy-6-methyl-benzoate. MS (ESI) 458 (M+H)<sup>+</sup>.

#### Example 64b

##### Ethyl 4-benzyloxy-2-[3-(2-cyanomethoxy-3-methyl-benzyloxy)-benzyloxy]-6-methyl-benzoate

The title compound is prepared using essentially the same procedure used in example 64 except using [2-(3-bromomethyl-phenoxy)-6-methyl-phenoxy]-acetonitrile (example 76) in place of 3-(quinolin-2-ylmethoxy)-benzyl chloride hydrochloride. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.30 (m, 8H), 7.12 (m, 1H), 7.06 (bs, 1H), 6.99 (d, 1H), 6.90 (dd, 1H), 6.42 (s, 2H), 5.07 (s, 2H), 5.04 (s, 2H), 5.02 (s, 2H), 4.71 (s, 2H), 4.33 (q, 2H), 2.39 (s, 3H), 2.30 (s, 3H), 1.31 (t, 3H). MS (ESI) 552 (M+H)<sup>+</sup>.

#### Example 64c

##### {2-Methyl-6-[3-(1,3,3-trimethyl-2-oxo-2,3-dihydro-1H-indol-6-ylloxymethyl)-phenoxy]-acetonitrile}

The title compound is prepared using essentially the same procedure used in example 64 except using [2-(3-bromomethyl-phenoxy)-6-methyl-phenoxy]-acetonitrile (example 76) in place of 3-(quinolin-2-ylmethoxy)-benzyl chloride hydrochloride and 6-hydroxy-1,3,3-trimethyl-1,3-dihydro-indol-2-one (example 80) in place of ethyl 4-benzyloxy-2-hydroxy-6-methyl-benzoate.

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.32 (dd, 2H), 7.24 (s, 1H), 7.11 (m, 4H), 6.96 (dd, 1H), 6.62 (dd, 1H), 6.54 (d, 1H), 5.10 (s, 2H), 5.06 (s, 2H), 4.72 (s, 2H), 3.18 (s, 3H), 2.39 (s, 3H), 1.34 (s, 6H). MS (ESI) 457 (M+H)<sup>+</sup>.

5 Example 65

Ethyl 2-hydroxy-6-methyl-4-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate

2,4-Dihydroxy-6-methyl-benzoic acid ethyl ester (315 mg, 1.6 mmol) is combined with 2-(3-chloromethyl-phenoxymethyl)-quinoline hydrochloride (0.51 g, 1.6 mmol, example 49),  
10 tetrabutylammonium iodide (55 mg, 0.15 mmol) and potassium carbonate (0.48 g, 3.5 mmol) in acetone (9 mL). The reaction is heated at reflux 48 h. The reaction is partitioned between ethyl acetate and saturated ammonium chloride. The organic phase is washed with brine, dried over magnesium sulfate, filtered and concentrated to yield the crude product. This material is purified by column chromatography (silica, 3% ether in dichloromethane) to the title compound; m.p. 127-128 °C, MS (ESI) 444 (M+H)<sup>+</sup>.

Example 66

Ethyl 2-methoxy-6-methyl-4-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate

To a solution of ethyl 2-hydroxy-6-methyl-4-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate (150 mg, 0.34 mmol, example 65) in DMF (5 mL) is added sodium hydride (60%, 14 mg, 0.34 mmol) and the reaction is stirred 20 min. Iodomethane (0.03 mL, 0.5 mmol) is added and the reaction is heated at 50 °C for 7 h. The reaction is concentrated *in vacuo* and the residue is partitioned between dichloromethane and aqueous ammonium chloride. The water layer is back-  
25 extracted with dichloromethane, the organic phases are combined, dried over magnesium sulfate, filtered and concentrated to yield crude product. The residue is purified by column chromatography (silica, 10 to 20% ethyl acetate in hexanes to provide the title compound; MS (ESI) 458 (M+H)<sup>+</sup>.

30 Example 66a

Ethyl 2-benzyloxy-6-methyl-4-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate

The title compound is prepared using essentially the same procedure used in example 66 except using benzyl bromide in place of iodomethane. MS (ESI) 534 (M+H)<sup>+</sup>.

#### Example 67

##### 4-Benzyloxy-2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoic acid

Ethyl 4-benzyloxy-2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate (2.4 g, 4.5 mmole, example 64) is added to ethanol (50 ml) and 10N sodium hydroxide (4.4 ml, 44 mmole) and refluxed for 8 hours. The solvent is removed *in vacuo* and the residue is dissolved in dichloromethane with a small amount of water and is acidified to pH6 with 1N HCl. The organic layer is dried over MgSO<sub>4</sub> and the solvent removed *in vacuo*. The crude product is purified by column chromatography (silica, 1% methanol in dichloromethane) to provide the title compound. m.p. 146-149°C; <sup>1</sup>H NMR (300 MHz, CD<sub>3</sub>OD) δ 8.34 (d, 1H), 8.04 (d, 1H), 7.91 (d, 1H), 7.78-7.70 (m, 2H), 7.61 (t, H), 7.37-7.20 (m, 7H), 7.05-6.91 (m, 2H), 6.51 (d, 1H), 6.47 (d, 1H), 5.35 (s, 2H), 5.09 (s, 2H), 5.03 (s, 2H), 2.29 (s, 3H); MS (ESI) 506 (M+H)<sup>+</sup>.

The following compounds are prepared using essentially the same procedure used in example 67 except using the cited ester in place of ethyl 4-benzyloxy-2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate.

#### Example 67a

##### 2-Methoxy-6-methyl-4-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoic acid

<sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.38 (d, 1H), 8.03 (d, 1H), 7.79 (d, H), 7.70-7.67 (m, 2H), 7.61 (t, H), 7.28 (t, 1H), 7.14 (s, 1H), 7.01 (t, 2H), 6.35 (t, 2H), 5.37 (s, 2H), 5.03 (s, 2H), 3.71 (s, 3H), 2.26 (s, 3H); MS (ESI) 430 (M+H)<sup>+</sup>. Prepared from ethyl 2-methoxy-6-methyl-4-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate (example 66).

#### Example 67b

##### 2-Benzyloxy-6-methyl-4-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoic acid

125-127 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.18 (d, 1H), 7.83 (d, 1H), 7.74-7.55 (m, 3H), 7.40 (s, 5H), 7.39-7.29 (m, 1H), 7.09 (s, 1H), 7.00 (m, 2H), 6.50 (s, 2H), 5.41 (s, 2H), 5.13 (s, 2H),



5.04 (s, 2H), 2.58 (s, 3H); MS (ESI) 506 (M+H)<sup>+</sup>. Prepared from ethyl 2-benzyloxy-6-methyl-4-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate (example 66a).

#### Example 67c

##### 4-Methoxy-2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoic acid

<sup>1</sup>H NMR (300 MHz, DMSO) δ 8.39 (d, 1H), 8.02-7.96 (m, 2H), 7.79-7.74 (m, 1H), 7.67-7.57 (m, 2H), 7.31-7.25 (m, 1H), 7.16 (s, 1H), 7.02-6.96 (m, 2H), 6.49 (s, 1H), 6.39 (s, 1H), 5.34 (s, 2H), 5.09 (s, 2H), 3.71 (s, 2H), 2.20 (s, 3H); MS (ESI) 429 (M+H)<sup>+</sup>. Prepared from ethyl 4-methoxy-2-methyl-6-[3-(quinolin-2-ylmethoxy)-benzyloxy]-benzoate (example 64a).

#### Example 68

##### 5-ethyl-2-chloromethyl pyridine

To a solution of 5-ethyl-2-methyl-pyridine-1-oxide (427 mg, 3.11 mmol, example 45c) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) is added a solution (0.2 mL) of phosphorous (v) trichloride oxide (327 μL) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL). Added simultaneously the remaining phosphorous (v) trichloride oxide solution and a solution of triethylamine (488 μL) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) at such a rate as to maintain a reflux. After the addition is complete, let reaction mixture cool to 20 °C and diluted with EtOAc. The organic layer is washed with sat. NaHCO<sub>3</sub> soln., brine, dried over MgSO<sub>4</sub> and concentrated. The residue is purified by flash chromatography (silica, 10% ethyl acetate in dichloromethane) to provide the title compound. MS (ESI) 156 (M+H)<sup>+</sup>.

#### Example 69

##### 2-(5-Ethyl-pyridin-2-yl)-ethanol

To a cooled solution (-10°C) of diisopropylamine (2.31 mL, 16.5mL) in THF ( 45 mL) is added dropwise (2.5M) n-butyllithium (6.6 mL, 16.5 mmol), let stir 10 min. then cooled to -78°C. To this mixture is added dropwise a solution of 5-ethyl-2-methylpyridine (1.98 mL, 15 mmol) in THF (3 mL) and let stir for 10min at -78°C. To the reaction mixture is added paraformaldehyde (1.13 g, 37.5 mmol), the cold bath removed and stirring continued for 1 hr. Quenched reaction with H<sub>2</sub>O, diluted with EtOAc and the organic layer washed with brine, dried over MgSO<sub>4</sub> and

concentrated. The residue is purified by flash chromatography (silica, 0.5% ammonia / 5% methanol / dichloromethane) to give the title compound as a pale yellow oil. MS (EI) 151 (M)<sup>+</sup>.

#### Example 69a

##### 2-quinolin-2-yl-ethanol

The title compound is prepared using essentially the same procedure used in example 69 except using 2-methyl-quinoline in place of 5-ethyl-2-methylpyridine. MS (ESI) 174 (M+H)<sup>+</sup>.

#### Example 70

##### Benzoic acid 3-[2-(5-ethyl-pyridin-2-yl)-ethoxy]-phenyl ester

To a solution of 2-(5-ethyl-pyridin-2-yl)-ethanol (480 mg, 3.17 mmol, example 69) in THF (10 mL) is added resorcinol monobenzoate (630 mg, 2.94 mmol), triphenylphosphine (850 mg, 3.24 mmol) and diethyl azodicarboxylate (510  $\mu$ L, 3.24 mmol). The resulting mixture is stirred for 1 h then concentrated. The residue is purified by flash chromatography (silica, 35% ethyl acetate in hexane) to give the title compound as a yellow oil. MS (ESI) 348 (M+H)<sup>+</sup>.

#### Example 70a

##### Benzoic acid 3-(2-pyridin-2-yl-ethoxy)-phenyl ester

The title compound is prepared using essentially the same procedure used in example 70 except using 2-(2-hydroxyethyl)pyridine in place of 2-(5-ethyl-pyridin-2-yl)-ethanol. MS (ESI) 320 (M+H)<sup>+</sup>.

#### Example 71

##### 3-[2-(5-ethyl-pyridin-2-yl)-ethoxy]-phenol

To a solution of benzoic acid 3-[2-(5-ethyl-pyridin-2-yl)-ethoxy]-phenyl ester (493 mg, 1.42 mmol, example 70) in 1:1 THF / CH<sub>3</sub>OH (5 mL) is added 10N NaOH soln. (0.5 mL) and water (50  $\mu$ L). The reaction mixture is stirred for 15 min then cooled to 5°C, adjusted to pH 7 with 2N HCl soln. and diluted with EtOAc. The organic layer is washed sequentially with brine, sat

NaHCO<sub>3</sub> soln. then dried over MgSO<sub>4</sub> and concentrated. The residue is purified by several triturations with hexane to give the title compound as a crystalline solid. MS (ESI) 244 (M+H)<sup>+</sup>.

#### Example 71a

##### 3-(2-pyridin-2-yl-ethoxy)-phenol

The title compound is prepared using essentially the same procedure used in example 71 except using benzoic acid 3-(2-pyridin-2-yl-ethoxy)-phenyl ester (example 70a) in place of 3-[2-(5-ethyl-pyridin-2-yl)-ethoxy]-phenyl ester. MS (ESI) 216 (M+H)<sup>+</sup>.

#### Example 72

##### [3-(2-Methoxy-ethoxymethoxy)-phenyl]-methanol

To a cooled suspension (0°C) of 60% NaH (660 mg, 16.5 mmol) in THF (35 mL) is added dropwise a solution of 3-hydroxybenzaldehyde (1.89 g, 15 mmol) in THF (15 mL) and the resulting mixture stirred for 20 min. To the mixture is added 2-methoxyethoxymethyl chloride (1.88 mL, 16.5 mmol) and DMPU (5 mL), the cold bath removed and stirred for 1 hr. The reaction mixture is cooled to 0 °C then slowly added 2M NaBH<sub>4</sub> (in triglyme) (3.75 mL, 7.5 mmol) and let stir for 1 hr. Slowly quenched with 2N HCl soln (3.9 mL) and diluted reaction mixture with ether. The organic layer is washed with brine, dried over MgSO<sub>4</sub> and concentrated. The residue is purified by flash chromatography (silica, 60% ethyl acetate in hexanes) to give the title compound as a pale yellow oil. MS (EI) 212 (M)<sup>+</sup>.

#### Example 73

##### 2-[3-(2-Methoxy-ethoxymethoxy)-benzyloxymethyl]-pyridine

To a cooled solution (0°C) of [3-(2-methoxy-ethoxymethoxy)-phenyl]-methanol (212 mg, 1 mmol, example 72) in THF (3 mL) is added 60% NaH (80 mg, 2 mmol) and the mixture stirred 10 min. Added 2-picolyl chloride hydrochloride (164 mg, 1 mmol) and DMPU (0.8 mL), removed cold bath and let reaction mixture stir for 2 hrs. Quenched reaction with sat NH<sub>4</sub>Cl soln. and diluted with EtOAc. The organic layer is washed with brine, dried over MgSO<sub>4</sub> and concentrated. The residue is purified by flash chromatography (silica, 70% ethyl acetate in hexanes) to give the title compound as a pale yellow oil. MS (ESI) 304 (M+H)<sup>+</sup>.

The following compounds are prepared using essentially the same procedure used in example 73 except using the cited halide in place of 2-picoyl chloride hydrochloride.

5 Example 73a

2-[3-(2-Methoxy-ethoxymethoxy)-benzyloxymethyl]-quinoline

MS (ESI) 354 (M+H)<sup>+</sup>. Prepared from 2-(chloromethyl)quinoline hydrochloride.

10 Example 73b

4-Chloro-2-[3-(2-methoxy-ethoxymethoxy)-benzyloxymethyl]-quinoline

MS (ESI) 388 (M+H)<sup>+</sup>. Prepared from 2-chloromethyl-4-chloroquinoline (example 46).

Example 73c

6-Methoxy-2-[3-(2-methoxy-ethoxymethoxy)-benzyloxymethyl]-quinoline

MS (ESI) 384 (M+H)<sup>+</sup>. Prepared from 2-chloromethyl-6-methoxyquinoline (example 46b).

Example 74

3-(Pyridin-2-ylmethoxymethyl)-phenol

To a solution of 2-[3-(2-methoxy-ethoxymethoxy)-benzyloxymethyl]-pyridine (171 mg, 0.56 mmol, example 73) in CH<sub>3</sub>OH (1.9 mL) is added p-toluenesulfonic acid monohydrate (148 mg, 0.78 mmol). The mixture is heated to 60°C and stirred for 1.5 hrs, then cooled to room temperature and diluted with EtOAc. The organic layer is washed with sat NaHCO<sub>3</sub>, brine, then dried over MgSO<sub>4</sub> and concentrated to give the title compound as a white crystalline solid. MS (ESI) 216 (M+H)<sup>+</sup>.

30 The following compounds are prepared using essentially the same procedure used in example 74 except using the cited MEM ether in place of 2-[3-(2-methoxy-ethoxymethoxy)-benzyloxymethyl]-pyridine.

Example 74a

3-(Quinolin-2-ylmethoxymethyl)-phenol

MS (ESI) 266 (M+H)<sup>+</sup>. Prepared from 2-[3-(2-methoxy-ethoxymethoxy)-benzyloxymethyl]-quinoline (example 73a).

## Example 74b

3-(4-Chloro-quinolin-2-ylmethoxymethyl)-phenol

MS (ESI) 300 (M+H)<sup>+</sup>. Prepared from 4-chloro-2-[3-(2-methoxy-ethoxymethoxy)-benzyloxymethyl]-quinoline (example 73b).

## Example 74c

3-(6-Methoxy-quinolin-2-ylmethoxymethyl)-phenol

MS (ESI) 296 (M+H)<sup>+</sup>. Prepared from 6-methoxy-2-[3-(2-methoxy-ethoxymethoxy)-benzyloxymethyl]-quinoline (example 73c).

## Example 75

[2-(3-Hydroxymethyl-phenoxymethyl)-6-methyl-phenoxy]-acetonitrile

To a solution of 3-hydroxybenzyl alcohol (202 mg, 1.63 mmol) in DMF (5.4 mL) is added K<sub>2</sub>CO<sub>3</sub> (247 mg, 1.79 mmol) and (2-bromomethyl-6-methyl-phenoxy)-acetonitrile (430 mg, 1.79 mmol, example 24). Heated resulting mixture to 60°C and stirred for 3 hrs then cooled to room temp and diluted with ether. Washed organic layer with water, brine, dried over MgSO<sub>4</sub> and concentrated. The residue is purified by flash chromatography (silica, 30% ethyl acetate in hexanes) to give the title compound. MS (EI) 283 (M)<sup>+</sup>.

## Example 76

[2-(3-Bromomethyl-phenoxymethyl)-6-methyl-phenoxy]-acetonitrile

To a solution of [2-(3-hydroxymethyl-phenoxymethyl)-6-methyl-phenoxy]-acetonitrile (230 mg, 0.81 mmol, example 75) in THF (3 mL) is added Ph<sub>3</sub>P (233 mg, 0.89 mmol) and stirred until homogeneous. Cooled solution to 0°C then added portionwise NBS (151 mg, 0.85 mmol) and let stir 45 min. Concentrated reaction mixture under reduced pressure. The residue is purified by

flash chromatography (silica, 40% CH<sub>2</sub>Cl<sub>2</sub> in hexanes) to give the title compound as a white crystalline solid. MS (EI) 345, 347 (M)<sup>+</sup>, Br pattern.

#### Example 77

##### 6-Methoxy-3-methyl-1,3-dihydro-indol-2-one

To a cooled solution (-78°C) of 6-methoxy-1,3-dihydro-indol-2-one (840 mg, 5.2 mmol, See Quallich, *Synthesis* **1993**, 51-53) in THF (20 mL) is added dropwise TMEDA (1.57 mL, 10.4 mL) followed by dropwise addition of 2.5M n-BuLi (4.16 mL, 10.4 mmol). The mixture is allowed to stir for 15 min then warmed to -25°C. Iodomethane (405 µL, 6.5 mmol) is added dropwise and stirred for 20 min. The reaction is quenched with sat NH<sub>4</sub>Cl soln, warmed to room temp and diluted with EtOAc. Washed organic layer with sat NH<sub>4</sub>Cl soln, brine, dried over MgSO<sub>4</sub> and concentrated. The residue is purified by flash chromatography (silica, 45% ethyl acetate in hexanes) to give the title compound. MS (ESI) 178 (M+H)<sup>+</sup>.

#### Example 78

##### 6-Methoxy-3,3-dimethyl-1,3-dihydro-indol-2-one

To a cooled solution (-78°C) of 6-methoxy-3-methyl-1,3-dihydro-indol-2-one (679 mg, 3.83 mmol, example 77) in THF (13 mL) is added TMEDA (1.16 mL, 7.66 mmol) followed by dropwise addition of 2.5M n-BuLi (3.06 mL, 7.66 mmol). The mixture is stirred 15 min then warmed to -25°C. Iodomethane (275 µL, 4.40 mmol) is added dropwise and stirred for 30 min. Reaction is quenched with sat NH<sub>4</sub>Cl soln, warmed to room temp and diluted with EtOAc. Washed organic layer with sat NH<sub>4</sub>Cl soln, brine, dried over MgSO<sub>4</sub> and concentrated. The residue is purified by flash chromatography (silica, 35% ethyl acetate in hexanes) to give the title compound as a white crystalline solid. MS (ESI) 192 (M+H)<sup>+</sup>.

#### Example 79

##### 6-Methoxy-1,3,3-trimethyl-1,3-dihydro-indol-2-one

To a cooled solution (-5°-0°C) of 6-methoxy-3,3-dimethyl-1,3-dihydro-indol-2-one (600 mg, 3.14 mmol, example 78) in THF (10.5 mL) is added 60% NaH (132 mg, 3.30 mmol) and is stirred for 15 min. Iodomethane (215 µL, 3.45 mmol) is added to the reaction mixture and stirred

for 2 hrs. Quenched reaction with sat  $\text{NH}_4\text{Cl}$  soln and diluted with EtOAc. Washed organic layer with sat  $\text{NH}_4\text{Cl}$  soln, brine, dried over  $\text{MgSO}_4$  and concentrated. The residue is purified by flash chromatography (silica, 30% ethyl acetate in hexanes) to give the title compound as a white crystalline solid. MS (ESI) 206 ( $\text{M}+\text{H}$ )<sup>+</sup>.

5

## Example 80

6-Hydroxy-1,3,3-trimethyl-1,3-dihydro-indol-2-one

To a solution of 6-methoxy-1,3,3-trimethyl-1,3-dihydro-indol-2-one (601 mg, 2.93 mmol, example 79) in acetic acid (880  $\mu\text{L}$ ) is added hydrobromic acid (48% in  $\text{H}_2\text{O}$ ) (8.8 mL). The resulting solution is heated to reflux (105°-110°C), stirred 2 hrs, then cooled to room temp and concentrated under reduced pressure. The residue is dissolved in EtOAc and the organic layer washed with water, brine, dried over  $\text{MgSO}_4$  and concentrated. The residue is purified by triturating with a small volume of ether to give the title compound as an off white solid. MS (ESI) 192 ( $\text{M}+\text{H}$ )<sup>+</sup>.

10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995

## Example 81

2-[3-(2-Methoxy-ethoxymethoxy)-benzyloxy]-quinoline

To a suspension of 60% NaH (44 mg, 1.1 mmol) in DMSO (2 mL) is added dropwise a solution of [3-(2-methoxy-ethoxymethoxy)-phenyl]-methanol (212 mg, 1.0 mmol, example 72) in DMSO (1 mL). Let stir 20 min. then added 2-chloroquinoline (180 mg, 1.1 mmol) and heated to 100 °C for 1 hr. Cooled reaction mixture to room temp. and diluted with EtOAc. The organic layer is washed with sat.  $\text{NH}_4\text{Cl}$  soln., brine, dried over  $\text{MgSO}_4$  and concentrated. The residue is purified by flash chromatography (silica, 25% ethyl acetate in hexanes) to give the title compound as a colorless oil. MS (ESI) 340 ( $\text{M}+\text{H}$ )<sup>+</sup>.

25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995

## Example 82

Isobutyl 2-[3-(methoxy)-phenylsulfanylmethyl]-6-methyl-benzoate

A 10N solution of sodium hydroxide (0.32 mL, 3.2 mmol) is added slowly to a solution of 3-methoxybenzenethiol (0.42 g, 3.0 mmol) in isobutanol (2 mL) followed by a solution of isobutyl 2-bromomethyl-6-methyl-benzoate (0.96 g, 3.3 mmol, example 2) in isobutanol (2 mL). The

30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995

reaction is allowed to stir 15 min and is then partitioned between ethyl acetate and dilute aqueous HCl. The organic phase is washed with water, dried over magnesium sulfate, concentrated and purified by column chromatography (silica, 40% dichloromethane in hexanes) to provide the title compound. MS (EI) 344 (M)<sup>+</sup>.

5

## Example 83

Isobutyl 2-[3-(hydroxy)-phenylsulfanylmethyl]-6-methyl-benzoate

Boron tribromide (1.3 mL, 1.0 M in dichloromethane, 1.3 mmol) is added to a solution of isobutyl 2-[3-(methoxy)-phenylsulfanylmethyl]-6-methyl-benzoate (194 mg, 0.56 mmol, example 82) in dichloromethane (3 mL) at 0 °C and then the reaction is stirred at room temperature 3h. The reaction is then partitioned between sodium bicarbonate solution and ethyl acetate. The organic phase is dried over magnesium sulfate, concentrated and purified by column chromatography (silica, 15 % ethyl acetate in hexanes) to provide the title compound. MS (EI) 330 (M)<sup>+</sup>.

10

15

## Example 84

Isobutyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenylsulfanylmethyl]-benzoate

The free base of 2-(chloromethyl)quinoline hydrochloride (148 mg, 0.69 mmol) is prepared by partitioning the material between ethyl ether and sodium bicarbonate and drying the organic phase with magnesium sulfate. This material is then dissolved with isobutyl 2-[3-(hydroxy)-phenylsulfanylmethyl]-6-methyl-benzoate (220 mg, 0.67 mmol, example 83) in DMF (2 mL) at 0 °C and sodium hydride (60%, 27 mg 0.67 mmol) is added. The reaction is allowed to stir 16 h and is then partitioned between ethyl acetate and water. The organic phase is washed with water (3X), dried over magnesium sulfate, concentrated and purified by column chromatography (silica, 10% ethyl acetate in hexanes) to provide the title compound. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.19 (d, 1H), 8.08 (d, 1H), 7.83 (d, 1H), 7.74 (t, 1H), 7.63 (d, 1H), 7.55 (t, 1H), 7.18-7.07 (m, 4H), 6.99 (d, 1H), 6.89 (d, 1H), 6.82 (dd, 1H), 5.33 (s, 2H), 4.18 (s, 2H), 4.10 (d, 2H), 2.36 (s, 3H), 2.07-2.01 (m, 1H), 0.98 (d, 6H); MS (ESI) 472 (M+H)<sup>+</sup>.

25

30

## Example 85



Isobutyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenylsulfanylmethyl]-benzoate

m-Chloroperbenzoic acid (<86%, 34 mg, 0.17 mmol) is added to a solution of isobutyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenylsulfanylmethyl]-benzoate (80 mg, 0.17 mmol, example 84) in dichloromethane (1 mL) and the reaction is stirred overnight. The reaction is partitioned between ethyl acetate and sodium bicarbonate and the organic phase is washed with additional bicarbonate solution, dried over magnesium sulfate, concentrated and purified by column chromatography (silica, 40 % ethyl acetate in hexanes) to provide the title compound. MS (ESI) 488 (M+H)<sup>+</sup>.

## Example 86

Isobutyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenylsulfonylmethyl]-benzoate

m-Chloroperbenzoic acid (<86%, 62 mg, 0.31 mmol) is added to a solution of isobutyl 2-methyl-6-[3-(quinolin-2-ylmethoxy)-phenylsulfanylmethyl]-benzoate (73 mg, 0.16 mmol, example 85) in dichloromethane (1 mL) and the reaction is stirred overnight. The reaction is partitioned between ethyl acetate and sodium bicarbonate and the organic phase is washed with additional bicarbonate solution, dried over magnesium sulfate, concentrated and purified by column chromatography (silica, 30 % ethyl acetate in hexanes) to provide the title compound. MS (ESI) 504 (M+H)<sup>+</sup>.

## Example 87

(1-Quinolin-2-ylmethyl-1H-imidazol-4-yl)-methanol and (3-quinolin-2-ylmethyl-3H-imidazol-4-yl)-methanol

2-Chloromethyl-quinoline hydrochloride (2.24 g, 10.5 mmol), 4-(hydroxymethyl)-imidazole hydrochloride (1.35 g, 10 mmol) and K<sub>2</sub>CO<sub>3</sub> (4.2 g, 30 mmol) are dissolved/suspended in anhyd. DMF (20 mL) and heated to 100 °C with rapid stirring overnight. The reaction is cooled to r.t. and poured into water (400 mL) and extracted with chloroform (3 x 150 mL). The organic fractions are pooled and washed with brine (2 x 200 mL), dried over MgSO<sub>4</sub>, filtered and reduced under vacuum to an oil. The crude material is purified by flash chromatography (silica, 5% methanol in dichloromethane) to give (3-quinolin-2-ylmethyl-3H-imidazol-4-yl)-methanol and (1-quinolin-2-ylmethyl-1H-imidazol-4-yl)-methanol in a 2:3 ratio. The identity of each

regioisomer was determined by NMR NOE experiments. MS (ESI) 240 (M+H)<sup>+</sup> found for both regioisomers.

#### Example 88

Isobutyl 2-methyl-6-(1-quinolin-2-ylmethyl-1H-imidazol-4-ylmethoxymethyl)-benzoate  
(1-Quinolin-2-ylmethyl-1H-imidazol-4-yl)-methanol (350 mg, 1.46 mmol, example 87) is dissolved in 20% DMPU in THF (5 mL) and cooled to 0 °C. Sodium hydride (60%, 60 mg, 1.50 mmol) is added portionwise, and the contents stirred for 15 min. Isobutyl 2-bromomethyl-6-methyl-benzoate (57%, 730 mg, 1.46 mmol, example 2) is added, the reaction is allowed to come to r.t. and stirred overnight. The contents are poured into water (200 mL) and extracted with dichloromethane (3 x 75 mL). The organic fractions are pooled and washed with brine (3 x 100 mL), dried over MgSO<sub>4</sub>, filtered and reduced under vacuum to an oil. The crude material is purified by flash chromatography (silica, 3% methanol in dichloromethane) to give the title compound. MS (ESI) 443 (M+H)<sup>+</sup>.

#### Example 88a

Isobutyl 2-methyl-6-(3-quinolin-2-ylmethyl-3H-imidazol-4-ylmethoxymethyl)-benzoate  
The title compound is prepared using essentially the same procedure used in example 88 except using (3-quinolin-2-ylmethyl-3H-imidazol-4-yl)-methanol in place of (1-quinolin-2-ylmethyl-1H-imidazol-4-yl)-methanol. MS (ESI) 443 (M+H)<sup>+</sup>.

#### Example 89

2-Methyl-6-(1-quinolin-2-ylmethyl-1H-imidazol-4-ylmethoxymethyl)-benzoic acid  
Isobutyl 2-methyl-6-(1-quinolin-2-ylmethyl-1H-imidazol-4-ylmethoxymethyl)-benzoate (300 mg, 0.68 mmol, example 88) is dissolved in ethanol (5 mL). 10 N NaOH (680 µL, 6.8 mmol) is added and the contents heated to 90 °C overnight. The reaction is cooled to r.t., 2 N HCl (3.4 mL, 6.8 mmol) is added and the pH adjusted to ~ 5 - 7. The contents are poured into water (100 mL) and extracted with chloroform (3 x 75 mL). The organic fractions are pooled, washed with brine (3 x 100 mL), dried over MgSO<sub>4</sub>, filtered and reduced under vacuum to an oil. The crude material is purified by HPLC (C-18, 25-50% acetonitrile in water over 15 min.) to give the title

compound as the TFA salt.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.93 (s, 1H), 8.31 (d, 1H), 8.07 (d, 1H), 7.86 (d, 1H), 7.78 (t, 1H), 7.63 (t, 1H), 7.51 (d, 1H), 7.44 (s, 1H), 7.17-7.12 (m, 1H), 7.06-7.01 (m, 2H), 5.64 (s, 2H), 4.66 (s, 2H), 4.48 (s, 2H), 2.29 (s, 3H). MS (ESI) 388 ( $\text{M}+\text{H}$ ) $^+$ .

#### 5 Example 89a

##### 2-Methyl-6-(3-quinolin-2-ylmethyl-3H-imidazol-4-ylmethoxymethyl)-benzoic acid

The title compound is prepared using essentially the same procedure used in example 89 except using isobutyl 2-methyl-6-(3-quinolin-2-ylmethyl-3H-imidazol-4-ylmethoxymethyl)-benzoate in place of isobutyl 2-methyl-6-(1-quinolin-2-ylmethyl-1H-imidazol-4-ylmethoxymethyl)-benzoate.

$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.95 (s, 1H), 8.27 (d, 1H), 8.05 (d, 1H), 7.82-7.76 (m, 2H), 7.64-7.59 (m, 1H), 7.46 (s, 1H), 7.38 (d, 1H), 7.10-6.99 (m, 2H), 6.92 (d, 1H), 5.91 (s, 2H), 4.52 (s, 2H), 4.49 (s, 2H), 2.23 (s, 3H). MS (ESI) 388 ( $\text{M}+\text{H}$ ) $^+$ .

#### Example 90

##### 2-[3-(1H-Indol-3-ylmethyl)-phenoxyethyl]-quinoline

Indole (230 mg, 2.0 mmol) is dissolved in tetrahydrofuran (3 mL) and ethylmagnesium bromide (1 M, 2.0 mL, 2.0 mmol) is added and the reaction is heated for 2 h at 65 °C. The free base of 2-(3-chloromethyl-phenoxyethyl)-quinoline hydrochloride (400 mg, 1.2 mmol, example 49) is prepared by partitioning the material between ethyl ether and sodium bicarbonate and drying the organic phase with magnesium sulfate. This free base is dissolved in tetrahydrofuran (2 mL) and is added to the cooled indole/Grignard solution, along with catalytic tetrabutylammonium iodide. This mixture is heated 6 h at 65 °C. The reaction is then cooled and partitioned between ethyl ether and ammonium chloride. The organic phase is washed with brine, dried over magnesium sulfate, concentrated and purified by column chromatography (silica, dichloromethane) to yield the title compound. MS (ESI) 365 ( $\text{M}+\text{H}$ ) $^+$ .

#### Example 91

##### {3-[3-(quinolin-2-ylmethoxy)-benzyl]-indol-1-yl}-acetic acid

Sodium hydride (60%, 22 mg, 0.55 mmol) is added to a solution of 2-[3-(1H-indol-3-ylmethyl)-phenoxyethyl]-quinoline (90 mg, 0.25 mmol, example 90) in DMF (2.5 mL). After stirring 5

minutes ethyl bromoacetate (0.1 mL, 0.9 mmol) is added and the reaction is allowed to stir 2h. The reaction is partitioned between ethyl acetate and ammonium chloride and the organic phase is washed with water. The organic phase is dried over magnesium sulfate, concentrated and then the solid is titrated with ethyl ether and ethyl acetate to provide the title compound as a solid.

5 m.p. 151-159 °C; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.20 (t, 2H), 7.83-7.80 (m, 1H), 7.77-7.71 (m, 1H), 7.67 (d, 1H), 7.61-7.53 (m, 1H), 7.45-7.42 (m, 1H), 7.29-7.14 (m, 3H), 7.05-6.99 (m, 2H), 6.94 (s, 1H), 6.85 (m, 2H), 5.16 (s, 2H), 4.84(s, 2H), 4.11 (s, 2H); MS (ESI) 423 (M+H)<sup>+</sup>.

#### Example 92

##### Ethyl (2-formyl-6-methyl-2-phenoxy)-acetate

Ethyl bromoacetate (4.5 mL, 40 mmol), 2-hydroxy-3-methyl-benzaldehyde (5 g, 37 mmol and potassium carbonate (8.1 g, 59 mmol) are combined in acetone (60 mL) and refluxed overnight. The reaction is filtered and the solvent removed from the filtrate under reduced pressure to give the title compound. MS (GC-MS) 222 (M)<sup>+</sup>.

#### Example 93

##### Ethyl 7-methyl-benzofuran-2-carboxylate

Sodium (0.52 g, 23 mmol) is dissolved in ethanol (60 mL) and to this is added ethyl (2-formyl-6-methyl-2-phenoxy)-acetate (5 g, 23 mmol, example 92). This mixture is refluxed for 3 hours and the solvent is removed in vacuo. The residue is dissolved in dichloromethane/water and acidified with 1N HCl. The organic layer is washed with water and brine and then dried over magnesium sulfate and the solvent removed in vacuo. The residue is purified by flash  
25 chromatography (silica, 1% methanol in dichloromethane) to give the title compound. MS (GC-EI) 176 (M)<sup>+</sup>.

#### Example 94

##### Ethyl 7-bromomethyl-benzofuran-2-carboxylate

Ethyl 7-methyl-benzofuran-2-carboxylate (0.5 g, 2.4 mmol, example 93), N-bromosuccinimide (0.48 g, 2.7 mmol) and benzoyl peroxide (0.06 g, 2.4 mmol) are combined in carbon tetrachloride (10 mL) and heated in an oil bath at 90°C overnight. The reaction is filtered and the

filtrate solvent is removed in vacuo. The residue is purified by flash chromatography (silica, 5% to 10% ethyl acetate in hexanes) to give the title compound. MS (GC-EI) 360, 362 ( $M^+$ , Br pattern).

#### 5 Example 95

##### Ethyl 2-methyl-6-trifluoromethanesulfonyloxy benzoate

Ethyl 6-methylsalicylate (2.5 g, 14 mmol, See, Hauser, Frank M., *Synthesis* **1980**, 10, 814-15) is dissolved in THF (20 mL) under nitrogen and cooled in an ice bath. Sodium hydride (60%, 0.56 g, 14 mmol) is added and the mixture is stirred 15 minutes. Then DMPU (0.20 mL) and N-phenyl-trifluoromethanesulfonimide (5.0 g, 14 mmol) are added and the reaction is stirred with cooling for 2 hours. The solvent is removed *in vacuo* and ether is added and the organics are washed with water, then dried over magnesium sulfate and concentrated under reduced pressure. The residue is purified by flash chromatography (silica, dichloromethane) to give the title compound.

#### Example 96

##### 3-(2-methoxy-ethoxymethoxy)-phenyliodide

To a suspension of 60% sodium hydride (1.76 g, 44 mmol) in THF (10 mL), cooled to 0°C, is added 3-iodophenol (8.8 g, 40 mmol) and methoxyethoxymethyl chloride (5 mL, 44 mmol) in THF (50 mL). Then DMPU (10 mL) is added, the cooling bath is removed and the reaction is stirred for an hour. The reaction is diluted with ether, washed with water and brine and the organic layer dried over magnesium sulfate. The solvent is removed *in vacuo* to give the title compound.

#### Example 97

##### [3-(2-Methoxy-ethoxymethoxy)-phenylethynyl]-trimethyl silane

3-(2-Methoxy-ethoxymethoxy)-phenyliodide (12.1 g, 39 mmol, 96) and tetrakis(triphenylphosphine)palladium (1.2 g, 1.0 mmol) and cuprous iodide (0.096 g, 0.5 mmol) are dissolved in THF (120 mL) and to this is added piperidine (12 mL) and (trimethylsilyl)acetylene (8 mL, 57 mmol). This mixture is degassed and is then stirred for 2 hours. The reaction is then diluted with ether

and washed twice with water and brine and the organic layer dried over magnesium sulfate. The solvent is removed *in vacuo* to give the title compound. MS (EI) 206 (M)<sup>+</sup>.

#### Example 98

##### Ethyl-2-[3-(2-methoxy-ethoxymethoxy)-phenylethynyl]-6-methyl-benzoate

[3-(2-Methoxy-ethoxymethoxy)-phenylethynyl]-trimethyl silane (0.57 g, 2 mmol, example 97) and 1.0 M tetrabutylammonium fluoride (2.1 mL, 2 mmol) are added to THF (10 mL) and acetic acid (0.13 g, 2 mmol) is added and this mixture is stirred at 20°C, under nitrogen. After 15 minutes, the solvent is removed in vacuo and the residue is azeotroped with benzene and purified by flash chromatography (silica, 20% ethyl acetate, 30% dichloromethane in hexanes) to give 1-ethynyl-3-(2-methoxy-ethoxymethoxy)-benzene (0.28 g, 1.4 mmol) which is dissolved in THF (8 mL), cooled to -78°C, under nitrogen and to this solution is added 2.5M n-butyl lithium (0.56 mL, 1.4 mmol) dropwise over 30 seconds. After stirring for 15 minutes, 1.0M zinc chloride in ether (1.4 mL, 1.4 mmol) is added dropwise over 30 seconds and this mixture is stirred for 30 minutes. Bis-(dibenzylideneacetone) palladium (0.04 g, 0.07 mmol) and bis (diphenyl phosphino) ferrocene (0.04 g, 0.07 mmol) is added and to this mixture is added ethyl-2-methyl-6-trifluoromethanesulfonyloxy benzoate (0.44 g, 1.4 mmol, 95) in THF (2 mL). The cooling bath is removed and the reaction allowed to warm to room temperature. Then the reaction mixture is heated in an oil bath at 65°C overnight. The reaction is then diluted with ethyl acetate (50 mL), washed with saturated ammonium chloride and brine and then dried over magnesium sulfate. The solvent is removed in vacuo and the residue purified by flash chromatography (silica, 10% ethyl acetate, 25% dichloromethane in hexanes) to give the title compound. MS (ESI) 369 (M+H)<sup>+</sup>.

#### Example 99

##### (3-Methyl-4-oxo-3,4-dihydro-quinazolin-2-yl)-methyl chloride.

To a suspension of isatoic anhydride (1.63g, 10 mmol) in dioxane (40 mL) is added methylamine solution (5 mL, 2M in THF). The resulting solution is stirred for 1h then concentrated under vacuum. The residue is taken up in toluene (30 mL) then pyridine (5.5 mL) is added followed by a solution of chloroacetyl chloride (2.7 mL, 34 mmol) in toluene (15 mL).

The resulting mixture is stirred for 15 h. The solid product is filtered, washed with water, then dried under vacuum to give 2.1 g of a tan solid. A portion of this product (452 mg, 2 mmol) is suspended in benzene (10 mL) then *p*-toluene sulphonic acid monohydrate (394 mg, 2 mmol) is added. This mixture is warmed to 70 °C and stirred at this temperature for 10 h. The mixture is then cooled to room temperature and the benzene solution decanted. The residual solid is mixed with sodium bicarbonate solution (sat.) and this mixture is extracted with ethyl acetate / methanol / dichloromethane. The combined extracts are washed with brine, dried over MgSO<sub>4</sub> and concentrated to give the title compound as a tan solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.28 (d, 1H), 7.75 (t, 1H), 7.67 (d, 1H), 7.51 (t, 1H), 4.62 (s, 3H), 3.76 (s, 3H).

#### Example 100

##### 3-(2-Hydroxymethyl-3-methyl-benzyloxy)-phenol

To a cooled (0 °C) solution of methyl 2-methyl-6-[(3-hydroxy-phenoxy)-methyl]-benzoate (220 mg, 0.76 mmol, example 5) in THF (2 mL) is added lithium aluminum hydride solution (1.5 mL, 1M in THF). The resulting solution is stirred for 10 min then warmed to room temperature and stirred for 40 min. This solution is then cooled to 0 °C and water (75 mL) added, dropwise, followed by sodium hydroxide solution (75 mL, 5N) and water (75 mL). The resulting suspension is diluted with ether, filtered through celite and the solid washed thoroughly with methanol (until the solid is free of product by TLC analysis). The combined filtrates are concentrated under vacuum to give the title compound as a white solid. MS (EI) 244 (M)<sup>+</sup>.

#### Example 101

##### 2-[3-(2-Hydroxymethyl-3-methyl-benzyloxy)-phenoxymethyl]-3-methyl-3H-quinazolin-4-one

To a solution of 3-(2-hydroxymethyl-3-methyl-benzyloxy)-phenol (87 mg, 0.38 mmol, example 100) and (3-methyl-4-oxo-3,4-dihydro-quinazolin-2-yl)-methyl chloride (94 mg, 0.45 mmol, example 99) in DMF (1 mL) is added powdered K<sub>2</sub>CO<sub>3</sub> (78 mg, 0.5 mmol). The resulting mixture is warmed to 60 °C and stirred at this temperature for 5h. This mixture is cooled to room temperature, diluted with ethyl acetate, washed with water and brine, dried over MgSO<sub>4</sub> and concentrated. The residue is purified by flash chromatography (silica, 40% ethyl acetate / 30% dichloromethane in hexanes) to give the title compound as a foam. MS (ESI) 417 (M+H)<sup>+</sup>.

## Example 101a

{2-[3-(5-Cyclobutyl-[1,2,4]oxadiazol-3-ylmethoxy)-phenoxy-methyl]-6-methyl-phenyl}-methanol

- 5 The title compound is prepared using essentially the same procedure used in example 101 except using 3-chloromethyl-5-cyclobutyl-[1,2,4]oxadiazole in place of (3-methyl-4-oxo-3,4-dihydro-quinazolin-2-yl)-methyl chloride. MS (ESI) 381 (M+H)<sup>+</sup>

## 10 Example 102

2-Methyl-6-[3-(3-methyl-4-oxo-3,4-dihydro-quinazolin-2-ylmethoxy)-phenoxy-methyl]-benzaldehyde

To a cooled (-78 °C) solution of oxalyl chloride (2.5 mL, 1.75 M in CH<sub>2</sub>Cl<sub>2</sub>) is added, dropwise, DMSO (80 mL). On complete addition, a solution of 2-[3-(2-hydroxymethyl-3-methyl-benzyloxy)-phenoxy-methyl]-3-methyl-3H-quinazolin-4-one (120 mg, 0.28 mmol, example 101) in dichloromethane (1 mL) is added dropwise. This solution is stirred for 5 min then triethylamine (276 mL, 2mmol) is added in one portion. The cold bath is removed and stirring continued for 10 min. The mixture is then diluted with ethyl acetate, washed with water and brine, dried over MgSO<sub>4</sub> and concentrated to give the title compound as a solid. MS (ESI) 415 (M+H)<sup>+</sup>.

## Example 102a

25 2-[3-(5-Cyclobutyl-[1,2,4]oxadiazol-3-ylmethoxy)-phenoxy-methyl]-6-methyl-benzaldehyde

The title compound is prepared using essentially the same procedure used in example 102 except using {2-[3-(5-cyclobutyl-[1,2,4]oxadiazol-3-ylmethoxy)-phenoxy-methyl]-6-methyl-phenyl}-methanol (example 101a) in place of 2-[3-(2-hydroxymethyl-3-methyl-benzyloxy)-phenoxy-methyl]-3-methyl-3H-quinazolin-4-one. MS (ESI) 379 (M+H)<sup>+</sup>.

30

## Example 103

2-Methyl-6-[3-(3-methyl-4-oxo-3,4-dihydro-quinazolin-2-ylmethoxy)-phenoxy-methyl]-benzoic acid



To a suspension of 2-methyl-6-[3-(3-methyl-4-oxo-3,4-dihydro-quinazolin-2-ylmethoxy)-phenoxymethyl]-benzaldehyde (120 mg, 0.28 mmol, example 102) in *t*-butanol (1.5 mL) is added iso-butene (0.5 mL) followed by NaClO<sub>2</sub> (220 mg, tech.grade 1.6 mmol) in water (1.5 mL) and NaH<sub>2</sub>PO<sub>4</sub>·H<sub>2</sub>O (220 mg, 1.6 mmol) in water (1.5 mL). This mixture is stirred for 1h (during which time the solids dissolve) then diluted with ethyl acetate, washed with water and brine, dried over MgSO<sub>4</sub> and concentrated. The residue is purified by flash chromatography (10% methanol in dichloromethane). This product was suspended in chloroform and filtered through celite. The filtrate is concentrated under reduced pressure to give the title compound as an amorphous solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 8.41 (d, 1H), 7.84 (m, 2H), 7.62 (m, 1H), 7.33 (m, 2H), 7.20 (m, 1H), 7.14 (t, 1H), 6.81 (m, 1H), 6.70 (m, 2H), 5.29 (s, 2H), 5.25 (s, 2H), 3.80 (s, 2H), 2.52 (s, 3H). MS (ESI) 430 (M+H)<sup>+</sup>.

#### Example 103a

##### 2-[3-(5-Cyclobutyl-[1,2,4]oxadiazol-3-ylmethoxy)-phenoxymethyl]-6-methyl-benzoic acid

The title compound is prepared using essentially the same procedure used in example 103 except using 2-[3-(5-Cyclobutyl-[1,2,4]oxadiazol-3-ylmethoxy)-phenoxymethyl]-6-methyl-benzaldehyde (example 102a) in place of 2-methyl-6-[3-(3-methyl-4-oxo-3,4-dihydro-quinazolin-2-ylmethoxy)-phenoxymethyl]-benzaldehyde. <sup>1</sup>H NMR (300MHz, DMSO) δ 7.10 (m, 4H), 6.68 (s, 1H), 6.60 (m, 2H), 5.19 (s, 2H), 5.13 (s, 2H), 3.86 (m, 1H), 2.36 (m, 4H), 2.28 (s, 3H), 2.08 (m, 1H), 1.96 (m, 1H). MS (ESI) 395 (M+H)<sup>+</sup>.

#### Example 104


##### 5-Phenyl-2-methylpyridine

To a cooled (-70°C) solution of 3-phenylpyridine (1.43 mL, 10.0 mmoles) in diethyl ether (7.5 mL) is added dropwise methylolithium (LiBr complex, 1.5 M in diethyl ether, 7.33 mL, 11.0 mmoles). After letting warm to room temperature over 16 hours the reaction is cooled (0°C) and quenched with distilled water (5 mL). The reaction is then extracted with methylene chloride, the organic layer isolated and concentrated, and the resulting residue purified by column chromatography (silica, 3:1 hexane: EtOAc) to yield the title compound as a pale yellow oil. MS(ESI) 170 (M+H)<sup>+</sup>.


### Synthesis of a compound of Formula (VI)

A compound of Formula (VI) is prepared in a multi-step synthesis illustrated in the below scheme. The key starting material is quinaldine. In the first stage it is chlorinated to form 2-chloromethylquinaldine which, without isolation, is reacted with hydroquinone to form the intermediate 4-(quinolin-2-yl-methoxy)phenol (VIII). This intermediate is then treated with  $\alpha, \alpha'$ -dichloro-*o*-xylene to form 2-[4-quinolin-2-yl-methoxy]phenoxyethyl]benzyl chloride, which is converted *in situ* to 2-[4-quinolin-2-yl-methoxy]phenoxyethyl]phenylacetonitrile (IX), the penultimate precursor to (VI).


(IX) is converted to (VI) crude, in a reaction with sodium azide and ammonium chloride which transforms the nitrile group into the tetrazole ring. The purification of the final product is accomplished by recrystallization of the crude material from methanol to afford pure (VI).



### Solid Phase Synthesis of a Compound of Formula:




#### 1. Acid Loading:



A 1L round bottom flask is charged with 4-(bromomethyl)benzoic acid (32.26 g, 150.0 mmole) and dichloromethane (650 mL). A stir bar is carefully added and the reaction flask is immersed in an ice-water bath. After approximately 15 minutes, oxalyl chloride (15.7 mL, 180 moles) is added. After approximately 15 minutes, N,N-dimethylformamide (500 mL, cat.) is added. The reaction began to bubble. After stirring for 1.5 hours, the ice-water bath is removed. After stirring for 3 hours at ambient temperature, the effervescence has ceased. At the end of this period, the stirbar is removed from the reaction mixture and the reaction solvent is removed *in vacuo*. After the solvent has been removed, more dichloromethane is added to the reaction flask and this too is removed *in vacuo*.


A three neck 3L round bottom flask is charged with dry N,N-dimethylformamide (1.3 L), N,N-diisopropylethylamine (39.19 mL, 225 mmole), 4-N,N-dimethylaminopyridine (3.67 g, 30 mmole) and MicroKANs [1456, 15 mg of Wang resin (1.7 mmole/g loading) per MicroKANs, 25.5 micromoles/microKAN, 37.1 mmole]. The flask is fitted with an overhead stirring apparatus. After stirring for approximately 15 minutes, a solution of the acid chloride as prepared above in dry N,N-dimethylformamide (200 mL) is transferred into the reaction flask. After 14 hours, the reaction solvent is removed. DMF (1.5 L) is added to the reaction flask. The flask was allowed to stir for approximately 15 minutes and the solvent is drained. The MicroKANs are washed, stirred for 20 minutes and drained in the following sequence repeatedly: DMF (2 x 6 L), THF (3 x 6 L), dichloromethane (3 x 6 L) and ether (2 x 6 L). After the final washing the MicroKANs are dried by blowing a stream of nitrogen through the flask with intermittent agitation. After sufficient drying, the MicroKANs are sorted for the next reaction.

## 2. Phenol Displacement:



A three neck 3L round bottom flask is charged with 3-chloro-4-hydroxybenzaldehyde (21.9 g, 140 mmol) and DMF (1.5 L). The reaction flask is fitted with an overhead stirrer and immersed in an ice-water bath. After approximately 15 minutes sodium hydride (60 % dispersion in oil, 6.48 g, 180 mmol) is carefully added. After approximately 30 minutes, the ice-water bath is removed and the reaction allowed to stir at ambient temperature for 1 hour. At the end of this time, the MicroKANs [1274, 25.5 micromol/microKAN, 32.5 mmol] and potassium iodide (1.0 g) are added to the reaction mixture. The reaction flask is immersed into an oil bath which is heated to 60°C. After 14 hours, the reaction flask is removed from the oil bath and allowed to cool to ambient temperature. The reaction solvent is removed. DMF (1.2 L) is added to the reaction flask. The flask is allowed to stir for approximately 15 minutes and the solvent is drained. DMF : water (1:1, 1.2 L) is added to the reaction flask. The flask is allowed to stir for approximately 15 minutes and the solvent is drained. This sequence is repeated at least three times or until the effluent from the washing is clear, the reaction flasks are washed repeatedly in the following sequence: THF (2 x 4 L), dichloromethane (1 x 4 L) then methanol (1 x 4 L), dichloromethane (1 x 4 L) then methanol (1 x 4 L), dichloromethane (1 x 4 L) and ether (1 x 4 L). After the final washing the MicroKANs are dried by blowing a stream of nitrogen through the flask with intermittent agitation. After sufficient drying, the MicroKANs are sorted for the next reaction.


### 3. Reductive Amination:



A three neck 2 L round bottom flask is charged with the MicroKANs [784, 25.5 micromol/microKAN, 20.0 mmol], trimethylorthoformate (850 mL) and 2-(2-aminoethyl)pyridine

20.79 g, 170 mmoles). The reaction flask is fitted with an overhead stirrer. After 2 hours, sodium cyanoborohydride (21.37 g, 340 mmoles) is added. After approximately 10 minutes, acetic acid (17.0 mL, 297 mmoles) is added. After stirring for an additional hour, the reaction flask is drained. Methanol (800 mL) is added to the flask. After stirring for approximately 10 minutes, the flask is drained. the reaction flask is washed repeatedly in the following sequence: DMF (3 x 4 L), dichloromethane (1 x 4 L) then methanol (1 x 4 L), dichloromethane (1 x 4 L) then methanol (1 x 4 L), dichloromethane (1 x 4 L) then methanol (1 x 4 L), dichloromethane (1 x 4 L) and ether (1 x 4 L). After the final washing the microKANs are dried by blowing a stream of nitrogen through the flask with intermittent agitation. After sufficient drying, the MicroKANs are sorted for the next reaction.

#### 4. Acylation:



A three neck 2 L round bottom flask is charged with the MicroKANs [784, 15 mg of resin (1.7 mmole/g loading) per MicroKAN, 25.5 micromoles/microKAN, 20.0 mmoles], and dichloromethane (800 mL). The reaction flask is fitted with an overhead stirrer. N,N-diisopropylethylamine (20.9 mL, 120 mmoles) and 4-N,N-dimethylaminopyridine (195 mg, 1.6 mmoles) are added. After approximately 15 minutes, the cyclopentanecarbonyl chloride (10.6 g, 80.0 mmoles) is added. The reaction was allowed to stir for 61 hours, the reaction flask is drained. Dichloromethane (800 mL) is added to the reaction flask. After stirring for approximately 10 minutes, the flask is drained. This is repeated. The MicroKANs from all of the acylation reactions are randomly combined into two separate large flasks and washed repeatedly in the following sequence: dichloromethane (1 x 4 L), THF (2 x 4 L), dichloromethane (1 x 4 L) then methanol (1 x 4 L), dichloromethane (1 x 4 L) then methanol (1 x 4 L), dichloromethane (1 x 4 L) then methanol (1 x 4 L), dichloromethane (1 x 4 L) and ether (1 x 4 L).

#### 5. Cleavage:

The MicroKAN is sorted into individual wells of IRORI AccuCleave 96 cleavage station. The well is charged with dichloromethane (600 mL) and then with a TFA: dichloromethane mixture (1:1, 600 mL). After agitating for approximately forty minutes, the reaction well is drained into 2 mL microtube in an 96-well format. The reaction well is again charged with dichloromethane (600 mL). After manual agitation, this too is drained into the 2 mL microtube in an 96-well format. The cleavage cocktail is

removed *in vacuo* using a Savant Speedvac. The concentrated products from the cleavage mother plates are reconstituted with THF and transferred into two daughter plates utilizing a Packard MultiProbe liquid handler. The daughter plates are concentrated *in vacuo* utilizing a GenieVac.

**Analytical:** MS: *m/z* 493 (*M*<sup>+</sup>).

5

The methods described above are used to prepare the following compounds of this invention.

5-[2-(4-(2-quinolinylmethoxy)phenoxy)methyl]benzyl]tetrazole (M.P. 108-111°C)

CALC: C, 59.87; H, 5.96; N, 13.96

10 FOUND: C, 59.67, 60.01; H, 5.62, 5.63; N, 13.73, 13.77

5-[4-Methoxy-3-(3-(2-quinolinylmethoxy)phenoxy)methyl]phenyl]tetrazole (M.P. 184-87°C)

CALC: C, 67.63; H, 4.88; N, 15.78

FOUND: C, 67.18; H, 5.13; N, 15.40

5-[3-(4-(2-quinolinylmethoxy)phenoxy)methyl]phenyl]tetrazole (M.P. 176-177°C)

CALC: C, 69.63; H, 4.75; N, 16.92

FOUND: C, 69.58, 69.64; H, 5.00, 4.98; N, 16.66, 16.63

5-[3-Methoxy-4-(4-(2-quinolinylmethoxy)benzyloxy)phenyl]tetrazole (M.P. 195-97°C)

CALC: C, 67.63; H, 4.88; N, 15.77

FOUND: C, 67.27; H, 4.89; N, 15.41

5-[4-(3-(2-quinolinylmethoxy)phenoxy)methyl]-3-methoxyphenyl]tetrazole (M.P. 189-91°C)

25 CALC: C, 66.95; H, 4.95; N, 15.61

FOUND: C, 66.48; H, 5.14; N, 14.93

5-[3-(4-(2-quinolinylmethoxy)phenoxy)methyl]benzyl]tetrazole (M.P. 139-44°C)

CALC: C, 70.53; H, 5.03; N, 16.45

30 FOUND: C, 70.33, 70.54; H, 5.25, 5.36; N, 16.38, 16.41

5-[4-(4-(2-quinolinylmethoxy)phenoxy)methyl]benzyl]tetrazole (M.P. 167-71°C)

CALC: C, 67.33; H, 5.31; N, 15.70

FOUND: C, 67.54, 67.67; H, 5.33, 5.33; N, 15.48, 15.52

35

5-[4-Methoxy-3-(4-(2-quinolinylmethoxy)phenylmethoxy)phenyl]tetrazole (M.P. 210-13°C)

CALC: C, 68.33; H, 4.82; N, 4.90

FOUND: C, 68.32; H, 4.90; N, 14.79

4-[3-(2-Quinolinylmethoxy)phenoxyethyl]phenoxyacetic acid

5 (M.P. 164 (dec))

CALC: C, 69.27; H, 5.35; N, 3.23

FOUND: C, 69.53, 69.65; H, 5.11, 5.05; N, 3.21, 3.12

5-[2-(4-(2-Quinolinylmethoxy)phenoxyethyl)phenoxyethyl]tetrazole (M.P. 183-85°C)

10 CALC: C, 65.63; H, 5.08; N, 15.31

FOUND: C, 65.77, 65.52; H, 4.99, 5.03; N, 14.92, 15.03

4-[4-(2-Quinolinylmethoxy)phenoxyethyl]phenoxyacetic acid

(176°C (dec))

15 CALC: C, 71.50; H, 5.16; N, 3.34

FOUND: C, 71.10, 71.17; H, 5.27, 5.33; N, 3.37, 3.34

4-[3-(2-Quinolinylmethoxy)phenoxyethyl]phenylacetic acid

(M.P. 158-60°C)

20 CALC: C, 75.17; H, 5.30; N, 3.51

FOUND: C, 74.89; H, 5.36; N, 3.37

2-[3-(3-(2-Quinolinylmethoxy)phenoxyethyl)phenoxy]pentanoic acid (M.P. 133-35°C)

CALC: C, 73.51; H, 5.95; N, 3.06

25 FOUND: C, 73.35, 73.60; H, 5.95, 5.98; N, 3.08, 3.05

2-[3-(2-Quinolinylmethoxy)phenoxyethyl]phenoxyacetic acid (M.P. 169-172°C)

CALC: C, 72.28; H, 5.10; N, 3.37

FOUND: C, 69.34, 69.69; H, 5.10, 5.13; N, 3.00, 3.08

30 CALC: C, 69.27; H, 5.35; N, 3.23 (as Hydrate)

2-[4-(2-Quinolinylmethoxy)phenoxyethyl]cinnamic acid (M.P. 175-178°C)

CALC: C, 75.90; H, 5.14; N, 3.40

FOUND: C, 73.92; H, 5.20; N, 3.01

35 CALC: C, 74.27; H, 5.27; N, 3.33 (as Hydrate)

6-Acetyl-2-propyl-3-[3-(2-quinolinylmethoxy)-benzyloxy]phenoxyacetic acid (M.P. 153-58°C)

CALC: C, 72.13; H, 5.85; N, 2.90

FOUND: C, 71.68, 72.08; H, 5.88, 5.83; N, 2.65, 2.70

5 2-[2-(4-(7-Chloroquinolin-2-ylmethoxy)-phenoxyethyl)phenoxy]propionic acid (M.P. 169-173°C)

CALC: C, 67.32; H, 4.78; N, 3.02; Cl, 7.64

FOUND: C, 65.18; H, 4.90; N, 2.84; Cl, 8.33

CALC: C, 65.41; H, 4.96; N, 2.93; Cl, 7.42 (as HYDRATE)

10 2-[4-(2-Quinolinylmethoxy)phenoxyethyl]phenylacetic acid (M.P. 181-83°C)

CALC: C, 75.17; H, 5.30; N, 3.51

FOUND: C, 75.12, 74.96; H, 5.50, 5.49; N, 3.16, 3.16

3-[3-(2-Quinolinylmethoxy)phenoxyethyl]phenoxyacetic acid (M.P. 146-51°C)

CALC: C, 72.28; H, 5.10; N, 3.37

FOUND: C, 71.82, 71.80; H, 5.24, 5.23; N, 2.98, 3.00

CALC: C, 71.50; H, 5.16; N, 3.34 (as HYDRATE)

2-[4-(2-Quinolinylmethoxy)phenoxyethyl]phenoxyacetic acid (M.P. 153-57°C)

CALC: C, 72.28; H, 5.10; N, 3.37

FOUND: C, 72.30, 71.72; H, 5.39, 5.30; N, 2.94, 2.89

5-[2-(4-(7-Chloroquinolin-2-ylmethoxy)-phenoxyethyl)benzyl]tetrazole (M.P. 159-63°C)

CALC: C, 65.57; H, 4.40; N, 15.29

25 FOUND: C, 64.16; H, 4.72; N, 14.98

CALC: C, 64.30; H, 4.53; N, 14.99 (as HYDRATE)

2-Carbomethoxy-5-[3-(2-quinolinylmethoxy)-phenoxyethyl]phenoxyacetic acid (M.P. 187-89°C)

CALC: C, 68.49; H, 4.90; N, 2.95

30 FOUND: C, 66.71; H, 4.96; N, 2.70

CALC: C, 66.59; H, 5.07; N, 2.87(as HYDRATE)

2-[3-(2-Quinolinylmethoxy)phenoxyethyl]-6-methylphenoxyacetic acid (M.P. 149-53°C)

CALC: C, 72.71; H, 5.40; N, 3.26

35 FOUND: C, 71.23; H, 5.46; N, 3.08

CALC: C, 71.22; H, 5.51; N, 3.19 (as HYDRATE)



2-[3-(3-(2-Quinolinylmethoxy)phenoxy)methyl]phenoxy]glutaric acid (M.P. 129-30°C)

CALC: C, 69.00; H, 5.17; N, 2.87

FOUND: C, 58.19; H, 4.93; N, 2.23

5 CALC: C, 58.23; H, 5.17; N, 2.43 (as HYDRATE)

2-[2-(2-Quinolinylmethoxy)phenoxy)methyl]benzylmalonic acid (M.P. 164-65°C)

CALC: C, 70.89; H, 4.08; N, 3.06

FOUND: C, 70.51, 70.61; H, 5.03, 5.24; N, 3.03, 2.90

10

2-[2-(3-(2-Quinolinylmethoxy)phenoxy)methyl]phenoxy]pentanoic acid (M.P. 118-20°C)

CALC: C, 73.51; H, 5.95; N, 3.06

FOUND: C, 73.26; H, 6.07; N, 2.79

15

2-[4-(2-Quinolinylmethoxy)phenoxy)methyl]-6-methylphenoxy acetic acid (M.P. 151-53°C)

CALC: C, 72.71; H, 5.40; N, 3.26

FOUND: C, 71.41; H, 5.58; N, 3.03

CALC: C, 71.22; H, 5.51; N, 3.19 (as HYDRATE)

20

2-[2-(4-(2-Quinolinylmethoxy)phenoxy)methyl]phenoxy]pentanoic acid (M.P. 85-92°C)

CALC: C, 73.51; H, 5.95; N, 3.06

FOUND: C, 71.73, 71.79; H, 5.96, 5.91; N, 3.06, 2.83

CALC: C, 72.09; H, 6.05; N, 3.00 (as HYDRATE)

25

2-Carbomethoxy-5-[4-(2-quinolinylmethoxy)-phenoxy)methyl]phenoxyacetic acid (M.P. 149-51°C)

CALC: C, 68.49; H, 4.90; N, 2.95

FOUND: C, 68.00, 68.08; H, 4.98, 5.04; N, 2.90, 2.90

2-[2-(4-(2-Quinolinylmethoxy)phenoxy)methyl]phenoxy]propionic acid (M.P. 161-64°C)

30

CALC: C, 72.71; H, 5.40; N, 3.26

FOUND: C, 70.96, 71.10; H, 5.51, 5.58; N, 3.08, 3.10

CALC: C, 71.22; H, 5.52; N, 3.19 (as HYDRATE)

2-[2-(3-(2-Quinolinylmethoxy)phenoxy)methyl]phenoxy]glutaric acid (M.P. 83°C dec)

35

CALC: C, 68.98; H, 5.17; N, 2.87

FOUND: C, 64.10, 63.75; H, 4.89, 4.92; N, 2.64, 2.69

CALC: C, 63.74; H, 5.63; N, 2.65(as HYDRATE)

2-(3-[2-Quinolinylmethyloxy]benzyloxy)phenoxyacetic acid (M.P. 153-55°C)

CALC: C, 72.28; H, 5.10; N, 3.37

5 FOUND: C, 71.75; H, 5.14; N, 3.38

CALC: C, 71.50; H, 5.16; N, 3.34 (as HYDRATE)

2-(2-[4-(2-Quinolinylmethyloxy)phenoxy]methyl]-4-chlorophenoxy)propionic acid (M.P. 196-99°C)

CALC: C, 67.32; H, 4.78; N, 3.02

10 FOUND: C, 67.40, 67.43; H, 4.89, 4.94; N, 3.01, 3.13

2-(2-[3-(2-Quinolinylmethyloxy)phenoxy]methyl]-4-chlorophenoxy)propionic acid (M.P. 169-71°C)

CALC: C, 67.32; H, 4.78; N, 3.02

FOUND: C, 65.47; H, 5.31; N, 2.78

15 CALC: C, 65.41; H, 4.96; N, 2.93 (as HYDRATE)

2-(2-[3-(2-Quinolinylmethyloxy)phenoxy]methyl]-4-chlorophenoxy)pentanoic acid (M.P. 144-45°C)

CALC: C, 68.36; H, 5.33; N, 2.85

FOUND: C, 67.74, 67.86; H, 5.39, 5.47; N, 2.91, 2.84

20 CALC: C, 67.74; H, 5.38; N, 2.82 (as HYDRATE)

2-(2-[4-(2-Quinolinylmethyloxy)phenoxy]methyl]-4-chlorophenoxy)pentanoic acid (M.P. 155-56°C)

CALC: C, 68.36; H, 5.33; N, 2.85

FOUND: C, 65.96; H, 5.59; N, 2.66

25 CALC: C, 65.95; H, 5.53; N, 2.75 (as HYDRATE)

2-(2-[4-(2-Quinolinylmethyloxy)phenoxy]methyl]-4-chlorophenoxy)pentanoic acid (M.P. 155-56°C)

CALC: C, 68.36; H, 5.33; N, 2.85

FOUND: C, 66.15; H, 5.58; N, 2.68

30 CALC: C, 65.95; H, 5.53; N, 2.75 (as HYDRATE)

2-(2-[4-(2-Quinolinylmethyloxy)phenoxy]methyl]-6-chlorophenoxy)pentanoic acid (M.P. 161-62°C)

CALC: C, 68.36; H, 5.33; N, 2.85

FOUND: C, 68.15; H, 5.36; N, 2.72

35

2-(2-[3-(2-Quinolinylmethyloxy)phenoxy]methyl]-6-chlorophenoxy)pentanoic acid (M.P. 169-70°C)

CALC: C, 68.36; H, 5.33; N, 2.85

FOUND: C, 68.10; H, 5.39; N, 2.72

2-(2-[3-(2-Quinolinylmethoxy)phenoxyethyl]-6-chlorophenoxy)-4-methylpentanoic acid (M.P.

5 164-66°C)

CALC: C, 68.84; H, 5.58; N, 2.77

FOUND: C, 68.84; H, 5.70; N, 2.69

2-(2-[4-(2-Quinolinylmethoxy)phenoxyethyl]-6-chlorophenoxy)-4-methylpentanoic acid (M.P.

10 167-69°C)

CALC: C, 68.84; H, 5.58; N, 2.77

FOUND: C, 68.78; H, 5.67; N, 2.68

5-[3-(3-(2-quinolinylmethoxy)benzyloxy)-4-methoxyphenyl]tetrazole (M.P. 204-07°C)

CALC: C, 67.63; H, 4.88; N, 15.78

FOUND: C, 67.11; H, 5.15; N, 15.86

N-[3-Methoxy-4-(3-(2-quinolinylmethoxy)benzyloxy)benzoyl]benzene sulfonamide hydrochloride  
(M.P. dec.88)

CALC: C, 62.99; H, 4.60; N, 4.74

FOUND: C, 63.88; H, 5.13; N, 4.80

5-Carboxy-2-(3-(2-quinolinylmethoxy)phenoxyethyl)phenoxy acetic acid (M.P. 226-28°C)

CALC: C, 61.90; H, 5.18; N, 2.77

25 FOUND: C, 61.62; H, 5.11; N, 2.67

5-[3-Methoxy-4-(3-(2-quinolinylmethoxy)benzyloxy)phenyl]tetrazole (M.P. 204-05°C)

CALC: C, 67.67; H, 5.14; N, 15.87















FOUND: C, 67.63; H, 4.88; N, 15.78
















30


5-(4-(3-(2-Quinolinylmethoxy)benzyloxy)phenyl)tetrazole (M.P. 233-36°C)













CALC: C, 69.58; H, 4.73; N, 16.91















FOUND: C, 69.59; H, 4.89; N, 16.91



















;




;




;




;




;




;




;




;




;




;




;

















;













;





;








of, CHN calc.  
C<sub>30</sub>H<sub>29</sub>NO<sub>7</sub>+0.5  
H<sub>2</sub>O: C 68.69, H  
5.76, N 2.67; found  
C 68.68, H 5.71, N  
2.86



CHN calc.  
C<sub>29</sub>H<sub>29</sub>NO +  
0.75 H<sub>2</sub>O:  
C 74.26, H 5.55,  
N 2.99;  
found C 73.97,  
H 6.31, N 2.89





Using a combination of the above Examples, various compounds may be made within the scope of this invention.

Compounds according to the invention exhibit marked pharmacological activities according to tests described in the literature which tests results are believed to correlate to pharmacological activity in humans and other mammals. The following pharmacological test results are typical characteristics of compounds of the present invention.

The compounds of the present invention have potent activity as PPAR ligand receptor binders and possess anti-diabetic, anti-lipidemic, anti-hypertensive, and anti-arteriosclerotic activity and are also anticipated to be effective in the treatment of diabetes, obesity and other related diseases.

#### hPPAR $\alpha$ Binding Assay

The activity of the compounds of the invention as PPAR $\alpha$  modulators may be examined in several relevant in vitro and in vivo preclinical assays, for example benchmarking with a known PPAR $\alpha$  modulator, for example, [ $^3$ H]-GW2331(2-(4-[2-(3-[2,4-Difluorophenyl]-1-heptylureido)-ethyl]phenoxy)-2-methylbutyric acid). (S. Kliewer, et al. Proc. Natl. Acad. Sci. USA 94 (1997).

#### Human peroxime proliferator-activated receptor a ligand binding domain(hPPAR $\alpha$ -LBD):

A binding assay for PPAR $\alpha$  could be carried out by the following procedure: cDNAs encoding the putative ligand binding domain of human PPAR $\alpha$  (amino acids 167-468) ( Sher,T., Yi, H.-F., McBride, O. W.& Gonzalez, F. J. (1993) *Biochemistry* 32, 5598-5604) are amplified by PCR (Polymerase

Chain Reaction) and inserted in frame into the BamHI site of pGEX-2T plasmid (Pharmacia). The soluble fraction of GST-hPPAR $\alpha$  fusion proteins or glutathione S-transferase (GST) alone are overexpressed in *E. coli* BL21(DE3)pLysS cells and purified from bacteria extracts as described in (S. Klierer, et al. Proc. Natl. Acad. Sci. USA 94 (1997), 4318-4323).

Gel-Filtration Assays: 30 ml of 90 nM GST-hPPAR $\alpha$ -LBD is mixed with 20 ml of 50 nM  $^3$ H-GW2331 with or without 5 ml of 10 mM test compounds in the binding buffer containing 10 mM Tris, 50 mM KCl, 0.05% Tween 20 and 10 mM DTT. The reaction mixtures are incubated in 96-well plates for 2h at room temperature. 50 ml of the reaction mixtures are then loaded on a 96-well gel filtration block (following manufacture instructions)(EdgeBioSystems). The block placed on top of a clean 96-well plate is centrifuged at 1,500 rpm for 2 min. The block is discarded. 100 ml of Scintillation fluid is added to each well of the 96-well plate. After overnight equilibration, the plate is counted in the Microbeta counter (Wallac.).

Homogenous Scintillation Proximity Binding Assay. For the Scarchard analysis, glutathione coated SPA beads (1.5 mg/ml )(Amersham) are mixed with GST-hPPAR $\alpha$ -LBD (10 mg/ml) in the binding buffer. The resulting slurry is incubated at room temperature with agitation for 15 min. 20 ml of the slurry is then added in 30 ml of binding buffer containing various amount  $^3$ H-GW2331(10–500 nM). Nonspecific binding is determined in the present of 100 mM of GW2331. For the competition binding assay, 20 ml of the slurry is then added in 30 ml of the binding buffer containing 75 nM of  $^3$ H-GW2331 and 0.03–20 mM of the test compounds. For the control experiments, the glutathione coated SPA beads (1.5 mg/ml) are coated with GST proteins (10 mg/ml). 20 ml of the slurry are mixed with 30 ml of 75 nM of  $^3$ H-GW2331 with or without 10 mM of GW2331. The above experiments are all performed in a 96-well plates. The sealed plates with the reaction mixtures are allowed to equilibrate for 2 h and counted in the Microbeta counter (Wallac.).

#### hPPAR $\gamma$ Binding Assay

The activity of the compounds of the invention as PPAR $\gamma$  modulators may be examined in several relevant in vitro and in vivo preclinical assays, for example benchmarking with a known PPAR $\gamma$  modulator, for example, [ $^3$ H]-BRL 49853 (Lehman L.J. et al, J. Biol. Chem. 270, 12953-12956; Lehman L.J. et al, J. Biol. Chem. 272, 3406-3410 (1997), and Nichols, J. S.; et al Analytical Biochemistry 257, 112-119(1998)).


#### Human peroxime proliferator-activated receptor a ligand binding domain(hPPAR $\gamma$ -LBD).

A binding assay for PPAR $\gamma$  could be carried out by the following procedure: cDNAs encoding the putative ligand binding domain of human PPAR $\gamma$  (amino acids 176-477) (Green, M.E. et al. Gene expression 281-299(1995)) are amplified by PCR (polymerase chain reaction) and inserted in frame into the BamHI site of pGEX-2T plasmid (Pharmacia). The soluble fraction of GST-hPPAR $\gamma$  fusion proteins or glutathione S-transferase (GST) alone are overexpressed in *E. coli* BL21(DE3)pLysS cells and purified from bacteria extracts.

**Binding Assay:** The fusion proteins, GST-PPAR $\gamma$  -LBD in PBS (5 mg/100ml/well) are incubated in the glutathione coated 96 well plates for 4 hours. Unbound proteins are then discarded and the plates are washed two times with the wash buffer (10 mM Tris, 50 mM KCl and 0.05% Tween-20). 100 ml of reaction mixtures containing 60 nM of  $^3\text{H}$ -BRL-49853 and 10 mM of the testing compounds (10 ml of 0.1mM compounds from each well of the child plates) in the binding buffer (10mM Tris, 50mM KCl and 10mM DTT) are then added and incubated at room temperature for 2.5h. The reaction mixtures are discarded and the plates are washed two times with the wash buffer. 100ml of scintillation fluid is added to each well and plates are counted on  $\beta$ -counter.

#### hPPAR $\delta$ Binding Assay

The activity of the compounds of the invention as PPAR $\delta$  modulators may be examined in several relevant in vitro and in vivo preclinical assays (See references WO 97/28149; Brown P. et al Chemistry & Biology, 4, 909-18, (1997)), for example benchmarking with a known PPAR $\delta$  modulator, for example [ $^3\text{H}$ ] GW2433 or [ $^3\text{H}$ ] Compound X



Compound X

The hPPAR $\delta$  binding assay comprises the steps of:

- preparing multiple test samples by incubating separate aliquots of the receptor hPPAR $\delta$  with a test compound in TEGM containing 5-10% COS-1 cell cytoplasmic lysate and 2.5 nM labeled ( $^3\text{H}$ )Compound X, 17 Ci/mmol) for a minimum of 12 hours, and preferably for about 16 hours, at 4°C, wherein the concentration of the test compound in each test sample is different, and preparing a control sample by incubating a further separate aliquot of the receptor hPPAR $\delta$  under the same conditions but without the test compound; then

- (b) removing unbound ligand by adding dextran/gelatin-coated charcoal to each sample while maintaining the samples at 4°C and allowing at least 10 minutes to pass, then
- (c) subjecting each of the test samples and control sample from step (b) to centrifugation at 4°C until the charcoal is pelleted; then
- (d) counting a portion of the supernatant fraction of each of the test samples and the control sample from step (c) in a liquid scintillation counter and analyzing the results to determine the IC<sub>50</sub> of the test compound.

In the hPPAR $\delta$  binding assay, preferably at least four test samples of varying concentrations of a single test compound are prepared in order to determine the IC<sub>50</sub>.

The compounds useful according to the invention can be administered to a patient in a variety of forms adapted to the chosen route of administration, i.e., orally, or parenterally. Parenteral administration in this respect includes administration by the following routes: intravenous, intramuscular, subcutaneous, intraocular, intrasynovial, transepthially including transdermal, ophthalmic, sublingual and buccal; topically including ophthalmic, dermal, ocular, rectal and nasal inhalation via insufflation and aerosol and rectal systemic.

The active compound may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed in hard or soft shell gelatin capsules, or it may be compressed into tablets, or it may be incorporated directly with the food of the diet. For oral therapeutic administration, the active compound may be incorporated with excipient and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.1% of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be from about 2% to about 6% of the weight of the unit. The amount of active compound in such therapeutically useful compositions is such that a suitable dosage will be obtained. Preferred compositions or preparations according to the present invention are prepared so that an oral dosage unit form contains between about 50 and 300 mg of active compound.

The tablets, troches, pills, capsules and the like may also contain the following: A binder such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be added or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present

as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. Of course, any material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compound may be incorporated into sustained-release preparations and formulations.

The active compound may also be administered parenterally or intraperitoneally. Solutions of the active compound as a free base or pharmacologically acceptable salt can be prepared in water suitably mixed with a surfactant such as hydroxypropyl-cellulose. Dispersion can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It may be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compound in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredient into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying technique which yield a powder of the active ingredient plus any additional desired ingredient from previously sterile-filtered solution thereof.



The therapeutic compounds useful according to this invention may be administered to a patient alone or in combination with pharmaceutically acceptable carriers, as noted above, the proportion of which is determined by the solubility and chemical nature of the compound, chosen route of administration and standard pharmaceutical practice.


The physician will determine the dosage of the present therapeutic agents which will be most suitable for prophylaxis or treatment and it will vary with the form of administration and the particular compound chosen, and also, it will vary with the particular patient under treatment. He will generally wish to initiate treatment with small dosages by small increments until the optimum effect under the circumstances is reached. The therapeutic dosage will generally be from 0.1 to 100 mM/day or from about 0.1mg to about 50 mg/kg of body weight per day, or 10mg to about 50 mg/kg of body weight per day, or more preferably 30mg to about 50 mg/kg of body weight per day, and higher, although it may be administered in several different dosage units. Higher dosages are required for oral administration.

The compounds useful according to the invention may be administered as frequently as necessary in order to obtain the desired therapeutic effect. Some patients may respond rapidly to a higher or lower dose and may find much weaker maintenance doses adequate. For other patients, it may be necessary to have long-term treatments at the rate of 1 to 4 doses per day, in accordance with the physiological requirements of each particular patient. Generally, the active product may be administered orally 1 to 4 times per day. It goes without saying that, for other patients, it will be necessary to prescribe not more than one or two doses per day.

One skilled in the art will readily appreciate that the present invention is well adapted to carry out the objects of the invention and obtain the ends and advantages mentioned, as well as those inherent therein. The compounds, compositions and methods described herein are presented as representative of the preferred embodiments, or intended to be exemplary and not intended as limitations on the scope of the present invention.

What is claimed is:

1. A compound of formula I




5 wherein:



, and are independently aryl, fused arylcycloalkenyl, fused arylcycloalkyl, fused arylheterocyclenyl, fused arylheterocyclyl, heteroaryl, fused heteroaryl cycloalkenyl, fused heteroaryl cycloalkyl, fused heteroaryl heterocyclenyl, or fused heteroaryl heterocyclyl;

A is -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sub>13</sub>-, -C(O)-, -N(R<sub>14</sub>)C(O)-, -C(O)N(R<sub>15</sub>)-, -N(R<sub>14</sub>)C(O)N(R<sub>15</sub>)-, -C(R<sub>14</sub>)=N-, a chemical bond,



B is -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sub>17</sub>-, a chemical bond, ethynylene, -C(O)-, -N(R<sub>18</sub>)C(O)-, or -C(O)NR<sub>18</sub>-;

D is -O-, -S-, -NR<sub>19</sub>-, a chemical bond, ethynylene, -N(R<sub>20</sub>)C(O)-, -C(O)-, or -C(O)N(R<sub>20</sub>)-;

E is a chemical bond or an ethylene group;

a is 0-4;

b is 0-4;

c is 0-4;

d is 0-5;

e is 0-4;

$f$  is 0-6;

g is 1-4;

h is 1-4;

R<sub>1</sub>, R<sub>3</sub>, R<sub>5</sub>, R<sub>7</sub>, R<sub>9</sub>, and R<sub>11</sub>, are independently hydrogen, halogen, alkyl, carboxyl, alkoxycarbonyl or aralkyl;

5 R<sub>2</sub>, R<sub>4</sub>, R<sub>6</sub>, R<sub>8</sub>, R<sub>10</sub> and R<sub>12</sub>, are independently  $-(CH_2)_q-X$ ;

q is 0-3;

X is hydrogen, halogen, alkyl, alkenyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, aralkyl, heteroaralkyl, hydroxy, alkoxy, aralkoxy, heteroaralkoxy, carboxyl, alkoxycarbonyl, tetrazolyl, acyl, acylHNSO<sub>2</sub><sup>-</sup>, -SR<sub>23</sub>, Y<sup>1</sup>Y<sup>2</sup>N- or Y<sup>3</sup>Y<sup>4</sup>NCO-;

10 Y<sup>1</sup> and Y<sup>2</sup> are independently hydrogen, alkyl, aryl, aralkyl or heteroaralkyl, or one of Y<sup>1</sup> and Y<sup>2</sup> is hydrogen or alkyl and the other of Y<sup>1</sup> and Y<sup>2</sup> is acyl or aroyl;

Y<sup>3</sup> and Y<sup>4</sup> are independently hydrogen, alkyl, aryl, aralkyl or heteroaralkyl;

Z is R<sub>21</sub>O<sub>2</sub>C-, R<sub>21</sub>OC-, cyclo-imide, -CN, R<sub>21</sub>O<sub>2</sub>SHNCO-, R<sub>21</sub>O<sub>2</sub>SHN-, (R<sub>21</sub>)<sub>2</sub>NCO-, R<sub>21</sub>O- 2,4-thiazolidinedionyl, or tetrazolyl; and

15 R<sub>19</sub> and R<sub>21</sub> are independently hydrogen, alkyl, aryl, cycloalkyl, or aralkyl;

R<sub>13</sub>, R<sub>17</sub>, R<sub>19</sub> and R<sub>23</sub> are independently R<sub>22</sub>OC-, R<sub>22</sub>NHOC-, hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, heteroaralkyl, or aralkyl;

R<sub>14</sub>, R<sub>15</sub>, R<sub>16</sub>, R<sub>18</sub> and R<sub>20</sub> are independently hydrogen, alkyl, aralkyl, carbonyl, or alkoxycarbonyl;

20 or R<sub>14</sub>, and R<sub>15</sub> taken together with the carbon and nitrogen atoms through which they are linked form a 5 or 6-membered azaheterocyclyl group; or

when a is 2-4, then vicinal R<sub>1</sub> radicals taken together with the carbon atoms to which the R<sub>1</sub> radicals are linked form an ethylene group; or

when b is 2-4, then vicinal R<sub>3</sub> radicals taken together with the carbon atoms to which the R<sub>3</sub> radicals are linked form an ethylene group; or


25 when c is 2-4, then vicinal R<sub>5</sub> radicals taken together with the carbon atoms to which the R<sub>5</sub> radicals are linked form an ethylene group; or

when d is 2-5, then vicinal R<sub>7</sub> radicals taken together with the carbon atoms to which the R<sub>7</sub> radicals are linked form an ethylene group; or


30 when e is 2-4, then vicinal R<sub>9</sub> radicals taken together with the carbon atoms to which the R<sub>9</sub> radicals are linked form an ethylene group; or

when f is 2-6, then vicinal  $R_{11}$  radicals taken together with the carbon atoms to which the  $R_{11}$  radicals are linked form an ethylene group; and


$R_{22}$  is hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, heteroaralkyl, or aralkyl; or a pharmaceutically acceptable salt thereof, an N-oxide thereof, a hydrate thereof or a solvate thereof.

2. A compound according to claim 1 wherein  is optionally substituted aryl,

optionally substituted azaheteroaryl, or optionally substituted fused arylheterocyclenyl; is optionally substituted aryl, optionally substituted heteroaryl, or optionally substituted fused

arylheterocyclenyl; and  is optionally substituted aryl, optionally substituted heteroaryl, optionally substituted fused arylheterocyclalkyl or optionally substituted fused arylheterocyclenyl.

3. A compound according to claim 1 wherein a = 1 or 2;  $R_1$  and  $R_2$  is hydrogen; A is a chemical bond; and b = 0.

4. A compound according to claim 1 wherein a = 0; A is ;  $R_{15}$  and  $R_{16}$  are hydrogen; g is 1, 2, or 3; and b = 0.

5. A compound according to claim 1 wherein a = 0; A is  $-NR_{13}-$ , b = 1,  $R_3$  and  $R_4$  are hydrogen.

6. A compound according to claim 1 wherein a = 2; vicinal  $R_1$  radicals taken together with the carbon atoms to which the  $R_1$  radicals are linked form an ethylene group;  $R_2$  is hydrogen; A is a chemical bond; and b=0.

7. A compound according to claim 1 wherein a = 1, 2 or 3;  $R_1$  and  $R_2$  are hydrogen; A is  $-O-$ ; and b = 0.

8. A compound according to claim 1 wherein a = 1;  $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$  are hydrogen; A is  $-O-$ ; and b = 1.

9. A compound according to claim 1 wherein c = 1 or 2;  $R_5$  and  $R_6$  are hydrogen or alkyl; B is a chemical bond; and d = 0.

10. A compound according to claim 1 wherein  $c = 2$ ; vicinal  $R_5$  radicals taken together with the carbon atoms to which the  $R_5$  radicals are linked form an ethylene group;  $R_6$  is hydrogen; B is a chemical bond; and  $d=0$ .

11. A compound according to claim 1 wherein  $c = 0$  or  $1$ ;  $R_5$  and  $R_6$  are hydrogen; B is  $-O-$ ; and  $d = 0$  or  $1$ .

12. A compound according to claim 1 wherein  $c = 0$ ; B is  $-C(O)-$  or  $-S(O)_2-$ ;  $d = 1$  and  $R_7$  and  $R_8$  are independently hydrogen or alkyl.

13. A compound according to claim 1 wherein  $e = 0$ ;  $f = 0$ ; D and E is a chemical bond; Z is  $R_{21}O_2SHNCO-$ , and  $R_{21}$  is phenyl.

14. A compound according to claim 1 wherein  $e = 0$ ;  $f = 0$  or  $1$ ; D and E is a chemical bond; Z is tetrazolyl,  $NH_2CO-$  or  $-CO_2R_{21}$ ; and  $R_{21}$  is hydrogen or lower alkyl.

15. A compound according to claim 1 wherein  $e = 0$ ;  $f = 0$  or  $1$ ; D is  $-O-$  or a chemical bond; E is a chemical bond; and Z is tetrazolyl,  $NH_2CO-$  or  $-CO_2R_{21}$ ; and  $R_{21}$  is hydrogen or lower alkyl.


16. A compound according to claim 1 wherein  $e = 0$ ;  $f = 1$ ; D is  $-O-$  or a chemical bond; E is a chemical bond;  $R_{11}$  and  $R_{12}$  are hydrogen or alkyl; and Z is tetrazolyl,  $NH_2CO-$  or  $-CO_2R_{21}$ ; and  $R_{21}$  is hydrogen or lower alkyl.

17. A compound according to claim 1 wherein  $e = 2$ , then vicinal  $R_9$  radicals taken together with the carbon atoms to which the  $R_9$  radicals are linked form an ethylene group;  $f = 0$ ; D and E is a chemical bond; and Z is  $-CO_2R_{21}$ ; and  $R_{21}$  is hydrogen.

18. A compound according to claim 1 wherein  $e = 0$ ;  $f = 3$ ; D is  $-O-$ ; E is a chemical bond;  $R_{11}$  and  $R_{12}$  are hydrogen or alkyl, or at least one of  $R_{11}$  is carboxyl or alkoxy carbonyl; Z is tetrazolyl, or  $-CO_2R_{21}$ ; and  $R_{21}$  is hydrogen or lower alkyl.


19. A compound according to claim 1 wherein  $e = 0$ ;  $f = 1, 2$ , or  $3$ ; D is  $-C(O)-$ ; E is a chemical bond;  $R_{11}$  and  $R_{12}$  are hydrogen or alkyl; Z is tetrazolyl or  $-CO_2R_{21}$ ; and  $R_{21}$  is hydrogen or lower alkyl.




20. A compound according to claim 1 wherein  is an optionally substituted quinolinyl, quinoxalinyl, quinazolinyl, isoquinolinyl, *N*-alkyl-quinolin-4-onyl, quinazolin-4-onyl, benzoxazolyl, benzimidazolyl, benzothiazolyl, benzofuranyl, benzothiophenyl, indolinyl, oxazolyl, thiazolyl, oxadiazolyl, isoxazolyl, imidazolyl, pyrazol-yl, thiadiazolyl, triazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, phenyl, or naphthalenyl group, wherein the substituent is a

ring system substituent as defined herein, more preferably a substituent selected from the group consisting of phenyl, substituted-phenyl, thienyl, substituted thienyl, cycloalkyl, lower alkyl, branched alkyl, fluoro, chloro, alkoxy, aralkyloxy, trifluoromethyl and trifluoromethyloxy.



21. A compound according to claim 1 wherein  is unsubstituted quinolin-2-yl, 3-substituted quinolin-2-yl, 4-substituted quinolin-2-yl, 6-substituted quinolin-2-yl or 7 substituted quinolin-2-yl; an unsubstituted quinoxalin-2-yl, 3-substituted quinoxalin-2-yl, 6-substituted quinoxalin-2-yl or 3,6-disubstituted quinoxalin-2-yl; unsubstituted quinazolin-2-yl, 4-substituted quinazolin-2-yl or 6-substituted quinazolin-2-yl; unsubstituted isoquinolin-3-yl, 6-substituted isoquinolin-3-yl or 7-substituted isoquinolin-3-yl; 3-substituted-quinazolin-4-on-2-yl; *N*-substituted quinolin-4-on-2-yl; 2-substituted-oxazol-4-yl or 2,5 disubstituted-oxazol-4-yl; 4-substituted oxazol-2-yl or 4,5-disubstituted-oxazol-2-yl; 2-substituted thiazol-4-yl or 2,5-disubstituted thiazol-4-yl; 4-substituted thiazol-2-yl or 4,5-disubstituted-thiazol-2-yl; 5-substituted-[1,2,4]oxadiazol-3-yl; 3-substituted-[1,2,4] oxadiazol-5-yl; 5-substituted-imidazol-3-yl or 3,5-disubstituted-imidazol-2-yl; 2-substituted-imidazol-5-yl or 2,3-disubstituted-imidazol-5-yl; 3-substituted-isoxazol-5-yl; 5-substituted-isoxazol-3-yl; 5-substituted-[1,2,4] thiadiazol-3-yl; 3-substituted-[1,2,4]-thiadiazol-5-yl; 2-substituted-[1,3,4]-thiadiazol-5-yl; 2-substituted-[1,3,4]-oxadiazol-5-yl; 1-substituted-pyrazol-3-yl; 3-substituted-pyrazol-5-yl; 3-substituted-[1,2,4]-triazol-5-yl; 1-substituted-[1,2,4]-triazol-3-yl; 3-substituted pyridin-2-yl, 5-substituted pyridin-2-yl, 6-substituted pyridin-2-yl or 3,5-disubstituted pyridin-2-yl; 3-substituted pyrazin-2-yl, 5-substituted pyrazin-2-yl, 6-substituted pyrazin-2-yl or 3,5 disubstituted-pyrazin-2-yl; 5-substituted pyrimidin-2-yl or 6-substituted-pyrimidin-2-yl; 6-substituted-pyridazin-3-yl or 4,6-disubstituted-pyridazin-3-yl; unsubstituted naphthalen-2-yl, 3-substituted naphthalen-2-yl, 4-substituted naphthalen-2-yl, 6-substituted naphthalen-2-yl or 7 substituted naphthalen-2-yl; 2-substituted phenyl, 4-substituted phenyl or 2,4-disubstituted phenyl; unsubstituted -benzothiazol-2-yl or 5-substituted-benzothiazol-2-yl; unsubstituted benzoxazol-2yl or 5-substituted-benzoxazol-2yl; unsubstituted -benzimidazol-2-yl or 5-substituted-benzimidazol-2-yl; unsubstituted -thiophen-2yl, 3-substituted -thiophen-2yl, 6-substituted -thiophen-2yl or 3,6-disubstituted-thiophen-2yl; unsubstituted -benzofuran-2-y, 3-substituted-benzofuran-2-yl, 6-substituted-benzofuran-2-yl or 3,6-disubstituted-benzofuran-2-yl; 3-substituted-benzofuran-6-yl or 3,7-disubstituted-benzofuran-6-yl, wherein the substituent is a ring system substituent.



22 A compound according to claim 21 wherein  is substituted by a substituent selected from the group consisting of phenyl, substituted-phenyl, thienyl, substituted thienyl, cycloalkyl, lower alkyl, branched alkyl, fluoro, chloro, alkoxy, aralkyloxy, trifluoromethyl and trifluoromethyloxy.


23. A compound according to claim 1 wherein  $R_1$  and  $R_2$  are hydrogen;  $a = 1$ ; A is  $-O-$ ; and  $b = 0$ .

24. A compound according to claim 1 wherein  $R_1$  and  $R_2$  are hydrogen;  $a = 2$ ; A is  $-O-$ ; and  $b = 0$ .


25. A compound according to claim 1 wherein  $a = 0$ ; A is  $-O-$  or  $-NR_{13}-$ ;  $R_{13}$  is hydrogen or alkyl;  $R_3$  and  $R_4$  are both independently hydrogen; and  $b = 1$ .

26. A compound according to claim 1 wherein  $a = 0$ ; A is  $-O-$  or  $-NR_{13}-$ ;  $R_{13}$  is hydrogen or



alkyl;  $R_3$  and  $R_4$  are both independently hydrogen;  $b = 1$ ; and  is 3-substituted quinolin-2-yl, 4-substituted quinolin-2-yl, 6-substituted quinolin-2-yl, 7 substituted quinolin-2-yl, unsubstituted quinoxalin-2-yl, 3-substituted quinoxalin-2-yl, 6-substituted quinoxalin-2-yl, 3,6-disubstituted quinoxalin-2-yl, unsubstituted quinazolin-2-yl, 4-substituted quinazolin-2-yl, 6-substituted quinazolin-2-yl, unsubstituted isoquinolin-3-yl, 6-substituted isoquinolin-3-yl, 7-substituted isoquinolin-3-yl, 4-substituted oxazol-2-yl, 4,5-disubstituted-oxazol-2-yl, 4-substituted-thiazol-2-yl, 4,5-disubstituted-thiazol-2-yl, 5-substituted -imidazol-2-yl, 3,5-disubstituted-imidazol-2-yl, 1-substituted-pyrazol-3-yl, 3-substituted-pyrazol-5-yl, 3-substituted pyridin-2-yl, 5-substituted pyridin-2-yl, 6-substituted pyridin-2-yl or 3,5-disubstituted pyridin-2-yl, 3-substituted pyrazin-2-yl, 5-substituted pyrazin-2-yl, 6-substituted pyrazin-2-yl, 3,5-disubstituted-pyrazin-2-yl, 5-substituted pyrimidin-2-yl, 6-substituted-pyrimidin-2-yl, 6-substituted-pyridazin-3-yl, 4,6-disubstituted-pyridazin-3-yl, unsubstituted-benzothiazol-2-yl, 5-substituted-benzothiazol-2-yl, unsubstituted-benzoxazol-2-yl, 5-substituted-benzoxazol-2-yl, unsubstituted benzimidazol-2-yl, 5-substituted-benzimidazol-2-yl, 3-substituted-benzofuran-6-yl or 3,7-disubstituted-benzofuran-6-yl.

27. A compound of formula (Ia)




wherein:



and

and are independently aryl, fused arylcycloalkenyl, fused arylcycloalkyl, fused arylheterocyclenyl, fused arylheterocyclyl, heteroaryl, fused heteroarylcycloalkenyl, fused heteroarylcycloalkyl, fused heteroarylheterocyclenyl, or fused heteroarylheterocyclyl;

A is -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sub>13</sub>-, -C(O)-, -N(R<sub>14</sub>)C(O)-, -C(O)N(R<sub>15</sub>)-, -N(R<sub>14</sub>)C(O)N(R<sub>15</sub>)-, -C(R<sub>14</sub>)=N-, a chemical bond,



10 B is -O-, -S-, -SO-, -SO<sub>2</sub>-, -NR<sub>17</sub>-, a chemical bond, ethynylene, -C(O)-, -N(R<sub>18</sub>)C(O)-, or -C(O)NR<sub>18</sub>-;

D is  $-O-$ ,  $-S-$ ,  $-NR_{19}-$ , a chemical bond, ethynylene,  $-N(R_{20})C(O)-$ ,  $-C(O)-$ , or  $-C(O)N(R_{20})-$ ;

E is a chemical bond or an ethylene group:

a is 0-4;

15    b is 0-4;

c is 0-4;



d is 0-5;

e is 0-4;

f is 0-6;

g is 1-4;

5 h is 1-4;

R<sub>1</sub>, R<sub>3</sub>, R<sub>5</sub>, R<sub>7</sub>, R<sub>9</sub>, and R<sub>11</sub>, are independently hydrogen, halogen, alkyl, carboxyl, alkoxycarbonyl or aralkyl;

R<sub>2</sub>, R<sub>4</sub>, R<sub>6</sub>, R<sub>8</sub>, R<sub>10</sub> and R<sub>12</sub>, are independently  $-(CH_2)_q-X$ ;

q is 0-3;

10 X is hydrogen, halogen, alkyl, alkenyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, aralkyl, heteroaralkyl, hydroxy, alkoxy, aralkoxy, heteroaralkoxy, carboxyl, alkoxycarbonyl, tetrazolyl, acyl, acylHNSO<sub>2</sub><sup>-</sup>, -SR<sub>23</sub>, Y<sup>1</sup>Y<sup>2</sup>N- or Y<sup>3</sup>Y<sup>4</sup>NCO-;

Y<sup>1</sup> and Y<sup>2</sup> are independently hydrogen, alkyl, aryl, aralkyl or heteroaralkyl, or one of Y<sup>1</sup> and Y<sup>2</sup> is hydrogen or alkyl and the other of Y<sup>1</sup> and Y<sup>2</sup> is aryl or aroyl;

15 Y<sup>3</sup> and Y<sup>4</sup> are independently hydrogen, alkyl, aryl, aralkyl or heteroaralkyl;

Z is R<sub>21</sub>O<sub>2</sub>C-, R<sub>21</sub>OC-, cyclo-imide, -CN, R<sub>21</sub>O<sub>2</sub>SHNCO-, R<sub>21</sub>O<sub>2</sub>SHN-, (R<sub>21</sub>)<sub>2</sub>NCO-, R<sub>21</sub>O- 2,4-thiazolidinedionyl, or tetrazolyl;

R' and R'' are ring system substituents;

R<sub>19</sub> and R<sub>21</sub> are independently hydrogen, alkyl, aryl, cycloalkyl, or aralkyl;

20 R<sub>13</sub>, R<sub>17</sub>, R<sub>19</sub> and R<sub>23</sub> are independently R<sub>22</sub>OC-, R<sub>22</sub>NHOC-, hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, heteroaralkyl, or aralkyl;

R<sub>14</sub>, R<sub>15</sub>, R<sub>16</sub>, R<sub>18</sub> and R<sub>20</sub> are independently hydrogen, alkyl, aralkyl, carbonyl, or alkoxycarbonyl;

or R<sub>14</sub>, and R<sub>15</sub> taken together with the carbon and nitrogen atoms through which they are linked

25 form a 5 or 6-membered azaheterocyclyl group; or

when a is 2-4, then vicinal R<sub>1</sub> radicals taken together with the carbon atoms to which the R<sub>1</sub> radicals are linked form an ethylene group; or

when b is 2-4, then vicinal R<sub>3</sub> radicals taken together with the carbon atoms to which the R<sub>3</sub> radicals are linked form an ethylene group; or

30 when c is 2-4, then vicinal R<sub>5</sub> radicals taken together with the carbon atoms to which the R<sub>5</sub> radicals are linked form an ethylene group; or

when d is 2-5, then vicinal  $R_7$  radicals taken together with the carbon atoms to which the  $R_7$  radicals are linked form an ethylene group; or


when e is 2-4, then vicinal  $R_9$  radicals taken together with the carbon atoms to which the  $R_9$  radicals are linked form an ethylene group; or

- 5 when f is 2-6, then vicinal  $R_{11}$  radicals taken together with the carbon atoms to which the  $R_{11}$  radicals are linked form an ethylene group; and

$R_{22}$  is hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, heteroaralkyl, or aralkyl; or a pharmaceutically acceptable salt thereof, an N-oxide thereof, a hydrate thereof or a solvate thereof.

10 28. A compound according to claim 27

wherein



are independently aryl, fused arylcycloalkenyl, fused arylcycloalkyl, fused arylheterocyclenyl, fused arylheterocyclyl, heteroaryl, fused heteroarylcyloalkenyl, fused heteroarylcyloalkyl, fused heteroarylheterocyclenyl, or fused heteroarylheterocyclyl;

c+d = 1 or 2;

B is  $-O-$ ;

$R_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$  are independently hydrogen;

e = 0;

20 f = 0;


D and E are a chemical bond;

Z is  $R_{21}O_2C-$ ,  $R_{21}OC-$ , cyclo-imide,  $-CN$ ,  $R_{21}O_2SHNCO-$ ,  $R_{21}O_2SHN-$ ,  $(R_{21})_2NCO-$ ,  $R_{21}O-$ , 2,4-thiazolidinedionyl, or tetrazolyl;

$R'$  is lower alkyl, halo, alkoxy, aryloxy or aralkyl; and

25  $R''$  is lower alkyl or halo.

29. A compound according to claim 27 wherein



are independently aryl, fused arylcycloalkenyl, fused arylcycloalkyl, fused arylheterocyclenyl, fused arylheterocyclyl, heteroaryl, fused heteroarylcyloalkenyl, fused heteroarylcyloalkyl, fused heteroarylheterocyclenyl, or fused heteroarylheterocyclyl;

$c+d = 1$  or  $2$ ;

B is  $-O-$ ;

$R_5, R_6, R_7, R_8$  are independently hydrogen;

$e = 0$ ;

5  $f = 0$ ;

D and E are a chemical bond;

Z is  $-CO_2H$ ;

$R'$  is lower alkyl, halo, alkoxy, aryloxy or aralkyl; and

$R''$  is lower alkyl or halo.

10

30. A compound according to claim 27 wherein

$a = 0-2$ ;

$b = 0-1$ ;

A is  $-O-$  or  $-NR_{13}-$ ;

15

$c+d = 1$  or  $2$ ;

B is  $-O-$ ;

$R_1, R_2, R_3, R_4, R_5, R_6, R_7$ , and  $R_8$  are independently hydrogen;

$R_{13}$  is hydrogen,  $R_{22}OC-$ , or alkyl;

$e = 0$ ;

20

$f = 0$ ;

D and E are a chemical bond;

Z is  $-CO_2H$ ;

$R'$  is lower alkyl, halo, alkoxy, aryloxy or aralkyl; and

$R''$  is lower alkyl or halo.

25

31. A compound according to claim 27 wherein

$a = 1$  or  $2$ ;

A is  $-O-$ ;

$b = 0$ ;

30

$R_1, R_2, R_7$  and  $R_8$  are independently hydrogen;



is optionally substituted phenyl;

c = 0;

B is -O-;

d = 1;

5 e = 0;

f = 0;

D and E are a chemical bond;

R' is hydrogen, halo or benzyloxy;

R'' is lower alkyl, preferably methyl;

Z is -CO<sub>2</sub>H.

32. A compound according to claim 27 wherein:

a = 1 or 2;

A is -O-;

b = 0;

R<sub>1</sub>, R<sub>2</sub>, R<sub>5</sub> and R<sub>6</sub> are independently hydrogen;



is optionally substituted phenyl;

c = 1;

B is -O-;

d = 0;

e = 0;

f = 0;

D and E are a chemical bond;

R' is hydrogen, halo or benzyloxy;

20 R'' is lower alkyl, preferably methyl;

Z is -CO<sub>2</sub>H.

33. A compound according to claim 27 wherein:

a = 1 or 2;

A is -O-;

b = 0;

R<sub>1</sub>, R<sub>2</sub>, R<sub>7</sub>, R<sub>8</sub>, R<sub>11</sub> and R<sub>12</sub> are independently hydrogen;

Ar II

is optionally substituted phenyl;

c = 0;

B is -O-;

d = 1;

e = 0;

f = 1;

D and E are a chemical bond;

R' is halo;

R'' is lower alkyl, preferably methyl;

Z is -CO<sub>2</sub>H.

34. A compound according to claim 27 wherein:

a = 1;

A is -O-;

b = 0;

c = 0-1;

B is -O-;

d = 0 or 1, wherein c+d = 1 or 2;

e = 0;

f = 0;

D and E are a chemical bond;

R' is hydrogen, aralkoxy, or halo;

R'' is lower alkyl, preferably methyl;

Z is -CO<sub>2</sub>H.

35. A compound according to claim 27 wherein:

a = 1;

A is -O-;

b = 0;

c = 0;

5 B is -O-;

d = 1;

e = 0;

f = 0;

D and E are a chemical bond;

10 R' is hydrogen;

R'' is lower alkyl;

Z is -CO<sub>2</sub>H.

36. A compound according to claim 27 wherein:



and



are aryl or heteroaryl;

a = 1;

A is -O-;

b = 0;

c = 0;

20 B is -O-;

d = 1;

e = 0;

f = 0;

D and E are a chemical bond;

25 R' is hydrogen;

R'' is lower alkyl;

Z is -CO<sub>2</sub>H.

37. A compound according to claim 27 wherein:



is optionally substituted azaheteroaryl;



is optionally substituted phenyl;

a = 1;

A is -O-;

b = 0;

c = 0;

B is -O-;

d = 1;

e = 0;

f = 0;

D and E are a chemical bond;

R' is hydrogen;

R'' is lower alkyl;

Z is CO<sub>2</sub>H.

38. A compound according to claim 27 wherein:



is optionally substituted quinolinyl, or a 5-membered heteroaryl group wherein the heteroaryl group is substituted by optionally substituted phenyl or optionally substituted cyclohexyl;



is optionally substituted phenyl;

a = 1;

A is -O-;

b = 0;

c = 0;

B is -O-;

d = 1;

e = 0;

f = 0;


D and E are a chemical bond;

R' is hydrogen;








5 R'' is lower alkyl;

Z is CO<sub>2</sub>H.

39. A compound according to claim 1 selected from the group


















5









;




;




;




;




;




;




;




;




;




;




;




;





;




;




;




;




;




;




;




;




;




;




;



;












;




;





231




oil; CHN calc.  
C30H29NO7+0.5  
H2O: C 68.69, H  
5.76, N 2.67, found  
C 68.68, H 5.71, N  
2.86





5
























40. A compound according to claim 1 selected from the group consisting of




















41. A compound according to claim 1 selected from the group consisting of







;




;




; and




42. A compound according to claim 1 selected from the group consisting of




;




;



;




;




and


43. A compound according to claim 1 selected from the group consisting of




44. A compound according to claim 1 selected from the group consisting of




45. A compound according to claim 1 selected from the group consisting of




46. A compound according to claim 1 selected from the group consisting of



47. A compound according to claim 1 selected from the group consisting of



48. A compound according to claim 1 of the formula



49. A pharmaceutical composition comprising a pharmaceutically acceptable amount of the compound according to claim 1 and a pharmaceutically acceptable carrier.

50. A method of treating a patient suffering from a physiological disorder capable of being modulated by a compound according to claim 1 having PPAR ligand binding activity, comprising administering to the patient a pharmaceutically effective amount of the compound, or a pharmaceutically acceptable salt thereof.

51. A method according to claim 50 wherein the disease is associated with a physiological detrimental blood level of insulin, glucose, free fatty acids (FFA), or triglycerides.

52. The method according to claim 51, wherein the physiological disorder is hyperglycemia.

53. The method according to claim 52, wherein the hyperglycemia is diabetes

54. The method according to claim 52, wherein the hyperglycemia is Type II diabetes.

55. The method according to claim 51, wherein the physiological disorder is

hyperinsulinism.

56. The method according to claim 55, wherein the hyperinsulinism is Syndrome X.

57. The method according to claim 51, wherein the physiological disorder is insulin resistance.

58. The method according to claim 51, wherein the physiological disorder is cardiovascular condition.

59. The method according to claim 58, wherein the cardiovascular condition is atherosclerosis.


60. The method according to claim 51, wherein the physiological disorder is hyperlipidemia.

61. The method according to claim 51, wherein the physiological disorder is hypertension.


62. The method according to claim 51, wherein the physiological disorder is an eating disorder.

63. The method according to claim 50 wherein the mediating is agonistic.


64. The method according to claim 50 wherein the mediating is antagonistic.
65. A method for mediating the activity of PPAR- $\gamma$  receptor comprising contacting said PPAR- $\gamma$  receptor with a compound of according to claim 1.
66. A pharmaceutical composition comprising a pharmaceutically acceptable amount of the compound according to claim 27 and a pharmaceutically acceptable carrier.
67. A method of treating a patient suffering from a physiological disorder capable of being modulated by a compound according to claim 27 having PPAR ligand binding activity, comprising administering to the patient a pharmaceutically effective amount of the compound, or a pharmaceutically acceptable salt thereof.
68. A method according to claim 67 wherein the disease is associated with a physiological detrimental blood level of insulin, glucose, free fatty acids (FFA), or tricyclerides.
69. The method according to claim 67, wherein the physiological disorder is hyperglycemia.
70. The method according to claim 69, wherein the hyperglycemia is diabetes
71. The method according to claim 69, wherein the hyperglycemia is Type II diabetes.
72. The method according to claim 67, wherein the physiological disorder is hyperinsulinism.
73. The method according to claim 72, wherein the hyperinsulinism is Syndrome X.
74. The method according to claim 67, wherein the physiological disorder is insulin resistance.
75. The method according to claim 67, wherein the physiological disorder is cardiovascular disorder.
76. The method according to claim 75, wherein the cardiovascular disorder is atherosclerosis.
77. The method according to claim 67, wherein the physiological disorder is hyperlipidemia.
78. The method according to claim 67, wherein the physiological disorder is hypertension.
79. The method according to claim 67, wherein the physiological disorder is an eating disorder.
80. The method according to claim 67 wherein the mediating is agonistic.
81. The method according to claim 67 wherein the mediating is antagonistic.
82. A method for mediating the activity of PPAR receptor comprising contacting said PPAR receptor with a compound of according to claim 27.
83. A method of treating a patient suffering from a physiological disorder capable of being modulated by a compound having PPAR $\alpha$  and PPAR $\gamma$  ligand binding activity, comprising administering to the patient a pharmaceutically effective amount of the compound, or a pharmaceutically acceptable salt thereof, wherein said compound is of the formula



84. A method of treating a patient suffering from a physiological disorder capable of being modulated by a compound having PPAR $\alpha$  ligand binding activity, comprising administering to the patient a pharmaceutically effective amount of the compound, or a pharmaceutically acceptable salt thereof, wherein said compound is selected from the group consisting of




85. A method of treating a patient suffering from a physiological disorder capable of being modulated by a compound having PPAR $\delta$  ligand binding activity, comprising administering to the patient a pharmaceutically effective amount of the compound, or a pharmaceutically acceptable salt thereof, wherein said compound is of the formula:




86. A method of treating a patient suffering from a physiological disorder capable of being modulated by a compound having PPAR $\alpha$  and PPAR $\delta$  ligand binding activity, comprising administering to the patient a pharmaceutically effective amount of the compound, or a


pharmaceutically acceptable salt thereof, wherein said compound is selected from the group consisting of:



87. A method of treating a patient suffering from a physiological disorder capable of being modulated by a compound having PPAR $\delta$  and PPAR $\gamma$  ligand binding activity, comprising administering to the patient a pharmaceutically effective amount of the compound, or a pharmaceutically acceptable salt thereof, wherein said compound is selected from the group consisting of:




88. A method of treating a patient suffering from a physiological disorder capable of being modulated by a compound having PPAR $\gamma$  ligand binding activity, comprising administering to the patient a pharmaceutically effective amount of the compound, or a pharmaceutically acceptable salt thereof, wherein said compound is selected from the group consisting of:





Abstract

This invention is directed to the use of triaryl acid derivatives of formula (I) and their pharmaceutical compositions as PPAR ligand receptor binders. The PPAR ligand receptor binders of this invention are useful as agonists or antagonists of the PPAR receptor.




wherein:




, and are independently aryl, fused arylcycloalkenyl, fused arylcycloalkyl, fused arylheterocyclenyl, fused arylheterocyclyl, heteroaryl, fused heteroarylalkenyl, fused heteroarylalkyl, fused heteroarylheterocyclenyl, or fused heteroarylheterocyclyl;

A is -O-, -S-, -SO<sub>2</sub>-, -SO<sub>2</sub>-, -NR<sub>13</sub>-, -C(O)-, -N(R<sub>14</sub>)C(O)-, -C(O)N(R<sub>13</sub>)-, -N(R<sub>14</sub>)C(O)N(R<sub>13</sub>)-, -C(R<sub>14</sub>)=N-,



a chemical bond,



B is -O-, -S-, -SO<sub>2</sub>-, -SO<sub>2</sub>-, -NR<sub>17</sub>-, a chemical bond, ethynylene, -C(O)-, -N(R<sub>18</sub>)C(O)-, or -C(O)NR<sub>18</sub>-;

D is -O-, -S-, -NR<sub>19</sub>-, a chemical bond, ethynylene, -C(O)-, -N(R<sub>20</sub>)C(O)-, or -C(O)N(R<sub>20</sub>)-;

E is a chemical bond or an ethylene group;

a is 0-4;

b is 0-4;

c is 0-4;

d is 0-5;

e is 0-4;

f is 0-6;

g is 1-4;

h is 1-4;

$R_1, R_3, R_5, R_7, R_9,$  and  $R_{11}$ , are independently hydrogen, halogen, alkyl, carboxyl, alkoxycarbonyl or aralkyl;

$R_2, R_4, R_6, R_8, R_{10}$  and  $R_{12}$ , are independently  $-(CH_2)_q-X$ ;

$q$  is 0-3;

- 5  $X$  is hydrogen, halogen, alkyl, alkenyl, cycloalkyl, heterocyclyl, aryl, heteroaryl, aralkyl, heteroaralkyl, hydroxy, alkoxy, aralkoxy, heteroaralkoxy, carboxyl, alkoxycarbonyl, tetrazolyl, acyl, acylHNSO<sub>2</sub>-, -SR<sub>23</sub>, Y<sup>1</sup>Y<sup>2</sup>N- or Y<sup>3</sup>Y<sup>4</sup>NCO-;

Y<sup>1</sup> and Y<sup>2</sup> are independently hydrogen, alkyl, aryl, aralkyl or heteroaralkyl, or one of Y<sup>1</sup> and Y<sup>2</sup> is hydrogen or alkyl and the other of Y<sup>1</sup> and Y<sup>2</sup> is acyl or aroyl;

- 10 Y<sup>3</sup> and Y<sup>4</sup> are independently hydrogen, alkyl, aryl, aralkyl or heteroaralkyl;

Z is R<sub>21</sub>O<sub>2</sub>C-, R<sub>21</sub>OC-, cyclo-imide, -CN, R<sub>21</sub>O<sub>2</sub>SHNCO-, R<sub>21</sub>O<sub>2</sub>SHN-, (R<sub>21</sub>)<sub>2</sub>NCO-, R<sub>21</sub>O- 2,4-thiazolidinedionyl, or tetrazolyl; and

R<sub>19</sub> and R<sub>21</sub> are independently hydrogen, alkyl, aryl, cycloalkyl, or aralkyl;

15 R<sub>13</sub>, R<sub>17</sub>, R<sub>19</sub> and R<sub>23</sub> are independently R<sub>22</sub>OC-, R<sub>22</sub>NHOC-, hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, heteroaralkyl, or aralkyl;

R<sub>14</sub>, R<sub>15</sub>, R<sub>16</sub>, R<sub>18</sub> and R<sub>20</sub> are independently hydrogen, alkyl, aralkyl, carbonyl, or alkoxycarbonyl; or R<sub>14</sub>, and R<sub>15</sub> taken together with the carbon and nitrogen atoms through which they are linked form a 5 or 6-membered azaheterocyclyl group; or

20 when a is 2-4, then vicinal R<sub>1</sub> radicals taken together with the carbon atoms to which the R<sub>1</sub> radicals are linked form an ethylene group; or

when b is 2-4, then vicinal R<sub>3</sub> radicals taken together with the carbon atoms to which the R<sub>3</sub> radicals are linked form an ethylene group; or

when c is 2-4, then vicinal R<sub>5</sub> radicals taken together with the carbon atoms to which the R<sub>5</sub> radicals are linked form an ethylene group; or

- 25 when d is 2-5, then vicinal R<sub>7</sub> radicals taken together with the carbon atoms to which the R<sub>7</sub> radicals are linked form an ethylene group; or

when e is 2-4, then vicinal R<sub>9</sub> radicals taken together with the carbon atoms to which the R<sub>9</sub> radicals are linked form an ethylene group; or

when f is 2-6, then vicinal R<sub>11</sub> radicals taken together with the carbon atoms to which the R<sub>11</sub> radicals are linked form an ethylene group; and

30

R<sub>22</sub> is hydrogen, alkyl, aryl, heteroaryl, cycloalkyl, heterocyclyl, heteroaralkyl, or aralkyl; or

a pharmaceutically acceptable salt thereof, an N-oxide thereof, a hydrate thereof or a solvate thereof.

# DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION

As a below named inventor(s), I/We hereby declare that:

My/Our residence(s), post office address(es) and citizenship(s) are as stated below my name(s).  
I/We verily believe I am/we are the original, first and sole/joint inventor(s) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

## THERAPEUTIC USES OF TRI-ARYL ACID DERIVATIVES

and the specification of which ☒ is attached hereto (Attorney Docket No. **A3536A-US**)  
(check one) ☐ was filed on \_\_\_\_\_ as U.S. Application Number \_\_\_\_\_  
☐ and was amended on \_\_\_\_\_ (if applicable).  
☐ was described and claimed in PCT int'l Application Number \_\_\_\_\_  
filed on \_\_\_\_\_ and as amended under PCT Article 19 on \_\_\_\_\_ (if \

I/We hereby state that I/We have reviewed and understand the contents of the above identified specification, including the claims as amended by any amendment referred to above.

I/We acknowledge the duty to disclose to the U.S. Patent and Trademark Office all information known to me/us to be material to patentability as defined in 37 C.F.R. 1.56.

☐ I/We hereby claim foreign priority benefits under Title 35, United States Code §119(a)-(d) or 365 (b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT international application which designated at least one other country other than the United States of America, listed below and having a filing date before that of the application on which priority is claimed. I/We have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Foreign Priority:	Number _____	Country _____	Day/Month/Year Filed _____
Prior Foreign Appln(s):	Number _____	Country _____	Day/Month/Year Filed _____

☒ I/We hereby claim the benefit under Title 35, United States Code §119(e) of any United States Provisional application(s) listed below:

**60/131,454** **28 April 1999 (28.04.99)**  
Number \_\_\_\_\_ Filing Date \_\_\_\_\_

☒ I/We hereby claim the benefit under Title 35, United States Code §120 or 365(c) of any United States application(s) or international application designating the United States listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code §112, I/We acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

**PCT/US00/11490** **28 April 2000** **Pending**  
Application Serial No. \_\_\_\_\_ Filing Date \_\_\_\_\_ Status (Patented, Pending) \_\_\_\_\_

I/We hereby appoint the attorneys and/or agents associated with the Customer No.(s) provided below as my/our attorneys and/or agents with full power to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

**Customer No.: 005487**

I/We hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Title 18, United States code §1001, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Inventors(s): ☐ Additional names and signatures are attached.

1. Full name: Zaid Jayyosi

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Country of Citizenship: France

Residence: US

(City and State/City and Country only)

P. O. Address: 108 Cherrywood Court

Collegeville, PA 19426

3. Full name: Michael F. Kelley

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Country of Citizenship: US

Residence: US

(City and State/City and Country only)

P. O. Address: 1086 Heartsease Drive

West Chester, PA 19382

5. Full name: Litao Zhang

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Country of Citizenship: US

Residence: US

(City and State/City and Country only)

P. O. Address: 456 Shakespere Drive

Collegeville, PA 19426

7. Full name: Daniel G. McGarry

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Country of Citizenship: GB

Residence: US

(City and State/City and Country only)

P. O. Address: 3000 Valley Forge Circle, Apt. 1148

King of Prussia, PA 19406

2. Full name: Gerard M. McGeehan

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Country of Citizenship: US

Residence: US

(City and State/City and Country only)

P. O. Address: 1711 Spring House Road

Chester Springs, PA 19425

4. Full name: Richard F. Labaudiniere

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Country of Citizenship: FR

Residence: US

(City and State/City and Country only)

P. O. Address: 220 Richard Way

Collegeville, PA 19426

6. Full name: Robert D. Groneberg

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Country of Citizenship: US

Residence: US

(City and State/City and Country only)

P. O. Address: 4173 Ironbridge Drive

Collegeville, PA 19426

8. Full name: Thomas J. Caulfield

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Country of Citizenship: : US

Residence: US

(City and State/City and Country only)

P. O. Address: 362 Vista Drive

Phoenixville, PA 19460

9. Full name: Anne Minnich

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Country of Citizenship: US

Residence: US

(City and State/City and Country only)

P. O. Address: 2107 Goodwin Lane

Montgomeryville, PA 19454

10. Full name: Mark Bobko

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Country of Citizenship: US

Residence: US

(City and State/City and Country only)

P. O. Address: 526 Summercroft Drive

Exton, PA 19341

11. Full name: Robert Morris

Signature: \_\_\_\_\_

Date: \_\_\_\_\_

Country of Citizenship: US

Residence: US

(City and State/City and Country only)

P. O. Address: 125 Conestoga Road

Wayne, PA 19087

100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200

Ayentis Pharmaceuticals Products Inc.

Patent Department

Route #202-206

P.O. Box 6800

Mail Code EMC-G1

Bridgewater, NJ 08807-0800

Telephone (908) 231-2985

Telefax (908) 231-2626